

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 49/00	A2	(11) International Publication Number: WO 99/58162 (43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/US99/06826 (22) International Filing Date: 29 March 1999 (29.03.99) (30) Priority Data: 60/080,150 31 March 1998 (31.03.98) US 60/112,715 18 December 1998 (18.12.98) US (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; 974 Centre Road, WR-1ST18, Wilmington, DE 19807 (US). (72) Inventors: RAJOPADHYE, Miland; 21 Honeysuckle Road, Westford, MA 01886 (US). EDWARDS, D., Scott; 123 Farms Drive, Burlington, MA (US). HARRIS, Thomas, D.; 56 Zion Hill Road, Samel, NH 03079 (US). HEMINWAY, Stuart, J.; 21 Fidler Terrace, Lowell, MA 01850 (US). LIU, Shuang; 17 Judith Road, Chelmsford, MA 01824 (US). SINGH, Prahlad, R.; 144 Charlton Street, Arlington, MA 02174 (US). (74) Agent: VANCE, David, H.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SI, SK, UA, VN, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PHARMACEUTICALS FOR THE IMAGING OF ANGIOGENIC DISORDERS (57) Abstract The present invention describes novel compounds of the formula: $(Q)_d-L_n-C_h$, useful for the diagnosis and treatment of cancer, methods of imaging tumors in a patient, and methods of treating cancer in a patient. The present invention also provides novel compounds useful for monitoring therapeutic angiogenesis treatment and destruction of new angiogenic vasculature. The pharmaceuticals are comprised of a targeting moiety that binds to a receptor that is upregulated during angiogenesis, an optional linking group, and a therapeutically effective radioisotope or diagnostically effective imageable moiety. The imageable moiety is a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, an X-ray contrast agent, or an ultrasound contrast agent.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TITLE

PHARMACEUTICALS FOR THE IMAGING OF ANGIOGENIC DISORDERS

5

FIELD OF THE INVENTION

The present invention provides novel pharmaceuticals useful for the diagnosis and treatment of cancer, methods of imaging tumors in a patient, and methods of treating cancer in a patient. The present invention also provides novel
10 pharmaceuticals useful for monitoring therapeutic angiogenesis treatment and destruction of new angiogenic vasculature. The pharmaceuticals are comprised of a targeting moiety that binds to a receptor that is upregulated during angiogenesis, an optional linking group, and a therapeutically effective
15 radioisotope or diagnostically effective imageable moiety. The therapeutically effective radioisotope emits a particle or electron sufficient to be cytotoxic. The imageable moiety is a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, an X-ray contrast agent, or
20 an ultrasound contrast agent.

BACKGROUND OF THE INVENTION

Cancer is a major public health concern in the United States and around the world. It is estimated that over 1
25 million new cases of invasive cancer will be diagnosed in the United States in 1998. The most prevalent forms of the disease are solid tumors of the lung, breast, prostate, colon and rectum. Cancer is typically diagnosed by a combination of in vitro tests and imaging procedures. The imaging procedures
30 include X-ray computed tomography, magnetic resonance imaging, ultrasound imaging and radionuclide scintigraphy. Frequently, a contrast agent is administered to the patient to enhance the image obtained by X-ray CT, MRI and ultrasound, and the administration of a radiopharmaceutical that localizes in
35 tumors is required for radionuclide scintigraphy.

Treatment of cancer typically involves the use of external beam radiation therapy and chemotherapy, either alone or in combination, depending on the type and extent of the

disease. A number of chemotherapeutic agents are available, but generally they all suffer from a lack of specificity for tumors versus normal tissues, resulting in considerable side-effects. The effectiveness of these treatment modalities is also limited, as evidenced by the high mortality rates for a number of cancer types, especially the more prevalent solid tumor diseases. More effective and specific treatment means continue to be needed.

Despite the variety of imaging procedures available for the diagnosis of cancer, there remains a need for improved methods. In particular, methods that can better differentiate between cancer and other pathologic conditions or benign physiologic abnormalities are needed. One means of achieving this desired improvement would be to administer to the patient a metallopharmaceutical that localizes specifically in the tumor by binding to a receptor expressed only in tumors or expressed to a significantly greater extent in tumors than in other tissue. The location of the metallopharmaceutical could then be detected externally either by its imageable emission. in the case of certain radiopharmaceuticals or by its effect on the relaxation rate of water in the immediate vicinity in the case of magnetic resonance imaging contrast agents.

This tumor specific metallopharmaceutical approach can also be used for the treatment of cancer when the metallopharmaceutical is comprised of a particle emitting radioisotope. The radioactive decay of the isotope at the site of the tumor results in sufficient ionizing radiation to be toxic to the tumor cells. The specificity of this approach for tumors minimizes the amount of normal tissue that is exposed to the cytotoxic agent and thus may provide more effective treatment with fewer side-effects.

Previous efforts to achieve these desired improvements in cancer imaging and treatment have centered on the use of radionuclide labeled monoclonal antibodies, antibody fragments and other proteins or polypeptides (i.e., molecular weight over 10,000 D) that bind to tumor cell surface receptors. The specificity of these radiopharmaceuticals is frequently very high, but they suffer from several disadvantages. First,

because of their high molecular weight, they are generally cleared from the blood stream very slowly, resulting in a prolonged blood background in the images. Also, due to their molecular weight they do not extravasate readily at the site of the tumor and then only slowly diffuse through the extravascular space to the tumor cell surface. This results in a very limited amount of the radiopharmaceutical reaching the receptors and thus very low signal intensity in imaging and insufficient cytotoxic effect for treatment.

Alternative approaches to cancer imaging and therapy have involved the use of small molecules, such as peptides, that bind to tumor cell surface receptors. An ^{111}In labeled somatostatin receptor binding peptide, ^{111}In -DTPA-D-Phe¹-octeotide, is in clinical use in many countries for imaging tumors that express the somatostatin receptor (Baker, et al., Life Sci., 1991, 49, 1583-91 and Krenning, et al., Eur. J. Nucl. Med., 1993, 20, 716-31). Higher doses of this radiopharmaceutical have been investigated for potential treatment of these types of cancer (Krenning, et al., Digestion, 1996, 57, 57-61). Several groups are investigating the use of Tc-99m labeled analogs of ^{111}In -DTPA-D-Phe¹-octeotide for imaging and Re-186 labeled analogs for therapy (Flanagan, et al., U.S. 5,556,939, Lyle, et al., U.S. 5,382,654, and Albert et al., U.S. 5,650,134).

Angiogenesis is the process by which new blood vessels are formed from pre-existing capillaries or post capillary venules; it is an important component of a variety of physiological processes including ovulation, embryonic development, wound repair, and collateral vascular generation in the myocardium. It is also central to a number of pathological conditions such as tumor growth and metastasis, diabetic retinopathy, and macular degeneration. The process begins with the activation of existing vascular endothelial cells in response to a variety of cytokines and growth factors. Tumor released cytokines or angiogenic factors stimulate vascular endothelial cells by interacting with specific cell surface receptors for the factors. The activated endothelial cells secrete enzymes that degrade the

basement membrane of the vessels. The endothelial cells then proliferate and invade into the tumor tissue. The endothelial cells differentiate to form lumens, making new vessel offshoots of pre-existing vessels. The new blood vessels then provide nutrients to the tumor permitting further growth and a route for metastasis.

Under normal conditions, endothelial cell proliferation is a very slow process, but it increases for a short period of time during embryogenesis, ovulation and wound healing. This temporary increase in cell turnover is governed by a combination of a number of growth stimulatory factors and growth suppressing factors. In pathological angiogenesis, this normal balance is disrupted resulting in continued increased endothelial cell proliferation. Some of the pro-angiogenic factors that have been identified include basic fibroblast growth factor (bFGF), angiogenin, TGF-alpha, TGF-beta, and vascular endothelium growth factor (VEGF), while interferon-alpha, interferon-beta and thrombospondin are examples of angiogenesis suppressors.

The proliferation and migration of endothelial cells in the extracellular matrix is mediated by interaction with a variety of cell adhesion molecules (Folkman, J., Nature Medicine, 1995, 1, 27-31). Integrins are a diverse family of heterodimeric cell surface receptors by which endothelial cells attach to the extracellular matrix, each other and other cells. The integrin $\alpha_v\beta_3$ is a receptor for a wide variety of extracellular matrix proteins with an exposed tripeptide Arg-Gly-Asp moiety and mediates cellular adhesion to its ligands: vitronectin, fibronectin, and fibrinogen, among others. The integrin $\alpha_v\beta_3$ is minimally expressed on normal blood vessels, but, is significantly upregulated on vascular cells within a variety of human tumors. The role of the $\alpha_v\beta_3$ receptors is to mediate the interaction of the endothelial cells and the extracellular matrix and facilitate the migration of the cells in the direction of the angiogenic signal, the tumor cell population. Angiogenesis induced by bFGF or TNF-alpha depend on the agency of the integrin $\alpha_v\beta_3$, while angiogenesis induced by VEGF depends on the integrin

$\alpha_v\beta_5$ (Cheresh et. al., Science, 1995, 270, 1500-2). Induction of expression of the integrins $\alpha_1\beta_1$ and $\alpha_2\beta_1$ on the endothelial cell surface is another important mechanism by which VEGF promotes angiogenesis (Senger, et. al., Proc. Natl. Acad. Sci USA, 1997, 94, 13612-7).

Angiogenic factors interact with endothelial cell surface receptors such as the receptor tyrosine kinases EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, and Axl. The receptors Flk-1/KDR, neuropilin-1, and Flt-1 recognize VEGF and these interactions play key roles in VEGF-induced angiogenesis. The Tie subfamily of receptor tyrosine kinases are also expressed prominently during blood vessel formation.

Because of the importance of angiogenesis to tumor growth and metastasis, a number of chemotherapeutic approaches are being developed to interfere with or prevent this process. One of these approaches, involves the use of anti-angiogenic proteins such as angiostatin and endostatin. Angiostatin is a 38 kDa fragment of plasminogen that has been shown in animal models to be a potent inhibitor of endothelial cell proliferation. (O'Reilly et. al., Cell, 1994, 79, 315-328) Endostatin is a 20 kDa C-terminal fragment of collagen XVIII that has also been shown to be a potent inhibitor. (O'Reilly et. al., Cell, 1997, 88, 277-285)

Systemic therapy with endostatin has been shown to result in strong anti-tumor activity in animal models. However, human clinical trials of these two chemotherapeutic agents of biological origin have been hampered by lack of availability.

Another approach to anti-angiogenic therapy is to use targeting moieties that interact with endothelial cell surface receptors expressed in the angiogenic vasculature to which are attached chemotherapeutic agents. Burrows and Thorpe (Proc. Nat. Acad. Sci, USA, 1993, 90, 8996-9000) described the use of an antibody-immunotoxin conjugate to eradicate tumors in a mouse model by destroying the tumor vasculature. The antibody was raised against an endothelial cell class II antigen of the major histocompatibility complex and was then conjugated with the cytotoxic agent, deglycosylated ricin A chain. The same

group (Clin. Can. Res., 1995, 1, 1623-1634) investigated the use of antibodies raised against the endothelial cell surface receptor, endoglin, conjugated to deglycosylated ricin A chain. Both of these conjugates exhibited potent anti-tumor activity in mouse models. However, both still suffer drawbacks to routine human use. As with most antibodies or other large, foreign proteins, there is considerable risk of immunologic toxicity which could limit or preclude administration to humans. Also, while the vasculature targeting may improve the local concentration of the attached chemotherapeutic agents, the agents still must be cleaved from the antibody carrier and be transported or diffuse into the cells to be cytotoxic.

Thus, it is desirable to provide anti-angiogenic pharmaceuticals and tumor or new vasculature imaging agents which don't suffer from poor diffusion or transportation, possible immunologic toxicity, limited availability, and/or a lack of specificity.

There is also a growing interest in therapeutic angiogenesis to improve blood flow in regions of the body that have become ischemic or poorly perfused. Several investigators are using growth factors administered locally to cause new vasculature to form either in the limbs or the heart. The growth factors VEGF and bFGF are the most common for this application. Recent publications include: Takeshita, S., et. al., J. Clin. Invest., 1994, 93, 662-670; and Schaper, W. and Schaper, J., Collateral Circulation: Heart, Brain, Kidney, Limbs, Kluwer Academic Publishers, Boston, 1993. The main applications that are under investigation in a number of laboratories are for improving cardiac blood flow and in improving peripheral vessel blood flow in the limbs. For example, Henry, T. et. al. (J. Amer. College Cardiology, 1998, 31, 65A) describe the use of recombinant human VEGF in patients for improving myocardial perfusion by therapeutic angiogenesis. Patients received infusions of rhVEGF and were monitored by nuclear perfusion imaging 30 and 60 days post treatment to determine improvement in myocardial perfusion. About 50% of patients showed improvement by nuclear perfusion

imaging whereas 5/7 showed new collatoralization by angiography.

Thus, it is desirable to discover a method of monitoring improved cardiac blood flow which is targeted to new
5 collatoral vessels themselves and not, as in nuclear perfusion imaging, a regional consequence of new collatoral vessels.

SUMMARY OF THE INVENTION

It is one object of the present invention to provide
10 anti-angiogenic pharmaceuticals, comprised of a targeting moiety that binds to a receptor that is expressed in tumor neovasculature, an optional linking group, and a radioactive metal ion that emits ionizing radiation such as beta particles, alpha particles and Auger or Coster-Kronig
15 electrons. The receptor binding compounds target the radioisotope to the tumor neovasculature. The beta or alpha-particle emitting radioisotope emits a cytotoxic amount of ionizing radiation which results in cell death. The penetrating ability of radiation obviates the requirement that
20 the cytotoxic agent diffuse or be transported into the cell to be cytotoxic.

It is another object of the present invention to provide pharmaceuticals to treat rheumatoid arthritis. These pharmaceuticals comprise a targeting moiety that binds to a
25 receptor that is upregulated during angiogenesis, an optional linking group, and a radioisotope that emits cytotoxic radiation (i.e., beta particles, alpha particles and Auger or Coster-Kronig electrons). In rheumatoid arthritis, the ingrowth of a highly vascularized pannus is caused by the
30 excessive production of angiogenic factors by the infiltrating macrophages, immune cells, or inflammatory cells. Therefore, the radiopharmaceuticals of the present invention that emit cytotoxic radiation could be used to destroy the new angiogenic vasculature that results and thus treat the
35 disease.

It is another object of the present invention to provide tumor imaging agents, comprised of targeting moiety that binds to a receptor that is upregulated during angiogenesis, an

optional linking group, and an imageable moiety, such as a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, an X-ray contrast agent, or an ultrasound contrast agent.

5 It is another object of the present invention to provide imaging agents for monitoring the progress and results of therapeutic angiogenesis treatment. These agents comprise of targeting moiety that binds to a receptor that is upregulated during angiogenesis, an optional linking group, and an
10 imageable moiety. Imaging agents of the present invention could be administered intravenously periodically after the administration of growth factors and imaging would be performed using standard techniques of the affected areas, heart or limbs, to monitor the progress and results of the
15 therapeutic angiogenesis treatment (i.e., image the formation of new blood vessels).

It is another object of the present invention to provide compounds useful for preparing the pharmaceuticals of the present invention. These compounds are comprised of a peptide
20 or peptidomimetic targeting moiety that binds to a receptor that is upregulated during angiogenesis, Q, an optional linking group, L_n , and a metal chelator or bonding moiety, C_h . The compounds may have one or more protecting groups attached to the metal chelator or bonding moiety. The protecting
25 groups provide improved stability to the reagents for long-term storage and are removed either immediately prior to or concurrent with the synthesis of the radiopharmaceuticals. Alternatively, the compounds of the present invention are comprised of a peptide or peptidomimetic targeting moiety that
30 binds to a receptor that is upregulated during angiogenesis, Q, an optional linking group, L_n , and a surfactant, S_f .

The pharmaceuticals of the present invention may be used for diagnostic and/or therapeutic purposes. Diagnostic radiopharmaceuticals of the present invention are
35 pharmaceuticals comprised of a diagnostically useful radionuclide (i.e., a radioactive metal ion that has imageable gamma ray or positron emissions). Therapeutic radiopharmaceuticals of the present invention are

pharmaceuticals comprised of a therapeutically useful radionuclide, a radioactive metal ion that emits ionizing radiation such as beta particles, alpha particles and Auger or Coster-Kronig electrons.

5 The pharmaceuticals comprising a gamma ray or positron emitting radioactive metal ion are useful for imaging tumors by gamma scintigraphy or positron emission tomography. The pharmaceuticals comprising a gamma ray or positron emitting radioactive metal ion are also useful for imaging therapeutic
10 angiogenesis by gamma scintigraphy or positron emission tomography. The pharmaceuticals comprising a particle emitting radioactive metal ion are useful for treating cancer by delivering a cytotoxic dose of radiation to the tumors. The pharmaceuticals comprising a particle emitting radioactive
15 metal ion are also useful for treating rheumatoid arthritis by destroying the formation of angiogenic vasculature. The pharmaceuticals comprising a paramagnetic metal ion are useful as magnetic resonance imaging contrast agents. The pharmaceuticals comprising one or more X-ray absorbing or
20 "heavy" atoms of atomic number 20 or greater are useful as X-ray contrast agents. The pharmaceuticals comprising a microbubble of a biocompatible gas, a liquid carrier, and a surfactant microsphere, are useful as ultrasound contrast agents.

25

DETAILED DESCRIPTION OF THE INVENTION

[1] Thus, in a first embodiment, the present invention provides a novel compound, comprising: a targeting moiety and a chelator, wherein the targeting moiety is bound to the
30 chelator, is a peptide or peptidomimetic, and binds to a receptor that is upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.


35

[2] In a preferred embodiment, the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie,


neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

5

[3] In a more preferred embodiment, the receptor is the integrin $\alpha_v\beta_3$ and the compound is of the formula:



10 wherein, Q is a peptide independently selected from the group:



15 K is an L-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine, δ -N-2-imidazolinylnornithine, δ -N-benzylcarbamoylnornithine, and β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

20

K' is a D-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine, δ -N-2-imidazolinylnornithine, δ -N-benzylcarbamoylnornithine, and β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

25

L is independently selected at each occurrence from the group: glycine, L-alanine, and D-alanine;

30

M is L-aspartic acid;

M' is D-aspartic acid;

R¹ is an amino acid substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, L-valine, D-valine, alanine, leucine,
isoleucine, norleucine, 2-aminobutyric acid,
5 2-aminohexanoic acid, tyrosine, phenylalanine,
thienylalanine, phenylglycine, cyclohexylalanine,
homophenylalanine, 1-naphthylalanine, lysine, serine,
ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
acid, cysteine, penicillamine, and methionine;

10

R² is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, valine, alanine, leucine, isoleucine,
norleucine, 2-aminobutyric acid, 2-aminohexanoic acid,
15 tyrosine, L-phenylalanine, D-phenylalanine,
thienylalanine, phenylglycine, biphenylglycine,
cyclohexylalanine, homophenylalanine,
L-1-naphthylalanine, D-1-naphthylalanine, lysine, serine,
ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
20 acid, cysteine, penicillamine, methionine, and
2-aminothiazole-4-acetic acid;

20

R³ is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
25 glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
D-phenylglycine, D-cyclohexylalanine,
D-homophenylalanine, D-1-naphthylalanine, D-lysine,
30 D-serine, D-ornithine, D-1,2-diaminobutyric acid,
D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
and D-methionine;

30

R⁴ is an amino acid, substituted with 0-1 bonds to L_n,
35 independently selected at each occurrence from the group:
glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
acid, D-tyrosine, D-phenylalanine, D-thienylalanine,

35

D-phenylglycine, D-cyclohexylalanine,
 D-homophenylalanine, D-1-naphthylalanine, D-lysine,
 D-serine, D-ornithine, D-1,2-diaminobutyric acid,
 D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
 5 D-methionine, and 2-aminothiazole-4-acetic acid;

R⁵ is an amino acid, substituted with 0-1 bonds to L_n,
 independently selected at each occurrence from the group:
 glycine, L-valine, L-alanine, L-leucine, L-isoleucine,
 10 L-norleucine, L-2-aminobutyric acid, L-2-aminohexanoic
 acid, L-tyrosine, L-phenylalanine, L-thienylalanine,
 L-phenylglycine, L-cyclohexylalanine,
 L-homophenylalanine, L-1-naphthylalanine, L-lysine,
 L-serine, L-ornithine, L-1,2-diaminobutyric acid,
 15 L-1,2-diaminopropionic acid, L-cysteine, L-penicillamine,
 L-methionine, and 2-aminothiazole-4-acetic acid;

provided that one of R¹, R², R³, R⁴, and R⁵ in each Q is
 substituted with a bond to L_n, further provided that when
 20 R² is 2-aminothiazole-4-acetic acid, K is
 N-methylarginine, further provided that when R⁴ is
 2-aminothiazole-4-acetic acid, K and K' are
 N-methylarginine, and still further provided that when R⁵
 is 2-aminothiazole-4-acetic acid, K' is N-methylarginine;
 25 d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

L_n is a linking group having the formula:

30 $(CR^6R^7)_g - (W)_h - (CR^{6a}R^{7a})_{g'} - (Z)_k - (W)_{h'} - (CR^8R^9)_{g''} - (W)_{h''} - (CR^{8a}R^{9a})_{g'''}$

provided that $g+h+g'+k+h'+g''+h''+g'''$ is other than 0;

W is independently selected at each occurrence from the group:
 35 O, S, NH, NHC(=O), C(=O)NH, C(=O), C(=O)O, OC(=O),
 NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'},
 (OCH₂CH₂CH₂)_{s''}, (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R¹⁰,
C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁰, and a 5-10
5 membered heterocyclic ring system containing 1-4
heteroatoms independently selected from N, S, and O and
substituted with 0-3 R¹⁰;


R⁶, R^{6a}, R⁷, R^{7a}, R⁸, R^{8a}, R⁹ and R^{9a} are independently selected
10 at each occurrence from the group: H, =O, COOH, SO₃H,
PO₃H, C₁₋₅ alkyl substituted with 0-3 R¹⁰, aryl
substituted with 0-3 R¹⁰, benzyl substituted with 0-3
R¹⁰, and C₁₋₅ alkoxy substituted with 0-3 R¹⁰,
NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and a
15 bond to C_H;

R¹⁰ is independently selected at each occurrence from the
group: a bond to C_H, COOR¹¹, OH, NHR¹¹, SO₃H, PO₃H, aryl
substituted with 0-3 R¹¹, C₁₋₅ alkyl substituted with 0-1
20 R¹², C₁₋₅ alkoxy substituted with 0-1 R¹², and a 5-10
membered heterocyclic ring system containing 1-4
heteroatoms independently selected from N, S, and O and
substituted with 0-3 R¹¹;

25 R¹¹ is independently selected at each occurrence from the
group: H, aryl substituted with 0-1 R¹², a 5-10 membered
heterocyclic ring system containing 1-4 heteroatoms
independently selected from N, S, and O and substituted
with 0-1 R¹², C₃₋₁₀ cycloalkyl substituted with 0-1 R¹²,
30 polyalkylene glycol substituted with 0-1 R¹²,
carbohydrate substituted with 0-1 R¹², cyclodextrin
substituted with 0-1 R¹², amino acid substituted with 0-1
R¹², polycarboxyalkyl substituted with 0-1 R¹²,
polyazaalkyl substituted with 0-1 R¹², peptide
35 substituted with 0-1 R¹², wherein the peptide is
comprised of 2-10 amino acids, and a bond to C_H;

R¹² is a bond to C_H;

- k is selected from 0, 1, and 2;
 h is selected from 0, 1, and 2;
 h' is selected from 0, 1, 2, 3, 4, and 5;
 5 h" is selected from 0, 1, 2, 3, 4, and 5;
 g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 g" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 g"' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 10 s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 15 C_h is a metal bonding unit having a formula selected from the group:



- A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently selected
 at each occurrence from the group: N, NR¹³, NR¹³R¹⁴, S,
 SH, S(Pg), O, OH, PR¹³, PR¹³R¹⁴, P(O)R¹⁵R¹⁶, and a bond to
 25 L_n;

E is a bond, CH, or a spacer group independently selected at
 each occurrence from the group: C₁-C₁₀ alkyl substituted
 with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₃₋₁₀

cycloalkyl substituted with 0-3 R¹⁷, heterocyclo-C₁₋₁₀ alkyl substituted with 0-3 R¹⁷, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆₋₁₀ aryl-C₁₋₁₀ alkyl substituted with 0-3 R¹⁷, C₁₋₁₀ alkyl-C₆₋₁₀ aryl- substituted with 0-3 R¹⁷, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;

R¹³, and R¹⁴ are each independently selected from the group: a bond to L_n, hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₁₋₁₀ cycloalkyl substituted with 0-3 R¹⁷, heterocyclo-C₁₋₁₀ alkyl substituted with 0-3 R¹⁷, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆₋₁₀ aryl-C₁₋₁₀ alkyl substituted with 0-3 R¹⁷, C₁₋₁₀ alkyl-C₆₋₁₀ aryl- substituted with 0-3 R¹⁷, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷, and an electron, provided that when one of R¹³ or R¹⁴ is an electron, then the other is also an electron;

alternatively, R¹³ and R¹⁴ combine to form =C(R²⁰)(R²¹);

R¹⁵ and R¹⁶ are each independently selected from the group: a bond to L_n, -OH, C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁷, heterocyclo-C₁₋₁₀ alkyl substituted with 0-3 R¹⁷, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆₋₁₀ aryl-C₁₋₁₀ alkyl substituted with 0-3 R¹⁷, C₁₋₁₀ alkyl-C₆₋₁₀ aryl- substituted with 0-3 R¹⁷, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms

independently selected from N, S, and O and substituted with 0-3 R¹⁷;


R¹⁷ is independently selected at each occurrence from the group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹⁸, -C(=O)R¹⁸, -C(=O)N(R¹⁸)₂, -CHO, -CH₂OR¹⁸, -OC(=O)R¹⁸, -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R¹⁸)₂, -NR¹⁹C(=O)R¹⁸, -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R¹⁸)₂, -NR¹⁹SO₂N(R¹⁸)₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a}, -SR¹⁸, -S(=O)R^{18a}, -SO₂N(R¹⁸)₂, -N(R¹⁸)₂, -NHC(=S)NHR¹⁸, =NOR¹⁸, NO₂, -C(=O)NHOR¹⁸, -C(=O)NHN(R¹⁸)R^{18a}, -OCH₂CO₂H, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆ alkoxyalkyl, aryl substituted with 0-2 R¹⁸, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O;

R¹⁸, R^{18a}, and R¹⁹ are independently selected at each occurrence from the group: a bond to L_n, H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy, halide, nitro, cyano, and trifluoromethyl;

Pg is a thiol protecting group;

R²⁰ and R²¹ are independently selected from the group: H, C₁-C₁₀ alkyl, -CN, -CO₂R²⁵, -C(=O)R²⁵, -C(=O)N(R²⁵)₂, C₂-C₁₀ 1-alkene substituted with 0-3 R²³, C₂-C₁₀ 1-alkyne substituted with 0-3 R²³, aryl substituted with 0-3 R²³, unsaturated 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³, and unsaturated C₃-10 carbocycle substituted with 0-3 R²³;

alternatively, R²⁰ and R²¹, taken together with the divalent carbon radical to which they are attached form:



R^{22} and R^{23} are independently selected from the group: H, R^{24} ,
 C₁-C₁₀ alkyl substituted with 0-3 R^{24} , C₂-C₁₀ alkenyl
 substituted with 0-3 R^{24} , C₂-C₁₀ alkynyl substituted with
 0-3 R^{24} , aryl substituted with 0-3 R^{24} , a 5-10 membered
 heterocyclic ring system containing 1-4 heteroatoms
 independently selected from N, S, and O and substituted
 with 0-3 R^{24} , and C₃-10 carbocycle substituted with 0-3
 R^{24} ;

10

alternatively, R^{22} , R^{23} taken together form a fused aromatic
 or a 5-10 membered heterocyclic ring system containing
 1-4 heteroatoms independently selected from N, S, and O;

15 **a** and **b** indicate the positions of optional double bonds and **n**
 is 0 or 1;

R^{24} is independently selected at each occurrence from the
 group: =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R²⁵, -C(=O)R²⁵,
 -C(=O)N(R²⁵)₂, -N(R²⁵)₃⁺, -CH₂OR²⁵, -OC(=O)R²⁵,
 -OC(=O)OR^{25a}, -OR²⁵, -OC(=O)N(R²⁵)₂, -NR²⁶C(=O)R²⁵,
 -NR²⁶C(=O)OR^{25a}, -NR²⁶C(=O)N(R²⁵)₂, -NR²⁶SO₂N(R²⁵)₂,
 -NR²⁶SO₂R^{25a}, -SO₃H, -SO₂R^{25a}, -SR²⁵, -S(=O)R^{25a},
 -SO₂N(R²⁵)₂, -N(R²⁵)₂, =NOR²⁵, -C(=O)NHOR²⁵, -OCH₂CO₂H,
 and 2-(1-morpholino)ethoxy; and,

25

R^{25} , R^{25a} , and R^{26} are each independently selected at each
 occurrence from the group: hydrogen and C₁-C₆ alkyl;

30 and a pharmaceutically acceptable salt thereof.

[4] In an even more preferred embodiment, the present invention provides a compound, wherein:

L is glycine;

5

R¹ is an amino acid, optionally substituted with a bond to L_n, independently selected at each occurrence from the group: L-valine, D-valine, alanine, leucine, isoleucine, norleucine, 2-aminobutyric acid, tyrosine, phenylalanine, phenylglycine, cyclohexylalanine, homophenylalanine, lysine, ornithine, 1,2-diaminobutyric acid, and 1,2-diaminopropionic acid;

10

R² is an amino acid, optionally substituted with a bond to L_n, independently selected at each occurrence from the group: valine, alanine, leucine, isoleucine, norleucine, 2-aminobutyric acid, tyrosine, L-phenylalanine, D-phenylalanine, thienylalanine, phenylglycine, biphenylglycine, cyclohexylalanine, homophenylalanine, L-1-naphthylalanine, D-1-naphthylalanine, lysine, ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic acid, and 2-aminothiazole-4-acetic acid;

20

R³ is an amino acid, optionally substituted with a bond to L_n, independently selected at each occurrence from the group: D-valine, D-alanine, D-leucine, D-isoleucine, D-norleucine, D-2-aminobutyric acid, D-tyrosine, D-phenylalanine, D-phenylglycine, D-cyclohexylalanine, D-homophenylalanine, D-lysine, D-serine, D-ornithine, D-1,2-diaminobutyric acid, and D-1,2-diaminopropionic acid;

25

30

R⁴ is an amino acid, optionally substituted with a bond to L_n, independently selected at each occurrence from the group: D-valine, D-alanine, D-leucine, D-isoleucine, D-norleucine, D-2-aminobutyric acid, D-tyrosine, D-phenylalanine, D-thienylalanine, D-phenylglycine, D-cyclohexylalanine, D-homophenylalanine,

35

D-1-naphthylalanine, D-lysine, D-ornithine,
D-1,2-diaminobutyric acid, D-1,2-diaminopropionic acid,
and 2-aminothiazole-4-acetic acid;

5 R⁵ is an amino acid, optionally substituted with a bond to L_n,
independently selected at each occurrence from the group:
L-valine, L-alanine, L-leucine, L-isoleucine,
L-norleucine, L-2-aminobutyric acid, L-tyrosine,
L-phenylalanine, L-thienylalanine, L-phenylglycine,
10 L-cyclohexylalanine, L-homophenylalanine,
L-1-naphthylalanine, L-lysine, L-ornithine,
L-1,2-diaminobutyric acid, L-1,2-diaminopropionic acid,
and 2-aminothiazole-4-acetic acid;

15 d is selected from 1, 2, and 3;

W is independently selected at each occurrence from the group:
O, NH, NHC(=O), C(=O)NH, C(=O), C(=O)O, OC(=O),
NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)_s, (CH₂CH₂O)_s,
20 (OCH₂CH₂CH₂)_s, and (CH₂CH₂CH₂O)_t,

Z is selected from the group: aryl substituted with 0-1 R¹⁰,
C₃₋₁₀ cycloalkyl substituted with 0-1 R¹⁰, and a 5-10
membered heterocyclic ring system containing 1-4
25 heteroatoms independently selected from N, S, and O and
substituted with 0-1 R¹⁰;

R⁶, R^{6a}, R⁷, R^{7a}, R⁸, R^{8a}, R⁹, and R^{9a} are independently
selected at each occurrence from the group: H, =O, COOH,
30 SO₃H, C₁-C₅ alkyl substituted with 0-1 R¹⁰, aryl
substituted with 0-1 R¹⁰, benzyl substituted with 0-1
R¹⁰, and C₁-C₅ alkoxy substituted with 0-1 R¹⁰,
NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and a
bond to C_h;

35

R¹⁰ is independently selected at each occurrence from the
group: COOR¹¹, OH, NHR¹¹, SO₃H, aryl substituted with
0-1 R¹¹, a 5-10 membered heterocyclic ring system

containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹¹, C₁-C₅ alkyl substituted with 0-1 R¹², C₁-C₅ alkoxy substituted with 0-1 R¹², and a bond to C_h;

5

R¹¹ is independently selected at each occurrence from the group: H, aryl substituted with 0-1 R¹², a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹², polyalkylene glycol substituted with 0-1 R¹², carbohydrate substituted with 0-1 R¹², cyclodextrin substituted with 0-1 R¹², amino acid substituted with 0-1 R¹², and a bond to C_h;

15 k is 0 or 1;

h is 0 or 1;

h' is 0 or 1;

s is selected from 0, 1, 2, 3, 4, and 5;

s' is selected from 0, 1, 2, 3, 4, and 5;

20 s" is selected from 0, 1, 2, 3, 4, and 5;

t is selected from 0, 1, 2, 3, 4, and 5;

A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently selected at each occurrence from the group: NR¹³, NR¹³R¹⁴, S, SH, S(Pg), OH, and a bond to L_n;

25

E is a bond, CH, or a spacer group independently selected at each occurrence from the group: C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₃-₁₀ cycloalkyl substituted with 0-3 R¹⁷, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;

35 R¹³, and R¹⁴ are each independently selected from the group: a bond to L_n, hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms

independently selected from N, S, and O and substituted with 0-3 R¹⁷, and an electron, provided that when one of R¹³ or R¹⁴ is an electron, then the other is also an electron;

5

alternatively, R¹³ and R¹⁴ combine to form =C(R²⁰)(R²¹);

R¹⁷ is independently selected at each occurrence from the group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹⁸, -C(=O)R¹⁸, -C(=O)N(R¹⁸)₂, -CH₂OR¹⁸, -OC(=O)R¹⁸,
 10 -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R¹⁸)₂, -NR¹⁹C(=O)R¹⁸, -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R¹⁸)₂, -NR¹⁹SO₂N(R¹⁸)₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a}, -S(=O)R^{18a}, -SO₂N(R¹⁸)₂,
 15 -N(R¹⁸)₂, -NHC(=S)NHR¹⁸, =NOR¹⁸, -C(=O)NHN(R¹⁸)R^{18a}, -OCH₂CO₂H, and 2-(1-morpholino)ethoxy;


R¹⁸, R^{18a}, and R¹⁹ are independently selected at each occurrence from the group: a bond to L_n, H, and C₁-C₆ alkyl;

20

R²⁰ and R²¹ are independently selected from the group: H, C₁-C₅ alkyl, -CO₂R²⁵, C₂-C₅ 1-alkene substituted with 0-3 R²³, C₂-C₅ 1-alkyne substituted with 0-3 R²³, aryl substituted with 0-3 R²³, and unsaturated 5-10 membered heterocyclic ring system containing 1-4 heteroatoms
 25 independently selected from N, S, and O and substituted with 0-3 R²³;

alternatively, R²⁰ and R²¹, taken together with the divalent carbon radical to which they are attached form:

30



R²² and R²³ are independently selected from the group: H, and R²⁴;

alternatively, R²², R²³ taken together form a fused aromatic
 5 or a 5-10 membered heterocyclic ring system containing
 1-4 heteroatoms independently selected from N, S, and O;


R²⁴ is independently selected at each occurrence from the
 group: -CO₂R²⁵, -C(=O)N(R²⁵)₂, -CH₂OR²⁵, -OC(=O)R²⁵,
 10 -OR²⁵, -SO₃H, -N(R²⁵)₂, and -OCH₂CO₂H; and,

R²⁵ is independently selected at each occurrence from the
 group: H and C₁-C₃ alkyl.

15

[5] In a still more preferred embodiment, the present
 invention provides a compound, wherein:

Q is a peptide selected from the group:
 20



R¹ is L-valine, D-valine, D-lysine optionally substituted on
 the ε amino group with a bond to L_n or L-lysine
 25 optionally substituted on the ε amino group with a bond
 to L_n;

R² is L-phenylalanine, D-phenylalanine, D-1-naphthylalanine,
 2-aminothiazole-4-acetic acid, L-lysine optionally
 30 substituted on the ε amino group with a bond to L_n or
 tyrosine, the tyrosine optionally substituted on the
 hydroxy group with a bond to L_n;

R³ is D-valine, D-phenylalanine, or L-lysine optionally
 35 substituted on the ε amino group with a bond to L_n;

R^4 is D-phenylalanine, D-tyrosine substituted on the hydroxy group with a bond to L_n , or L-lysine optionally substituted on the ϵ amino group with a bond to L_n ;

5

provided that one of R^1 and R^2 in each Q is substituted with a bond to L_n , and further provided that when R^2 is 2-aminothiazole-4-acetic acid, K is N-methylarginine;

10 d is 1 or 2;

W is independently selected at each occurrence from the group: $NHC(=O)$, $C(=O)NH$, $C(=O)$, $(CH_2CH_2O)_s$, and $(CH_2CH_2CH_2O)_t$;

15 R^6 , R^{6a} , R^7 , R^{7a} , R^8 , R^{8a} , R^9 , and R^{9a} are independently selected at each occurrence from the group: H, $NHC(=O)R^{11}$, and a bond to C_h ;

k is 0;

20 h" is selected from 0, 1, 2, and 3;

g is selected from 0, 1, 2, 3, 4, and 5;


g' is selected from 0, 1, 2, 3, 4, and 5;

g" is selected from 0, 1, 2, 3, 4, and 5;

g''' is selected from 0, 1, 2, 3, 4, and 5;

25 s' is 1 or 2;

t is 1 or 2;



30 A^1 is selected from the group: OH, and a bond to L_n ;

A^2 , A^4 , and A^6 are each N;

A^3 , A^5 , and A^8 are each OH;

35

A^7 is a bond to L_n or NH-bond to L_n ;

E is a C₂ alkyl substituted with 0-1 R¹⁷;

R¹⁷ is =O;

5

alternatively, C_h is A^1-E-A^2 ;

A¹ is NH₂ or N=C(R²⁰)(R²¹);

10 E is a bond;

A² is NHR¹³;

15 R¹³ is a heterocycle substituted with R¹⁷, the heterocycle being selected from pyridine and pyrimidine;


R¹⁷ is selected from a bond to L_n, C(=O)NHR¹⁸, and C(=O)R¹⁸;

R¹⁸ is a bond to L_n;

20

R²⁴ is selected from the group: -CO₂R²⁵, -OR²⁵, -SO₃H, and -N(R²⁵)₂;

25 R²⁵ is independently selected at each occurrence from the group: hydrogen and methyl;



alternatively, C_h is

A¹, A², A³, and A⁴ are each N;

A⁵, A⁶, and A⁸ are each OH;

A⁷ is a bond to L_n;

5

E is a C₂ alkyl substituted with 0-1 R¹⁷; and,

R¹⁷ is =O.

10

[6] In another even more preferred embodiment, the present invention provides a compound selected from the group:

- 15 (a) cyclo{Arg-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 20 (b) cyclo{Arg-Gly-Asp-D-Tyr((N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-18-amino-14-aza-4,7,10-oxy-15-oxo-octadecoyl)-3-aminopropyl)-Val};
- 25 (c) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp};
- 30 (d) cyclo{Arg-Gly-Asp-D-Tyr-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])});
- (e) cyclo{Arg-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])});
- 35 (f) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};

- (g) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Phe-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};
- 5 (h) cyclo{Arg-Gly-Asp-D-Nal-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])});
- (i) [2-[[[5-[carbonyl]-2-pyridinyl]-hydrazono)methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal};
- 10 (j) cyclo{Arg-Gly-Asp-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])-D-Val};
- 15 (k) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-cyclo{Lys-D-Val-Arg-Gly-Asp};
- 20 (l) {cyclo{Arg-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly};
- (m) cyclo{D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])-D-Phe-D-Asp-Gly-Arg};
- 25 (n) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Glu(cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg})-cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg};
- 30 (o) cyclo{D-Phe-D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])-D-Asp-Gly-Arg};
- 35 (p) cyclo{N-Me-Arg-Gly-Asp-ATA-D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])});

- (q) cyclo{Cit-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]]});
- 5 (r) 2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)-1-cyclododecyl)acetyl-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};
- (s) cyclo{Arg-Gly-Asp-D-Phe-Lys(DTPA)};
- 10 (t) cyclo{Arg-Gly-Asp-D-Phe-Lys}₂(DTPA);
- (u) Cyclo{Arg-Gly-Asp-D-Tyr(N-DTPA-3-aminopropyl)-Val};
- 15 (v) cyclo{Orn(d-N-2-Imidazolinyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- (w) cyclo{Lys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 20 (x) cyclo{Cys(2-aminoethyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 25 (y) cyclo{HomoLys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 30 (z) cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- (aa) cyclo{Dap(b-(2-benzimidazolylacetyl))-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 35

- (bb) cyclo{Orn(d-N-2-Imidazoliny1)-Gly-Asp-D-Phe-Lys(N-[2-
[[[5-[carbonyl]-2-pyridiny1]hydrazono)methyl]-
benzenesulfonic acid]]);
- 5 (cc) cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Phe-Lys(N-[2-
[[[5-[carbonyl]-2-pyridiny1]hydrazono)methyl]-
benzenesulfonic acid]]);
- (dd) cyclo{Lys-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-
10 pyridiny1]hydrazono)methyl]-benzenesulfonic acid]-3-
aminopropyl)-D-Asp-Gly};
- (ee) cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(N-[2-[[[5-
[carbonyl]-2-pyridiny1]hydrazono)methyl]-benzenesulfonic
15 acid]-3-aminopropyl)-D-Asp-Gly}; and,
- (ff) cyclo{Orn(d-N-2-Imidazoliny1)-D-Val-D-Tyr(N-[2-[[[5-
[carbonyl]-2-pyridiny1]hydrazono)methyl]-benzenesulfonic
acid]-3-aminopropyl)-D-Asp-Gly};
20
- or a pharmaceutically acceptable salt form thereof.
- [7] In a further preferred embodiment, the present invention
25 provides a kit comprising a compound of the present
invention.
- [8] In an even further preferred embodiment, the kit further
30 comprises one or more ancillary ligands and a reducing
agent.
- [9] In a still further preferred embodiment, the ancillary
35 ligands are tricine and TPPTS.

[10] In another still further preferred embodiment, the reducing agent is tin(II).

5 [11] In a second embodiment, the present invention provides a novel diagnostic or therapeutic metallopharmaceutical composition, comprising: a metal, a chelator capable of chelating the metal and a targeting moiety, wherein the targeting moiety is bound to the chelator, is a peptide or peptidomimetic and binds to a receptor that is
10 upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.

15 [12] In another preferred embodiment, the metallopharmaceutical is a diagnostic radiopharmaceutical, the metal is a radioisotope selected from the group: ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{64}Cu , ^{67}Ga , and ^{68}Ga , the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group:
20 EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

25

[13] In another more preferred embodiment, the targeting moiety is a cyclic pentapeptide and the receptor is $\alpha_v\beta_3$.

30

[14] In another even more preferred embodiment, the radioisotope is ^{99m}Tc or ^{95}Tc , the radiopharmaceutical further comprises a first ancillary ligand and a second ancillary ligand capable of stabilizing the
35 radiopharmaceutical.

[15] In another still more preferred embodiment, the radioisotope is ^{99m}Tc .

5 [16] In another further preferred embodiment, the radiopharmaceutical is selected from the group:

- ^{99m}Tc (tricine)(TPPTS)(cyclo(Arg-Gly-Asp-D-Tyr(N-[[5-[carbonyl]-2-pyridinyl]diazenido]-3-aminopropyl)-Val));
- 10 ^{99m}Tc (tricine)(TPPMS)(cyclo(Arg-D-Val-D-Tyr(N-[[5-[carbonyl]-2-pyridinyl]diazenido]-3-aminopropyl)-D-Asp-Gly));
- ^{99m}Tc (tricine)(TPPDS)(cyclo(Arg-D-Val-D-Tyr(N-[[5-[carbonyl]-2-pyridinyl]diazenido]-3-aminopropyl)-D-Asp-Gly));
- 15 ^{99m}Tc (tricine)(TPPTS)(cyclo(Arg-D-Val-D-Tyr(N-[[5-[carbonyl]-2-pyridinyl]diazenido]-3-aminopropyl)-D-Asp-Gly));
- ^{99m}Tc (tricine)(TPPTS)(cyclo(Arg-Gly-Asp-D-Phe-Lys(N-[[5-[carbonyl]-2-pyridinyl]diazenido]]));
- 20 ^{99m}Tc (tricine)(TPPTS)(cyclo(Arg-Gly-Asp-D-Tyr-Lys(N-[[5-[carbonyl]-2-pyridinyl]diazenido]]));
- 25 ^{99m}Tc (tricine)(TPPTS)((2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid)-Phe-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe});
- 30 ^{99m}Tc (tricine)(TPPTS)(cyclo{Arg-Gly-Asp-D-Nal-Lys}([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]]));
- 35 ^{99m}Tc (tricine)(TPPTS)((2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid)-Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal});

- ^{99m}Tc(tricine) (TPPTS) (cyclo(Arg-Gly-Asp-D-Tyr(N-[5-[carbonyl]-2-pyridinyl]diazenido)-18-amino-14-aza-4,7,10-oxy-15-oxo-octadecoyl)-3-aminopropyl)-Val));
- 5 ^{99m}Tc(tricine) (TPPTS) (N-[5-[carbonyl]-2-pyridinyl]diazenido)-Glu(O-cyclo(Lys-Arg-Gly-Asp-D-Phe))-O-cyclo(Lys-Arg-Gly-Asp-D-Phe));
- 10 ^{99m}Tc(tricine) (TPPTS) (N-[5-[carbonyl]-2-pyridinyl]diazenido)-Glu(O-cyclo(D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp))-O-cyclo(D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp));
- 15 ^{99m}Tc(tricine) (TPPTS) (cyclo(Arg-Gly-Asp-Lys(N-[5-[carbonyl]-2-pyridinyl]diazenido))-D-Val));
- ^{99m}Tc(tricine) (TPPTS) (cyclo(D-Lys([2-[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid))-D-Phe-D-Asp-Gly-Arg));
- 20 ^{99m}Tc(tricine) (TPPTS) ([2-[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid)-Glu(cyclo(D-Lys-D-Phe-D-Asp-Gly-Arg))-cyclo(D-Lys-D-Phe-D-Asp-Gly-Arg));
- 25 ^{99m}Tc(tricine) (TPPTS) (cyclo(D-Phe-D-Lys([2-[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid))-D-Asp-Gly-Arg));
- 30 ^{99m}Tc(tricine) (TPPTS) (cyclo(N-Me-Arg-Gly-Asp-ATA-D-Lys(N-[5-[carbonyl]-2-pyridinyl]diazenido))));
- ^{99m}Tc(tricine) (TPPTS) (cyclo(Cit-Gly-Asp-D-Phe-Lys([2-[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid)))); and,
- 35 ^{99m}Tc(tricine) (1,2,4-triazole) (cyclo(Arg-Gly-Asp-D-Tyr(N-[5-[carbonyl]-2-pyridinyl]diazenido)-3-aminopropyl)-Val)).

[17] In another even more preferred embodiment, the radioisotope is ^{111}In .

5

[18] In another still more preferred embodiment, the radiopharmaceutical is selected from the group:

(DOTA- ^{111}In)-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};

10

cyclo(Arg-Gly-Asp-D-Phe-Lys(DTPA- ^{111}In)); and,

cyclo(Arg-Gly-Asp-D-Phe-Lys) $_2$ (DTPA- ^{111}In).

15

[19] In another preferred embodiment, the metallopharmaceutical is a therapeutic radiopharmaceutical, the metal is a radioisotope selected from the group: ^{186}Re , ^{188}Re , ^{153}Sm , ^{166}Ho , ^{177}Lu , ^{149}Pm , ^{90}Y , ^{212}Bi , ^{103}Pd , ^{109}Pd , ^{159}Gd , ^{140}La , ^{198}Au , ^{199}Au , ^{169}Yb , ^{175}Yb , ^{165}Dy , ^{166}Dy , ^{67}Cu , ^{105}Rh , ^{111}Ag , and ^{192}Ir , the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

20

25

30

[20] In another more preferred embodiment, the targeting moiety is a cyclic pentapeptide and the receptor is $\alpha_v\beta_3$.

35 [21] In another even more preferred embodiment, the radioisotope is ^{153}Sm .

[22] In another still more preferred embodiment, the radiopharmaceutical is selected from the group:

cyclo(Arg-Gly-Asp-D-Phe-Lys(DTPA-¹⁵³Sm));

5

cyclo(Arg-Gly-Asp-D-Phe-Lys)₂(DTPA-¹⁵³Sm); and,

cyclo(Arg-Gly-Asp-D-Tyr(N-DTPA(¹⁵³Sm)-3-aminopropyl)-Val).

10

[23] In another even more preferred embodiment, the radioisotope is ¹⁷⁷Lu.

15 [24] In another still more preferred embodiment, the radiopharmaceutical is selected from the group:

cyclo(Arg-Gly-Asp-D-Phe-Lys(DTPA-¹⁷⁷Lu));

20 (DOTA-¹⁷⁷Lu)-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};

cyclo(Arg-Gly-Asp-D-Phe-Lys)₂(DTPA-¹⁷⁷Lu); and,

25 cyclo(Arg-Gly-Asp-D-Tyr(N-DTPA(¹⁷⁷Lu)-3-aminopropyl)-Val).

[25] In another even more preferred embodiment, the radioisotope is ⁹⁰Y.

30

[26] In another still more preferred embodiment, the radiopharmaceutical is:

35 (DOTA-⁹⁰Y)-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};

[27] In another preferred embodiment, the metallopharmaceutical is a MRI contrast agent, the metal is a paramagnetic metal ion selected from the group: Gd(III), Dy(III), Fe(III), and Mn(II), the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

[28] In another more preferred embodiment, the targeting moiety is a cyclic pentapeptide and the receptor is $\alpha_v\beta_3$.

[29] In another even more preferred embodiment, the metal ion is Gd(III).

[30] In another still more preferred embodiment, the contrast agent is:

cyclo(Arg-Gly-Asp-D-Tyr(N-DTPA(Gd(III)))-3-aminopropyl)-Val).

[31] In another preferred embodiment, the metallopharmaceutical is a X-ray contrast agent, the metal is selected from the group: Re, Sm, Ho, Lu, Pm, Y, Bi, Pd, Gd, La, Au, Au, Yb, Dy, Cu, Rh, Ag, and Ir, the targeting moiety is a cyclic pentapeptide, the receptor is $\alpha_v\beta_3$, and the linking group is present between the targeting moiety and chelator.

[32] In another even more preferred embodiment, the present invention provides a novel method of treating rheumatoid arthritis in a patient comprising: administering a therapeutic

radiopharmaceutical of the present invention capable of localizing in new angiogenic vasculature to a patient by injection or infusion.

5

[33] In another even more preferred embodiment, the present invention provides a novel method of treating cancer in a patient comprising: administering to a patient in need thereof a therapeutic radiopharmaceutical of the present invention by injection or infusion.

[34] In another even more preferred embodiment, the present invention provides a novel method of imaging formation of new blood vessels in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of the present invention to a patient by injection or infusion; (2) imaging the area of the patient wherein the desired formation of new blood vessels is located.

20

[35] In another even more preferred embodiment, the present invention provides a novel method of imaging cancer in a patient comprising: (1) administering a diagnostic radiopharmaceutical of the present invention to a patient by injection or infusion; (2) imaging the patient using planar or SPECT gamma scintigraphy, or positron emission tomography.

[36] In another even more preferred embodiment, the present invention provides a novel method of imaging cancer in a patient comprising: (1) administering a MRI contrast agent of the present invention; and (2) imaging the patient using magnetic resonance imaging.

35

[37] In another even more preferred embodiment, the present invention provides a novel method of imaging cancer in a

patient comprising: (1) administering a X-ray contrast agent of the present invention; and (2) imaging the patient using X-ray computed tomography.

5


[38] In a third embodiment, the present invention provides a novel compound capable of being used in an ultrasound contrast composition, comprising: a targeting moiety and a surfactant, wherein the targeting moiety is bound to the surfactant, is a peptide or peptidomimetic, and binds to a receptor that is upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and surfactant.

[39] In a preferred embodiment, the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and surfactant.

[40] In a more preferred embodiment, the receptor is the integrin $\alpha_v\beta_3$ and the compound is of the formula:

(Q)_d-L_n-S_f

wherein, Q is a cyclic pentapeptide independently selected from the group:



K is an L-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine, δ -N-2-imidazolinylnornithine,

35

δ -N-benzylcarbamoylornithine, and
 β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

5 K' is a D-amino acid independently selected at each occurrence
from the group: arginine, citrulline, N-methylarginine,
lysine, homolysine, 2-aminoethylcysteine,
 δ -N-2-imidazolinylnornithine,
 δ -N-benzylcarbamoylornithine, and
 β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

10

L is independently selected at each occurrence from the group:
glycine, L-alanine, and D-alanine;

M is L-aspartic acid;

15

M' is D-aspartic acid;

R¹ is an amino acid substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
20 glycine, L-valine, D-valine, alanine, leucine,
isoleucine, norleucine, 2-aminobutyric acid,
2-aminohexanoic acid, tyrosine, phenylalanine,
thienylalanine, phenylglycine, cyclohexylalanine,
homophenylalanine, 1-naphthylalanine, lysine, serine,
25 ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
acid, cysteine, penicillamine, and methionine;

R² is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
30 glycine, valine, alanine, leucine, isoleucine,
norleucine, 2-aminobutyric acid, 2-aminohexanoic acid,
tyrosine, L-phenylalanine, D-phenylalanine,
thienylalanine, phenylglycine, biphenylglycine,
cyclohexylalanine, homophenylalanine,
35 L-1-naphthylalanine, D-1-naphthylalanine, lysine, serine,
ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
acid, cysteine, penicillamine, methionine, and
2-aminothiazole-4-acetic acid;

R³ is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
5 D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
D-phenylglycine, D-cyclohexylalanine,
D-homophenylalanine, D-1-naphthylalanine, D-lysine,
D-serine, D-ornithine, D-1,2-diaminobutyric acid,
10 D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
and D-methionine;

R⁴ is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
15 glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
D-phenylglycine, D-cyclohexylalanine,
D-homophenylalanine, D-1-naphthylalanine, D-lysine,
20 D-serine, D-ornithine, D-1,2-diaminobutyric acid,
D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
D-methionine, and 2-aminothiazole-4-acetic acid;

R⁵ is an amino acid, substituted with 0-1 bonds to L_n,
25 independently selected at each occurrence from the group:
glycine, L-valine, L-alanine, L-leucine, L-isoleucine,
L-norleucine, L-2-aminobutyric acid, L-2-aminohexanoic
acid, L-tyrosine, L-phenylalanine, L-thienylalanine,
L-phenylglycine, L-cyclohexylalanine,
30 L-homophenylalanine, L-1-naphthylalanine, L-lysine,
L-serine, L-ornithine, L-1,2-diaminobutyric acid,
L-1,2-diaminopropionic acid, L-cysteine, L-penicillamine,
L-methionine, and 2-aminothiazole-4-acetic acid;

35 provided that one of R¹, R², R³, R⁴, and R⁵ in each Q is
substituted with a bond to L_n, further provided that when
R² is 2-aminothiazole-4-acetic acid, K is
N-methylarginine, further provided that when R⁴ is

2-aminothiazole-4-acetic acid, K and K' are N-methylarginine, and still further provided that when R⁵ is 2-aminothiazole-4-acetic acid, K' is N-methylarginine;

5 d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

S_f is a surfactant which is a lipid or a compound of the

formula: $A^9-E^1-A^{10}$;

10 A⁹ is selected from the group: OH and OR²⁷;

A¹⁰ is OR²⁷;

R²⁷ is C(=O)C₁₋₂₀ alkyl;

15

E¹ is C₁₋₁₀ alkylene substituted with 1-3 R²⁸;

R²⁸ is independently selected at each occurrence from the group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹,

20 -C(=O)N(R²⁹)₂, -CH₂OR²⁹, -OR²⁹, -N(R²⁹)₂, C₁-C₅ alkyl, and C₂-C₄ alkenyl;

R²⁹ is independently selected at each occurrence from the group: R³⁰, H, C₁-C₆ alkyl, phenyl, benzyl, and

25 trifluoromethyl;

R³⁰ is a bond to L_n;

L_n is a linking group having the formula:

30

$(CR^6R^7)_g-(W)_h-(CR^{6a}R^{7a})_g-(Z)_k-(W)_h-(CR^8R^9)_g-(W)_h-(CR^{8a}R^{9a})_g$.

W is independently selected at each occurrence from the group:

35 O, S, NH, NHC(=O), C(=O)NH, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂O)₂₀₋₂₀₀, (OCH₂CH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂CH₂O)₂₀₋₂₀₀, and (aa)_t;

aa is independently at each occurrence an amino acid;

5 Z is selected from the group: aryl substituted with 0-3 R^{10} ,
C₃₋₁₀ cycloalkyl substituted with 0-3 R^{10} , and a 5-10
membered heterocyclic ring system containing 1-4
heteroatoms independently selected from N, S, and O and
substituted with 0-3 R^{10} ;

10 R^6 , R^{6a} , R^7 , R^{7a} , R^8 , R^{8a} , R^9 and R^{9a} are independently selected
at each occurrence from the group: H, =O, COOH, SO₃H,
PO₃H, C₁₋₅ alkyl substituted with 0-3 R^{10} , aryl
substituted with 0-3 R^{10} , benzyl substituted with 0-3
 R^{10} , and C₁₋₅ alkoxy substituted with 0-3 R^{10} ,
15 NHC(=O) R^{11} , C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R^{11} , and a
bond to S_f ;

20 R^{10} is independently selected at each occurrence from the
group: a bond to S_f , COOR¹¹, OH, NHR¹¹, SO₃H, PO₃H, aryl
substituted with 0-3 R^{11} , C₁₋₅ alkyl substituted with 0-1
 R^{12} , C₁₋₅ alkoxy substituted with 0-1 R^{12} , and a 5-10
membered heterocyclic ring system containing 1-4
heteroatoms independently selected from N, S, and O and
substituted with 0-3 R^{11} ;

25 R^{11} is independently selected at each occurrence from the
group: H, aryl substituted with 0-1 R^{12} , a 5-10 membered
heterocyclic ring system containing 1-4 heteroatoms
independently selected from N, S, and O and substituted
with 0-1 R^{12} , C₃₋₁₀ cycloalkyl substituted with 0-1 R^{12} ,
30 amino acid substituted with 0-1 R^{12} , and a bond to S_f ;

R^{12} is a bond to S_f ;


k is selected from 0, 1, and 2;
35 h is selected from 0, 1, and 2;
h' is selected from 0, 1, 2, 3, 4, and 5;
h'' is selected from 0, 1, 2, 3, 4, and 5;
g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 g" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 g"' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

5


and a pharmaceutically acceptable salt thereof.

[41] In another even more preferred embodiment, the compound
 10 is of the formula:



wherein, Q is a cyclic pentapeptide independently selected
 from the group:

15



K is an L-amino acid independently selected at each occurrence
 from the group: arginine, citrulline, N-methylarginine,
 20 lysine, homolysine, 2-aminoethylcysteine,
 δ -N-2-imidazolinylnornithine,
 δ -N-benzylcarbamoylnornithine, and
 β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

25 K' is a D-amino acid independently selected at each occurrence
 from the group: arginine, citrulline, N-methylarginine,
 lysine, homolysine, 2-aminoethylcysteine,
 δ -N-2-imidazolinylnornithine,
 δ -N-benzylcarbamoylnornithine, and
 30 β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

L is independently selected at each occurrence from the group:
 glycine, L-alanine, and D-alanine;

35 M is L-aspartic acid;

M' is D-aspartic acid;

R¹ is an amino acid substituted with 0-1 bonds to L_n,

5 independently selected at each occurrence from the group:
glycine, L-valine, D-valine, alanine, leucine,
isoleucine, norleucine, 2-aminobutyric acid,
2-aminohexanoic acid, tyrosine, phenylalanine,
thienylalanine, phenylglycine, cyclohexylalanine,
10 homophenylalanine, 1-naphthylalanine, lysine, serine,
ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
acid, cysteine, penicillamine, and methionine;

R² is an amino acid, substituted with 0-1 bonds to L_n,

15 independently selected at each occurrence from the group:
glycine, valine, alanine, leucine, isoleucine,
norleucine, 2-aminobutyric acid, 2-aminohexanoic acid,
tyrosine, L-phenylalanine, D-phenylalanine,
thienylalanine, phenylglycine, biphenylglycine,
20 cyclohexylalanine, homophenylalanine,
L-1-naphthylalanine, D-1-naphthylalanine, lysine, serine,
ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
acid, cysteine, penicillamine, methionine, and
2-aminothiazole-4-acetic acid;

25 R³ is an amino acid, substituted with 0-1 bonds to L_n,

independently selected at each occurrence from the group:
glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
30 acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
D-phenylglycine, D-cyclohexylalanine,
D-homophenylalanine, D-1-naphthylalanine, D-lysine,
D-serine, D-ornithine, D-1,2-diaminobutyric acid,
D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
35 and D-methionine;

R⁴ is an amino acid, substituted with 0-1 bonds to L_n,

independently selected at each occurrence from the group:

glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
 D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
 acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
 D-phenylglycine, D-cyclohexylalanine,
 5 D-homophenylalanine, D-1-naphthylalanine, D-lysine,
 D-serine, D-ornithine, D-1,2-diaminobutyric acid,
 D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
 D-methionine, and 2-aminothiazole-4-acetic acid;

10 R⁵ is an amino acid, substituted with 0-1 bonds to L_n,
 independently selected at each occurrence from the group:
 glycine, L-valine, L-alanine, L-leucine, L-isoleucine,
 L-norleucine, L-2-aminobutyric acid, L-2-aminohexanoic
 acid, L-tyrosine, L-phenylalanine, L-thienylalanine,
 15 L-phenylglycine, L-cyclohexylalanine,
 L-homophenylalanine, L-1-naphthylalanine, L-lysine,
 L-serine, L-ornithine, L-1,2-diaminobutyric acid,
 L-1,2-diaminopropionic acid, L-cysteine, L-penicillamine,
 L-methionine, and 2-aminothiazole-4-acetic acid;

20 provided that one of R¹, R², R³, R⁴, and R⁵ in each Q is
 substituted with a bond to L_n, further provided that when
 R² is 2-aminothiazole-4-acetic acid, K is
 N-methylarginine, further provided that when R⁴ is
 25 2-aminothiazole-4-acetic acid, K and K' are
 N-methylarginine, and still further provided that when R⁵
 is 2-aminothiazole-4-acetic acid, K' is N-methylarginine;

S_f is a surfactant which is a lipid or a compound of the

30 formula: $A^9 - E^L - A^{10}$;

A⁹ is OR²⁷;

A¹⁰ is OR²⁷;

35 R²⁷ is C(=O)C₁₋₁₅ alkyl;

E¹ is C₁₋₄ alkylene substituted with 1-3 R²⁸;

R²⁸ is independently selected at each occurrence from the
group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹, -CH₂OR²⁹,
5 -OR²⁹, and C₁₋₅ alkyl;

R²⁹ is independently selected at each occurrence from the
group: R³⁰, H, C₁₋₆ alkyl, phenyl, and benzyl;

10 R³⁰ is a bond to L_n;

L_n is a linking group having the formula:

(CR⁶R⁷)_g-(W)_h-(CR^{6a}R^{7a})_{g'}-(Z)_k-(W)_{h'}-(CR⁸R⁹)_{g''}-(W)_{h''}-(CR^{8a}R^{9a})_{g'''}.
15

W is independently selected at each occurrence from the group:
O, S, NH, NHC(=O), C(=O)NH, C(=O), C(=O)O, OC(=O),
NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)₂₀₋₂₀₀,
(CH₂CH₂O)₂₀₋₂₀₀, (OCH₂CH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂CH₂O)₂₀₋₂₀₀, and
20 (aa)_t;

aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R¹⁰,
25 C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁰, and a 5-10
membered heterocyclic ring system containing 1-4
heteroatoms independently selected from N, S, and O and
substituted with 0-3 R¹⁰;

30 R⁶, R^{6a}, R⁷, R^{7a}, R⁸, R^{8a}, R⁹ and R^{9a} are independently selected
at each occurrence from the group: H, =O, C₁₋₅ alkyl
substituted with 0-3 R¹⁰, and C₁₋₅ alkoxy substituted
with 0-3 R¹⁰, and a bond to S_f;

35 R¹⁰ is independently selected at each occurrence from the
group: a bond to S_f, COOR¹¹, OH, NHR¹¹, C₁₋₅ alkyl
substituted with 0-1 R¹², and C₁₋₅ alkoxy substituted
with 0-1 R¹²;

R¹¹ is independently selected at each occurrence from the
group: H, aryl substituted with 0-1 R¹², C₃₋₁₀ cycloalkyl
substituted with 0-1 R¹², amino acid substituted with 0-1
5 R¹², and a bond to S_f;

R¹² is a bond to S_f;

k is selected from 0, 1, and 2;
10 h is selected from 0, 1, and 2;
h' is selected from 0, 1, 2, 3, 4, and 5;
h'' is selected from 0, 1, 2, 3, 4, and 5;
g is selected from 0, 1, 2, 3, 4, and 5;
g' is selected from 0, 1, 2, 3, 4, and 5;
15 g'' is selected from 0, 1, 2, 3, 4, and 5;
g''' is selected from 0, 1, 2, 3, 4, and 5;
s is selected from 0, 1, 2, 3, 4, and 5;
s' is selected from 0, 1, 2, 3, 4, and 5;
s'' is selected from 0, 1, 2, 3, 4, and 5;
20 t is selected from 0, 1, 2, 3, 4, and 5;
t' is selected from 0, 1, 2, 3, 4, and 5;

and a pharmaceutically acceptable salt thereof.

25

[42] In another still more preferred embodiment, the present
invention provides a compound selected from the group:

1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-
30 (cyclo(Arg-Gly-Asp-D-Phe-Lys))-dodecane-1,12-dione;

1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-((ω-
amino-PEG₃₄₀₀-α-carbonyl)-cyclo(Arg-Gly-Asp-D-Phe-Lys))-
dodecane-1,12-dione; and,

35

1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-((ω-
amino-PEG₃₄₀₀-α-carbonyl)-Glu-(cyclo(Arg-Gly-Asp-D-Phe-
Lys))₂)-Dodecane-1,12-dione.

[43] In another even more preferred embodiment, the present invention provides a novel ultrasound contrast agent composition, comprising:

- (a) a compound comprising: a cyclic pentapeptide that binds to the integrin $\alpha_v\beta_3$, a surfactant and a linking group between the cyclicpentapeptide and the surfactant;
- (b) a parenterally acceptable carrier; and,
- (c) an echogenic gas.

[44] In another still more preferred embodiment, the ultrasound contrast agent further comprises: 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine.

[45] In another further preferred embodiment, the echogenic gas is a C₂₋₅ perfluorocarbon.

[46] In another even more preferred embodiment, the present invention provides a method of imaging cancer in a patient comprising: (1) administering, by injection or infusion, a ultrasound contrast agent composition of the present invention to a patient; and (2) imaging the patient using sonography.

[47] In another even more preferred embodiment, the present invention provides a novel method of imaging formation of new blood vessels in a patient comprising: (1) administering, by injection or infusion, a ultrasound contrast agent composition of the present invention to a patient; (2) imaging the area of the patient wherein the desired formation of new blood vessels is located.

[48] In another even more preferred embodiment, the present invention provides a novel therapeutic radiopharmaceutical composition, comprising:

- (a) a therapeutic radiopharmaceutical of the present invention; and,
- (b) a parenterally acceptable carrier.

10

[49] In another even more preferred embodiment, the present invention provides a novel diagnostic radiopharmaceutical composition, comprising:

- (a) a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of the present invention; and,
- (b) a parenterally acceptable carrier.

20 [50] In another even more preferred embodiment, the present invention provides a novel therapeutic radiopharmaceutical composition, comprising: a radiolabelled targeting moiety, wherein the targeting moiety is a compound Q and the radiolabel is a therapeutic isotope selected from the group:
25 ^{35}S , ^{32}P , ^{125}I , ^{131}I , and ^{211}At .

[51] In another further preferred embodiment, the present invention provides a novel therapeutic radiopharmaceutical composition, comprising: a radiolabelled targeting moiety, wherein the targeting moiety is a compound Q and the radiolabel is a therapeutic isotope which is ^{131}I .

35 Another embodiment of the present invention is diagnostic kits for the preparation of radiopharmaceuticals useful as imaging agents for cancer or imaging agents for imaging formation of new blood vessels. Diagnostic kits of the

present invention comprise one or more vials containing the sterile, non-pyrogenic, formulation comprised of a predetermined amount of a reagent of the present invention, and optionally other components such as one or two ancillary
5 ligands, reducing agents, transfer ligands, buffers, lyophilization aids, stabilization aids, solubilization aids and bacteriostats. The inclusion of one or more optional components in the formulation will frequently improve the ease of synthesis of the radiopharmaceutical by the practicing end
10 user, the ease of manufacturing the kit, the shelf-life of the kit, or the stability and shelf-life of the radiopharmaceutical. The inclusion of one or two ancillary ligands is required for diagnostic kits comprising reagent comprising a hydrazine or hydrazone bonding moiety. The one
15 or more vials that contain all or part of the formulation can independently be in the form of a sterile solution or a lyophilized solid.

DEFINITIONS

20 The compounds herein described may have asymmetric centers. Unless otherwise indicated, all chiral, diastereomeric and racemic forms are included in the present invention. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds
25 described herein, and all such stable isomers are contemplated in the present invention. It will be appreciated that compounds of the present invention contain asymmetrically substituted carbon atoms, and may be isolated in optically active or racemic forms. It is well known in the art how to
30 prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Two distinct isomers (cis and trans) of the peptide bond are known to occur; both can also be present in the compounds described herein, and all such stable isomers
35 are contemplated in the present invention. The D and L-isomers of a particular amino acid are designated herein using the conventional 3-letter abbreviation of the amino

acid, as indicated by the following examples: D-Leu, or L-Leu.

When any variable occurs more than one time in any substituent or in any formula, its definition on each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R^{52} , then said group may optionally be substituted with up to two R^{52} , and R^{52} at each occurrence is selected independently from the defined list of possible R^{52} . Also, by way of example, for the group $-N(R^{53})_2$, each of the two R^{53} substituents on N is independently selected from the defined list of possible R^{53} . Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. When a bond to a substituent is shown to cross the bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring.

By "reagent" is meant a compound of this invention capable of direct transformation into a metallopharmaceutical of this invention. Reagents may be utilized directly for the preparation of the metallopharmaceuticals of this invention or may be a component in a kit of this invention.

The term "binding agent" means a metallopharmaceutical of this invention having affinity for and capable of binding to the vitronectin receptor. The binding agents of this invention preferably have $K_i < 1000 \text{ nM}$.

Metallopharmaceutical as used herein is intended to refer to a pharmaceutically acceptable compound containing a metal, wherein the compound is useful for imaging, magnetic resonance imaging, contrast imaging, or x-ray imaging. The metal is the cause of the imageable signal in diagnostic applications and the source of the cytotoxic radiation in radiotherapeutic applications. Radiopharmaceuticals are metallopharmaceuticals in which the metal is a radioisotope.

By "stable compound" or "stable structure" is meant herein a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious pharmaceutical agent.

The term "substituted", as used herein, means that one or more hydrogens on the designated atom or group is replaced with a selection from the indicated group, provided that the designated atom's or group's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

The term "bond", as used herein, means either a single or double bond.

The term "salt", as used herein, is used as defined in the CRC Handbook of Chemistry and Physics, 65th Edition, CRC Press, Boca Raton, Fla, 1984, as any substance which yields ions, other than hydrogen or hydroxyl ions. As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds modified by making acid or base salts. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts

include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

10 The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. C₁₋₁₀ alkyl, is intended to include C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, and C₁₀ alkyl groups. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, and s-pentyl. "Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example -C_vF_w where v = 1 to 3 and w = 1 to (2v+1)). Examples of haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, and pentachloroethyl. "Alkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. C₁₋₁₀ alkoxy, is intended to include C₁, C₂, C₃, C₄,

C₅, C₆, C₇, C₈, C₉, and C₁₀ alkoxy groups. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxy, and s-pentoxy. "Cycloalkyl" is intended to include saturated ring groups, such as cyclopropyl, cyclobutyl, or cyclopentyl. C₃₋₇ cycloalkyl, is intended to include C₃, C₄, C₅, C₆, and C₇ cycloalkyl groups. "Alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl and propenyl. C₂₋₁₀ alkenyl, is intended to include C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, and C₁₀ alkenyl groups. "Alkynyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more triple carbon-carbon bonds which may occur in any stable point along the chain, such as ethynyl and propynyl. C₂₋₁₀ alkynyl, is intended to include C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, and C₁₀ alkynyl groups.

As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3, 4, 5, 6, or 7-membered monocyclic or bicyclic or 7, 8, 9, 10, 11, 12, or 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane, [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl.

As used herein, the term "alkaryl" means an aryl group bearing an alkyl group of 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms; the term "aralkyl" means an alkyl group of 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms bearing an aryl group; the term "arylalkaryl" means an aryl group bearing an alkyl group of 1-10 carbon atoms bearing an aryl group; and the term "heterocycloalkyl" means an alkyl group of 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms bearing a heterocycle.

As used herein, the term "heterocycle" or "heterocyclic system" is intended to mean a stable 5, 6, or 7-membered

monocyclic or bicyclic or 7, 8, 9, or 10-membered bicyclic heterocyclic ring which is saturated, partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and 1, 2, 3, or 4 heteroatoms independently selected from the group consisting of N, NH, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" or "heteroaryl" is intended to mean a stable 5, 6, or 7-membered monocyclic or bicyclic or 7, 8, 9, or 10-membered bicyclic heterocyclic aromatic ring which consists of carbon atoms and 1, 2, 3, or 4 heterotams independently selected from the group consisting of N, NH, O and S. It is to be noted that total number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl, indolenyl, indolinyl, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, methylenedioxyphenyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-

oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,
oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl,
phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl,
phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl,
5 piperidinyl, piperidonyl, 4-piperidonyl, piperonyl,
pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl,
pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole,
pyridoimidazole, pyridothiazole, pyridinyl, pyridyl,
pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl,
10 quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxalinyl,
quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl,
tetrahydroquinolinyl, tetrazolyl, 6H-1,2,5-thiadiazinyl,
1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl,
1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl,
15 thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl,
triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl,
1,3,4-triazolyl, and xanthenyl. Preferred heterocycles
include, but are not limited to, pyridinyl, furanyl, thienyl,
pyrrolyl, pyrazolyl, pyrrolidinyl, imidazolyl, indolyl,
20 benzimidazolyl, 1H-indazolyl, oxazolidinyl, benzotriazolyl,
benzisoxazolyl, oxindolyl, benzoxazoliny, and isatinoyl.
Also included are fused ring and spiro compounds containing,
for example, the above heterocycles.

A "polyalkylene glycol" is a polyethylene glycol,
25 polypropylene glycol or polybutylene glycol having a molecular
weight of less than about 5000, terminating in either a
hydroxy or alkyl ether moiety.

A "carbohydrate" is a polyhydroxy aldehyde, ketone,
alcohol or acid, or derivatives thereof, including polymers
30 thereof having polymeric linkages of the acetal type.

A "cyclodextrin" is a cyclic oligosaccharide. Examples
of cyclodextrins include, but are not limited to,
 α -cyclodextrin, hydroxyethyl- α -cyclodextrin,
hydroxypropyl- α -cyclodextrin, β -cyclodextrin,
35 hydroxypropyl- β -cyclodextrin, carboxymethyl- β -cyclodextrin,
dihydroxypropyl- β -cyclodextrin, hydroxyethyl- β -cyclodextrin,
2,6 di-O-methyl- β -cyclodextrin, sulfated- β -cyclodextrin,
 γ -cyclodextrin, hydroxypropyl- γ -cyclodextrin,

dihydroxypropyl- γ -cyclodextrin, hydroxyethyl- γ -cyclodextrin, and sulfated γ -cyclodextrin.

As used herein, the term "polycarboxyalkyl" means an alkyl group having between two and about 100 carbon atoms and a plurality of carboxyl substituents; and the term
5 "polyazaalkyl" means a linear or branched alkyl group having between two and about 100 carbon atoms, interrupted by or substituted with a plurality of amine groups.

A "reducing agent" is a compound that reacts with a
10 radionuclide, which is typically obtained as a relatively unreactive, high oxidation state compound, to lower its oxidation state by transferring electron(s) to the radionuclide, thereby making it more reactive. Reducing agents useful in the preparation of radiopharmaceuticals and
15 in diagnostic kits useful for the preparation of said radiopharmaceuticals include but are not limited to stannous chloride, stannous fluoride, formamidine sulfinic acid, ascorbic acid, cysteine, phosphines, and cuprous or ferrous salts. Other reducing agents are described in Brodack et.
20 al., PCT Application 94/22496, which is incorporated herein by reference.

A "transfer ligand" is a ligand that forms an intermediate complex with a metal ion that is stable enough to prevent unwanted side-reactions but labile enough to be
25 converted to a metallopharmaceutical. The formation of the intermediate complex is kinetically favored while the formation of the metallopharmaceutical is thermodynamically favored. Transfer ligands useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the
30 preparation of diagnostic radiopharmaceuticals include but are not limited to gluconate, glucoheptonate, mannitol, glucarate, N,N,N',N'-ethylenediaminetetraacetic acid, pyrophosphate and methylenediphosphonate. In general, transfer ligands are comprised of oxygen or nitrogen donor atoms.

35 The term "donor atom" refers to the atom directly attached to a metal by a chemical bond.

"Ancillary" or "co-ligands" are ligands that are incorporated into a radiopharmaceutical during its synthesis.

They serve to complete the coordination sphere of the radionuclide together with the chelator or radionuclide bonding unit of the reagent. For radiopharmaceuticals comprised of a binary ligand system, the radionuclide coordination sphere is composed of one or more chelators or bonding units from one or more reagents and one or more ancillary or co-ligands, provided that there are a total of two types of ligands, chelators or bonding units. For example, a radiopharmaceutical comprised of one chelator or bonding unit from one reagent and two of the same ancillary or co-ligands and a radiopharmaceutical comprised of two chelators or bonding units from one or two reagents and one ancillary or co-ligand are both considered to be comprised of binary ligand systems. For radiopharmaceuticals comprised of a ternary ligand system, the radionuclide coordination sphere is composed of one or more chelators or bonding units from one or more reagents and one or more of two different types of ancillary or co-ligands, provided that there are a total of three types of ligands, chelators or bonding units. For example, a radiopharmaceutical comprised of one chelator or bonding unit from one reagent and two different ancillary or co-ligands is considered to be comprised of a ternary ligand system.

Ancillary or co-ligands useful in the preparation of radiopharmaceuticals and in diagnostic kits useful for the preparation of said radiopharmaceuticals are comprised of one or more oxygen, nitrogen, carbon, sulfur, phosphorus, arsenic, selenium, and tellurium donor atoms. A ligand can be a transfer ligand in the synthesis of a radiopharmaceutical and also serve as an ancillary or co-ligand in another radiopharmaceutical. Whether a ligand is termed a transfer or ancillary or co-ligand depends on whether the ligand remains in the radionuclide coordination sphere in the radiopharmaceutical, which is determined by the coordination chemistry of the radionuclide and the chelator or bonding unit of the reagent or reagents.

A "chelator" or "bonding unit" is the moiety or group on a reagent that binds to a metal ion through the formation of chemical bonds with one or more donor atoms.

The term "binding site" means the site in vivo or in vitro that binds a biologically active molecule.

A "diagnostic kit" or "kit" comprises a collection of components, termed the formulation, in one or more vials which are used by the practicing end user in a clinical or pharmacy setting to synthesize diagnostic radiopharmaceuticals. The kit provides all the requisite components to synthesize and use the diagnostic radiopharmaceutical except those that are commonly available to the practicing end user, such as water or saline for injection, a solution of the radionuclide, equipment for heating the kit during the synthesis of the radiopharmaceutical, if required, equipment necessary for administering the radiopharmaceutical to the patient such as syringes and shielding, and imaging equipment.

Therapeutic radiopharmaceuticals, X-ray contrast agent pharmaceuticals, ultrasound contrast agent pharmaceuticals and metallopharmaceuticals for magnetic resonance imaging contrast are provided to the end user in their final form in a formulation contained typically in one vial, as either a lyophilized solid or an aqueous solution. The end user reconstitutes the lyophilized with water or saline and withdraws the patient dose or just withdraws the dose from the aqueous solution formulation as provided.

A "lyophilization aid" is a component that has favorable physical properties for lyophilization, such as the glass transition temperature, and is added to the formulation to improve the physical properties of the combination of all the components of the formulation for lyophilization.

A "stabilization aid" is a component that is added to the metallopharmaceutical or to the diagnostic kit either to stabilize the metallopharmaceutical or to prolong the shelf-life of the kit before it must be used. Stabilization aids can be antioxidants, reducing agents or radical scavengers and can provide improved stability by reacting

preferentially with species that degrade other components or the metallopharmaceutical.

A "solubilization aid" is a component that improves the solubility of one or more other components in the medium
5 required for the formulation.

A "bacteriostat" is a component that inhibits the growth of bacteria in a formulation either during its storage before use of after a diagnostic kit is used to synthesize a radiopharmaceutical.

10 The term "amino acid" as used herein means an organic compound containing both a basic amino group and an acidic carboxyl group. Included within this term are natural amino acids (e.g., L-amino acids), modified and unusual amino acids (e.g., D-amino acids), as well as amino acids which are known
15 to occur biologically in free or combined form but usually do not occur in proteins. Included within this term are modified and unusual amino acids, such as those disclosed in, for example, Roberts and Vellaccio (1983) The Peptides, 5: 342-429, the teaching of which is hereby incorporated by
20 reference. Natural protein occurring amino acids include, but are not limited to, alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, serine, threonine, tyrosine, tryptophan, proline,
25 and valine. Natural non-protein amino acids include, but are not limited to arginosuccinic acid, citrulline, cysteine sulfinic acid, 3,4-dihydroxyphenylalanine, homocysteine, homoserine, ornithine, 3-monoiodotyrosine, 3,5-diiodotyrosine, 3,5,5'-triiodothyronine, and 3,3',5,5'-tetraiodothyronine.
30 Modified or unusual amino acids which can be used to practice the invention include, but are not limited to, D-amino acids, hydroxylysine, 4-hydroxyproline, an N-Cbz-protected amino acid, 2,4-diaminobutyric acid, homoarginine, norleucine, N-methylaminobutyric acid, naphthylalanine, phenylglycine,
35 β -phenylproline, tert-leucine, 4-aminocyclohexylalanine, N-methyl-norleucine, 3,4-dehydroproline, N,N-dimethylaminoglycine, N-methylaminoglycine, 4-aminopiperidine-4-carboxylic acid, 6-aminocaproic acid,

trans-4-(aminomethyl)-cyclohexanecarboxylic acid, 2-, 3-, and 4-(aminomethyl)-benzoic acid, 1-aminocyclopentanecarboxylic acid, 1-aminocyclopropanecarboxylic acid, and 2-benzyl-5-aminopentanoic acid.

5 The term "peptide" as used herein means a linear compound that consists of two or more amino acids (as defined herein) that are linked by means of a peptide bond. A "peptide" as used in the presently claimed invention is intended to refer to a moiety with a molecular weight of less than 10,000
10 Daltons, preferable less than 5,000 Daltons, and more preferably less than 2,500 Daltons. The term "peptide" also includes compounds containing both peptide and non-peptide components, such as pseudopeptide or peptidomimetic residues or other non-amino acid components. Such a compound
15 containing both peptide and non-peptide components may also be referred to as a "peptide analog".

A "pseudopeptide" or "peptidomimetic" is a compound which mimics the structure of an amino acid residue or a peptide, for example, by using linking groups other than amide linkages
20 between the peptide mimetic and an amino acid residue (pseudopeptide bonds) and/or by using non-amino acid substituents and/or a modified amino acid residue. A "pseudopeptide residue" means that portion of an pseudopeptide or peptidomimetic that is present in a peptide.

25 The term "peptide bond" means a covalent amide linkage formed by loss of a molecule of water between the carboxyl group of one amino acid and the amino group of a second amino acid.

The term "pseudopeptide bonds" includes peptide bond
30 isosteres which may be used in place of or as substitutes for the normal amide linkage. These substitute or amide "equivalent" linkages are formed from combinations of atoms not normally found in peptides or proteins which mimic the spatial requirements of the amide bond and which should
35 stabilize the molecule to enzymatic degradation.

The following abbreviations are used herein:

Acm	acetamidomethyl
-----	-----------------

	b-Ala, beta-Ala	
	or bAla	3-aminopropionic acid
	ATA	2-aminothiazole-5-acetic acid or 2-aminothiazole-5-acetyl group
5	Boc	t-butyloxycarbonyl
	CBZ, Cbz or Z	Carbobenzyloxy
	Cit	citrulline
	Dap	2,3-diaminopropionic acid
	DCC	dicyclohexylcarbodiimide
10	DIEA	diisopropylethylamine
	DMAP	4-dimethylaminopyridine
	EOE	ethoxyethyl
	HBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
15	hynic	boc-hydrazinonicotinyl group or 2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid,
	NMeArg or MeArg	a-N-methyl arginine
	NMeAsp	a-N-methyl aspartic acid
20	NMM	N-methylmorpholine
	OcHex	O-cyclohexyl
	OBzl	O-benzyl
	oSu	O-succinimidyl
25	TBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
	THF	tetrahydrofuran
	THP	tetrahydropyran
	Tos	tosyl
	Tr	trityl

30

The following conventional three-letter amino acid abbreviations are used herein; the conventional one-letter amino acid abbreviations are NOT used herein:

35	Ala	=	alanine
	Arg	=	arginine
	Asn	=	asparagine
	Asp	=	aspartic acid

	Cys	=	cysteine
	Gln	=	glutamine
	Glu	=	glutamic acid
	Gly	=	glycine
5	His	=	histidine
	Ile	=	isoleucine
	Leu	=	leucine
	Lys	=	lysine
	Met	=	methionine
10	Nle	=	norleucine
	Orn	=	ornithine
	Phe	=	phenylalanine
	Phg	=	phenylglycine
	Pro	=	proline
15	Sar	=	sarcosine
	Ser	=	serine
	Thr	=	threonine
	Trp	=	tryptophan
	Tyr	=	tyrosine
20	Val	=	valine

The pharmaceuticals of the present invention are comprised of a targeting moiety for a receptor that is expressed or upregulated in angiogenic tumor vasculature. For targeting the VEGF receptors, Flk-1/KDR, Flt-1, and neuropilin-1, the targeting moieties are comprised of peptides or peptidomimetics that bind with high affinity to the receptors. For example, peptides comprised of a 23 amino acid portion of the C-terminal domain of VEGF have been synthesized which competitively inhibit binding of VEGF to VEGFR (Soker, et. al., J. Biol. Chem., 1997, 272, 31582-8). Linear peptides of 11 to 23 amino acid residues that bind to the basic FGF receptor (bFGFR) are described by Cosic et. al., Mol. and Cell. Biochem., 1994, 130, 1-9. A preferred linear peptide antagonist of the bFGFR is the 16 amino acid peptide, Met-Trp-Tyr-Arg-Pro-Asp-Leu-Asp-Glu-Arg-Lys-Gln-Gln-Lys-Arg-Glu. Gho et. al. (Cancer Research, 1997, 57, 3733-40) describe the identification of small peptides that bind with high affinity

to the angiogenin receptor on the surface of endothelial cells. A preferred peptide is Ala-Gln-Leu-Ala-Gly-Glu-Cys-Arg-Glu-Asn-Val-Cys-Met-Gly-Ile-Glu-Gly-Arg, in which the two Cys residues form an intramolecular disulfide bond. Yayon et. al. (Proc. Natl. Acad. Sci, USA, 1993, 90, 10643-7) describe other linear peptide antagonists of FGFR, identified from a random phage-displayed peptide library. Two linear octapeptides, Ala-Pro-Ser-Gly-His-Tyr-Lys-Gly and Lys-Arg-Thr-Gly-Gln-Tyr-Lys- Leu are preferred for inhibiting binding of bFGF to it receptor.

Targeting moieties for integrins expressed in tumor vasculature include peptides and peptidomimetics that bind to $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$. Pierschbacher and Rouslahti (J. Biol. Chem., 1987, 262, 17294-8) describe peptides that bind selectively to $\alpha_5\beta_1$ and $\alpha_v\beta_3$. U.S. 5,536,814 describe peptides that bind with high affinity to the integrin $\alpha_5\beta_1$. Burgess and Lim (J. Med. Chem., 1996, 39, 4520-6) disclose the synthesis three peptides that bind with high affinity to $\alpha_v\beta_3$: cyclo[Arg-Gly-Asp-Arg-Gly-Asp], cyclo[Arg-Gly-Asp-Arg-Gly-D-Asp] and the linear peptide Arg-Gly-Asp-Arg-Gly-Asp. U.S. 5,770,565 and U.S. 5,766,591 disclose peptides that bind with high affinity to $\alpha_v\beta_3$. U.S. 5,767,071 and U.S. 5,780,426, disclose cyclic peptides that have an exocyclic Arg amino acid that have high affinity for $\alpha_v\beta_3$. Srivatsa et. al., (Cardiovascular Res., 1997, 36, 408-28) describe the cyclic peptide antagonist for $\alpha_v\beta_3$, cyclo[Ala-Arg-Gly-Asp-Mamb]. Tran et. al., (Bioorg. Med. Chem. Lett., 1997, 7, 997-1002) disclose the cyclic peptide cyclo[Arg-Gly-Asp-Val-Gly-Ser-BTD-Ser-Gly-Val-Ala] that binds with high affinity to $\alpha_v\beta_3$. Arap et. al. (Science, 1998, 279, 377-80) describe cyclic peptides that bind to $\alpha_v\beta_3$ and $\alpha_v\beta_5$, Cys-Asp-Cys-Arg-Gly-Asp-Cys-Phe-Cys, and cyclo[Cys-Asn-Gly-Asp-Cys]. Corbett et. al. (Bioorg. Med. Chem. Lett., 1997, 7, 1371-6) describe a series of $\alpha_v\beta_3$ selective peptidomimetics. And Haubner et. al., (Angew. Chem. Int. Ed. Engl., 1997, 36, 1374-89) disclose peptides and peptidomimetic $\alpha_v\beta_3$ antagonists obtained from peptide libraries.

The targeting moieties of the present invention, preferably, have a binding affinity for the integrin $\alpha_v\beta_3$ of less than 1000nM. More preferably, the targeting moieties of the present invention, preferably, have a binding affinity for the integrin $\alpha_v\beta_3$ of less than 100nM. Even more preferably, the targeting moieties of the present invention, preferably, have a binding affinity for the integrin $\alpha_v\beta_3$ of less than 10nM.

The ultrasound contrast agents of the present invention comprise a plurality of angiogenic tumor vasculature targeting moieties attached to or incorporated into a microbubble of a biocompatible gas, a liquid carrier, and a surfactant microsphere, further comprising an optional linking moiety, L_n , between the targeting moieties and the microbubble. In this context, the term liquid carrier means aqueous solution and the term surfactant means any amphiphilic material which produces a reduction in interfacial tension in a solution. A list of suitable surfactants for forming surfactant microspheres is disclosed in EP0727225A2, herein incorporated by reference. The term surfactant microsphere includes nanospheres, liposomes, vesicles and the like. The biocompatible gas can be air, or a fluorocarbon, such as a C_3 - C_5 perfluoroalkane, for example, perfluoropropane, perfluorobutane, or perfluoropentane, which provides the difference in echogenicity and thus the contrast in ultrasound imaging. The gas is encapsulated or contained in the microsphere to which is attached the biodirecting group, optionally via a linking group. The attachment can be covalent, ionic or by van der Waals forces. Specific examples of such contrast agents include lipid encapsulated perfluorocarbons with a plurality of tumor neovasculature receptor binding peptides or peptidomimetics.

S_f as used herein is a surfactant which is either a lipid or a compound of the formula A^1-E-A^2 , defined above. The surfactant is intended to form a vesicle (e.g., a microsphere) capable of containing an echogenic gas. The ultrasound contrast agent compositions of the present invention are intended to be capable upon agitation (e.g., shaking,

stirring, etc...) of encapsulating an echogenic gas in a vesicle in such a way as to allow for the resultant product to be useful as an ultrasound contrast agent.

"Vesicle" refers to a spherical entity which is
5 characterized by the presence of an internal void. Preferred vesicles are formulated from lipids, including the various lipids described herein. In any given vesicle, the lipids may be in the form of a monolayer or bilayer, and the mono- or bilayer lipids may be used to form one of more mono- or
10 bilayers. In the case of more than one mono- or bilayer, the mono- or bilayers are generally concentric. The lipid vesicles described herein include such entities commonly referred to as liposomes, micelles, bubbles, microbubbles, microspheres and the like. Thus, the lipids may be used to
15 form a unilamellar vesicle (comprised of one monolayer or bilayer), an oligolamellar vesicle (comprised of about two or about three monolayers or bilayers) or a multilamellar vesicle (comprised of more than about three monolayers or bilayers). The internal void of the vesicles may be filled with a liquid,
20 including, for example, an aqueous liquid, a gas, a gaseous precursor, and/or a solid or solute material, including, for example, a bioactive agent, as desired.

"Vesicular composition" refers to a composition which is formulate from lipids and which comprises vesicles.

25 "Vesicle formulation" refers to a composition which comprises vesicles and a bioactive agent.

Microsphere, as used herein, is preferably a sphere of less than or equal to 10 microns. Liposome, as used herein, may include a single lipid layer (a lipid monolayer), two
30 lipid layers (a lipid bilayer) or more than two lipid layers (a lipid multilayer). "Lipsomes" refers to a generally spherical cluster or aggregate of amphipathic compounds, including lipid compounds, typically in the form of one or more concentric layers, for example, bilayers. They may also
35 be referred to herein as lipid vesicles.

The term "bubbles", as used herein, refers to vesicles which are generally characterized by the presence of one or more membranes or walls surrounding an internal void that is

filled with a gas or precursor thereto. Exemplary bubbles include, for example, liposomes, micelles and the like.

"Lipid" refers to a synthetic or naturally-occurring amphipathic compound which comprises a hydrophilic component
5 and a hydrophobic component. Lipids include, for example, fatty acids, neutral fats, phosphatides, glycolipids, aliphatic alcohols and waxes, terpenes and steroids.

"Lipid composition" refers to a composition which comprises a lipid compound. Exemplary lipid compositions
10 include suspensions, emulsions and vesicular compositions.

"Lipid formulation" refers to a composition which comprises a lipid compound and a bioactive agent.

Examples of classes of suitable lipids and specific suitable lipids include: phosphatidylcholines, such as
15 dioleoylphosphatidylcholine, dimyristoylphosphatidylcholine, dipalmitoylphosphatidylcholine (DPPC), and distearoylphosphatidylcholine; phosphatidylethanolamines, such as dipalmitoylphosphatidylethanolamine (DPPE), dioleoylphosphatidylethanolamine and N-succinyl-
20 dioleoylphosphatidylethanolamine; phosphatidylserines; phosphatidylglycerols; sphingolipids; glycolipids, such as ganglioside GM1; glucolipids; sulfatides; glycosphingolipids; phosphatidic acids, such as dipalmitoylphosphatidic acid (DPPA); palmitic fatty acids; stearic fatty acids; arachidonic
25 fatty acids; lauric fatty acids; myristic fatty acids; lauroleic fatty acids; physeteric fatty acids; myristoleic fatty acids; palmitoleic fatty acids; petroselinic fatty acids; oleic fatty acids; isolauric fatty acids; isomyristic fatty acids; isopalmitic fatty acids; isostearic fatty acids;
30 cholesterol and cholesterol derivatives, such as cholesterol hemisuccinate, cholesterol sulfate, and cholesteryl-(4'-trimethylammonio)-butanoate; polyoxyethylene fatty acid esters; polyoxyethylene fatty acid alcohols; polyoxyethylene fatty acid alcohol ethers; polyoxyethylated sorbitan fatty
35 acid esters; glycerol polyethylene glycol oxystearate; glycerol polyethylene glycol ricinoleate; ethoxylated soybean sterols; ethoxylated castor oil; polyoxyethylene-polyoxypropylene fatty acid polymers; polyoxyethylene fatty

acid stearates; 12-(((7'-diethylaminocoumarin-3-yl)-carbonyl)-methylamino)-octadecanoic acid; N-[12-(((7'-diethylamino-coumarin-3-yl)-carbonyl)-methyl-amino)octadecanoyl]-2-amino-palmitic acid; 1,2-dioleoyl-sn-glycerol; 1,2-dipalmitoyl-sn-3-
5 succinylglycerol; 1,3-dipalmitoyl-2-succinyl-glycerol; and 1-hexadecyl-2-palmitoyl-glycerophosphoethanolamine and palmitoylhomocysteine; lauryltrimethylammonium bromide; cetyltrimethylammonium bromide; myristyltrimethylammonium bromide; alkyl dimethylbenzylammonium chlorides, such as
10 wherein alkyl is a C₁₂, C₁₄ or C₁₆ alkyl; benzyldimethyldodecylammonium bromide; benzyldimethyldodecylammonium chloride, benzyldimethylhexadecylammonium bromide; benzyldimethylhexadecylammonium chloride;
15 benzyldimethyltetradecylammonium bromide; benzyldimethyltetradecylammonium chloride; cetyl dimethylethylammonium bromide; cetyl dimethylethylammonium chloride; cetylpyridinium bromide; cetylpyridinium chloride; N-[1,2,3-dioleoyloxy)-propyl]-N,N,N-trimethylammonium chloride
20 (DOTMA); 1,2-dioleoyloxy-3-(trimethylammonio)propane (DOTAP); and 1,2-dioleoyl-c-(4'-trimethylammonio)-butanoyl-sn-glycerol (DOTB).

The echogenic gas may be one gas or mixture of gases, such as CF₄, C₂F₆, C₃F₈, cyclo-C₄F₈, C₄F₁₀, C₅F₁₂, cyclo-
25 C₅F₁₀, cyclo-C₄F₇ (1-trifluoromethyl), propane (2-trifluoromethyl)-1,1,1,3,3,3 hexafluoro, and butane (2-trifluoromethyl)-1,1,1,3,3,3,4,4,4 nonafluoro. Also preferred are the the corresponding unsaturated versions of the above compounds, for example C₂F₄, C₃F₆, the isomers of C₄F₈. Also,
30 mixtures of these gases, especially mixtures of perfluorocarbons with other perfluorocarbons and mixtures of perfluorocarbons with other inert gases, such as air, N₂, O₂, He, would be useful. Examples of these can be found in Quay, U.S. Patent No. 5,595,723, the contents of which are herein
35 incorporated by reference.

X-ray contrast agents of the present invention are comprised of one or more angiogenic tumor vasculature targeting moieties attached to one or more X-ray absorbing or

"heavy" atoms of atomic number 20 or greater, further comprising an optional linking moiety, L_n , between the targeting moieties and the X-ray absorbing atoms. The frequently used heavy atom in X-ray contrast agents is iodine.

5 Recently, X-ray contrast agents comprised of metal chelates (Wallace, R., U.S. 5,417,959) and polychelates comprised of a plurality of metal ions (Love, D., U.S. 5,679,810) have been disclosed. More recently, multinuclear cluster complexes have been disclosed as X-ray contrast agents (U.S. 5,804,161, PCT
10 WO91/14460, and PCT WO 92/17215). Examples of X-ray agents include the non-radioactive or naturally occurring analogs of the above listed radionuclides (e.g., Re, Sm, Ho, Lu, Pm, Y, Bi, Pd, Gd, La, Au, Au, Yb, Dy, Cu, Rh, Ag, and Ir).

MRI contrast agents of the present invention are
15 comprised of one or more angiogenic tumor vasculature targeting moieties attached to one or more paramagnetic metal ions, further comprising an optional linking moiety, L_n , between the targeting moieties and the paramagnetic metal ions. The paramagnetic metal ions are present in the form of
20 metal complexes or metal oxide particles. U.S. 5,412,148, and 5,760,191, describe examples of chelators for paramagnetic metal ions for use in MRI contrast agents. U.S. 5,801,228, U.S. 5,567,411, and U.S. 5,281,704, describe examples of polychelants useful for complexing more than one paramagnetic
25 metal ion for use in MRI contrast agents. U.S. 5,520,904, describes particulate compositions comprised of paramagnetic metal ions for use as MRI contrast agents.

The pharmaceuticals of the present invention have the formulae, $(Q)_d-L_n-(C_h-X)$, $(Q)_d-L_n-(C_h-X^1)_d'$, $(Q)_d-L_n-(X^2)_d''$, and
30 $(Q)_d-L_n-(X^3)$, wherein Q represents a peptide or peptidomimetic that binds to a receptor expressed in angiogenic tumor vasculature, d is 1-10, L_n represents an optional linking group, C_h represents a metal chelator or bonding moiety, X represents a radioisotope, X^1 represents paramagnetic metal
35 ion, X^2 represents a paramagnetic metal ion or heavy atom containing insoluble solid particle, d'' is 1-100, and X^3 represents a surfactant microsphere of an echogenic gas. Preferred pharmaceuticals of the present invention are

comprised of targeting moieties, Q, that are peptides and peptidomimetics that bind to the vitronectin receptors $\alpha_v\beta_3$ and $\alpha_v\beta_5$. More preferred pharmaceuticals of the present invention are comprised of targeting moieties, Q, that are peptides and peptidomimetics that bind to $\alpha_v\beta_3$. Most preferred pharmaceuticals of the present invention are comprised of $\alpha_v\beta_3$ targeting moieties, Q, that are comprised of one to ten cyclic pentapeptides or peptidomimetics, independently attached to a therapeutic radioisotope or imageable moiety, further comprising an optional linking moiety, L_n , between the targeting moieties and the therapeutic radioisotopes or imageable moieties. The cyclic peptides are comprised of a tripeptide sequence that binds to the $\alpha_v\beta_3$ receptor and two amino acids either one of which can be attached to L_n , C_h , X^2 , or X^3 . The interaction of the tripeptide recognition sequences of the cyclic peptide or peptidomimetic portion of the pharmaceuticals with the $\alpha_v\beta_3$ receptor results in localization of the pharmaceuticals in angiogenic tumor vasculature, which express the $\alpha_v\beta_3$ receptor.

The pharmaceuticals of the present invention can be synthesized by several approaches. One approach involves the synthesis of the targeting peptide or peptidomimetic moiety, Q, and direct attachment of one or more moieties, Q, to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble. Another approach involves the attachment of one or more moieties, Q, to the linking group, L_n , which is then attached to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble. Another approach, useful in the synthesis of pharmaceuticals wherein d is 1, involves the synthesis of the moiety, Q- L_n , together, by incorporating an amino acid or amino acid mimetic residue bearing L_n into the synthesis of the peptide or peptidomimetic. The resulting moiety, Q- L_n , is then attached to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas

microbubble. Another approach involves the synthesis of a peptide or peptidomimetic, Q, bearing a fragment of the linking group, L_n, one or more of which are then attached to the remainder of the linking group and then to one or more metal chelators or bonding moieties, C_n, or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble.

The peptides or peptidomimetics, Q, optionally bearing a linking group, L_n, or a fragment of the linking group, can be synthesized using standard synthetic methods known to those skilled in the art. Preferred methods include but are not limited to those methods described below.

Generally, peptides and peptidomimetics are elongated by deprotecting the alpha-amine of the C-terminal residue and coupling the next suitably protected amino acid through a peptide linkage using the methods described. This deprotection and coupling procedure is repeated until the desired sequence is obtained. This coupling can be performed with the constituent amino acids in a stepwise fashion, or condensation of fragments (two to several amino acids), or combination of both processes, or by solid phase peptide synthesis according to the method originally described by Merrifield, J. Am. Chem. Soc., 85, 2149-2154 (1963), the disclosure of which is hereby incorporated by reference.

The peptides and peptidomimetics may also be synthesized using automated synthesizing equipment. In addition to the foregoing, procedures for peptide and peptidomimetic synthesis are described in Stewart and Young, "Solid Phase Peptide Synthesis", 2nd ed, Pierce Chemical Co., Rockford, IL (1984); Gross, Meienhofer, Udenfriend, Eds., "The Peptides: Analysis, Synthesis, Biology, Vol. 1, 2, 3, 5, and 9, Academic Press, New York, (1980-1987); Bodanszky, "Peptide Chemistry: A Practical Textbook", Springer-Verlag, New York (1988); and Bodanszky et al. "The Practice of Peptide Synthesis" Springer-Verlag, New York (1984), the disclosures of which are hereby incorporated by reference.

The coupling between two amino acid derivatives, an amino acid and a peptide or peptidomimetic, two peptide or

peptidomimetic fragments, or the cyclization of a peptide or peptidomimetic can be carried out using standard coupling procedures such as the azide method, mixed carbonic acid anhydride (isobutyl chloroformate) method, carbodiimide (dicyclohexylcarbodiimide, diisopropylcarbodiimide, or water-soluble carbodiimides) method, active ester (p-nitrophenyl ester, N-hydroxysuccinic imido ester) method, Woodward reagent K method, carbonyldiimidazole method, phosphorus reagents such as BOP-Cl, or oxidation-reduction method. Some of these methods (especially the carbodiimide) can be enhanced by the addition of 1-hydroxybenzotriazole. These coupling reactions may be performed in either solution (liquid phase) or solid phase.

The functional groups of the constituent amino acids or amino acid mimetics must be protected during the coupling reactions to avoid undesired bonds being formed. The protecting groups that can be used are listed in Greene, "Protective Groups in Organic Synthesis" John Wiley & Sons, New York (1981) and "The Peptides: Analysis, Synthesis, Biology, Vol. 3, Academic Press, New York (1981), the disclosure of which is hereby incorporated by reference.

The alpha-carboxyl group of the C-terminal residue is usually protected by an ester that can be cleaved to give the carboxylic acid. These protecting groups include: 1) alkyl esters such as methyl and t-butyl, 2) aryl esters such as benzyl and substituted benzyl, or 3) esters which can be cleaved by mild base treatment or mild reductive means such as trichloroethyl and phenacyl esters. In the solid phase case, the C-terminal amino acid is attached to an insoluble carrier (usually polystyrene). These insoluble carriers contain a group which will react with the carboxyl group to form a bond which is stable to the elongation conditions but readily cleaved later. Examples of which are: oxime resin (DeGrado and Kaiser (1980) *J. Org. Chem.* **45**, 1295-1300) chloro or bromomethyl resin, hydroxymethyl resin, and aminomethyl resin. Many of these resins are commercially available with the desired C-terminal amino acid already incorporated.

The alpha-amino group of each amino acid must be protected. Any protecting group known in the art can be used. Examples of these are: 1) acyl types such as formyl, trifluoroacetyl, phthalyl, and p-toluenesulfonyl; 2) aromatic carbamate types such as benzyloxycarbonyl (Cbz) and substituted benzyloxycarbonyls, 1-(p-biphenyl)-1-methylethoxycarbonyl, and 9-fluorenylmethyloxycarbonyl (Fmoc); 3) aliphatic carbamate types such as tert-butyloxycarbonyl (Boc), ethoxycarbonyl, diisopropylmethoxycarbonyl, and allyloxycarbonyl; 4) cyclic alkyl carbamate types such as cyclopentyloxycarbonyl and adamantyloxycarbonyl; 5) alkyl types such as triphenylmethyl and benzyl; 6) trialkylsilane such as trimethylsilane; and 7) thiol containing types such as phenylthiocarbonyl and dithiasuccinoyl. The preferred alpha-amino protecting group is either Boc or Fmoc. Many amino acid or amino acid mimetic derivatives suitably protected for peptide synthesis are commercially available.

The alpha-amino protecting group is cleaved prior to the coupling of the next amino acid. When the Boc group is used, the methods of choice are trifluoroacetic acid, neat or in dichloromethane, or HCl in dioxane. The resulting ammonium salt is then neutralized either prior to the coupling or in situ with basic solutions such as aqueous buffers, or tertiary amines in dichloromethane or dimethylformamide. When the Fmoc group is used, the reagents of choice are piperidine or substituted piperidines in dimethylformamide, but any secondary amine or aqueous basic solutions can be used. The deprotection is carried out at a temperature between 0 °C and room temperature.

Any of the amino acids or amino acid mimetics bearing side chain functionalities must be protected during the preparation of the peptide using any of the above-identified groups. Those skilled in the art will appreciate that the selection and use of appropriate protecting groups for these side chain functionalities will depend upon the amino acid or amino acid mimetic and presence of other protecting groups in the peptide or peptidomimetic. The selection of such a

protecting group is important in that it must not be removed during the deprotection and coupling of the alpha-amino group.

For example, when Boc is chosen for the alpha-amine protection the following protecting groups are acceptable:

5 p-toluenesulfonyl (tosyl) moieties and nitro for arginine; benzyloxycarbonyl, substituted benzyloxycarbonyls, tosyl or trifluoroacetyl for lysine; benzyl or alkyl esters such as cyclopentyl for glutamic and aspartic acids; benzyl ethers for serine and threonine; benzyl ethers, substituted benzyl ethers
10 or 2-bromobenzyloxycarbonyl for tyrosine; p-methylbenzyl, p-methoxybenzyl, acetamidomethyl, benzyl, or t-butylsulfonyl for cysteine; and the indole of tryptophan can either be left unprotected or protected with a formyl group.

When Fmoc is chosen for the alpha-amine protection
15 usually tert-butyl based protecting groups are acceptable. For instance, Boc can be used for lysine, tert-butyl ether for serine, threonine and tyrosine, and tert-butyl ester for glutamic and aspartic acids.

Once the elongation of the peptide or peptidomimetic, or
20 the elongation and cyclization of a cyclic peptide or peptidomimetic is completed all of the protecting groups are removed. For the liquid phase synthesis the protecting groups are removed in whatever manner as dictated by the choice of protecting groups. These procedures are well known to those
25 skilled in the art.

When a solid phase synthesis is used to synthesize a cyclic peptide or peptidomimetic, the peptide or peptidomimetic should be removed from the resin without simultaneously removing protecting groups from functional
30 groups that might interfere with the cyclization process. Thus, if the peptide or peptidomimetic is to be cyclized in solution, the cleavage conditions need to be chosen such that a free α -carboxylate and a free α -amino group are generated without simultaneously removing other protecting groups.
35 Alternatively, the peptide or peptidomimetic may be removed from the resin by hydrazinolysis, and then coupled by the azide method. Another very convenient method involves the synthesis of peptides or peptidomimetics on an oxime resin,

followed by intramolecular nucleophilic displacement from the resin, which generates a cyclic peptide or peptidomimetic (Osapay, Profit, and Taylor (1990) *Tetrahedron Letters* **43**, 6121-6124). When the oxime resin is employed, the Boc
5 protection scheme is generally chosen. Then, the preferred method for removing side chain protecting groups generally involves treatment with anhydrous HF containing additives such as dimethyl sulfide, anisole, thioanisole, or p-cresol at 0 °C. The cleavage of the peptide or peptidomimetic can also be
10 accomplished by other acid reagents such as trifluoromethanesulfonic acid/trifluoroacetic acid mixtures.

Unusual amino acids used in this invention can be synthesized by standard methods familiar to those skilled in the art ("The Peptides: Analysis, Synthesis, Biology, Vol. 5,
15 pp. 342-449, Academic Press, New York (1981)). N-Alkyl amino acids can be prepared using procedures described in previously (Cheung et al., (1977) *Can. J. Chem.* **55**, 906; Freidinger et al., (1982) *J. Org. Chem.* **48**, 77 (1982)), which are incorporated herein by reference.

20 Additional synthetic procedures that can be used by one of skill in the art to synthesize the peptides and peptidomimetics targeting moieties are described in PCT WO94/22910, the contents of which are herein incorporated by reference.


25 The attachment of linking groups, L_n , to the peptides and peptidomimetics, Q; chelators or bonding units, C_h , to the peptides and peptidomimetics, Q, or to the linking groups, L_n ; and peptides and peptidomimetics bearing a fragment of the linking group to the remainder of the linking group, in
30 combination forming the moiety, $(Q)_d-L_n$, and then to the moiety C_h ; can all be performed by standard techniques. These include, but are not limited to, amidation, esterification, alkylation, and the formation of ureas or thioureas. Procedures for performing these attachments can be found in
35 Brinkley, M., *Bioconjugate Chemistry* **1992**, 3(1), which is incorporated herein by reference.

A number of methods can be used to attach the peptides and peptidomimetics, Q, to paramagnetic metal ion or heavy

atom containing solid particles, X^2 , by one of skill in the art of the surface modification of solid particles. In general, the targeting moiety Q or the combination $(Q)_dL_n$ is attached to a coupling group that react with a constituent of the surface of the solid particle. The coupling groups can be any of a number of silanes which react with surface hydroxyl groups on the solid particle surface, as described in co-pending U.S.A.N 60/092,360, and can also include polyphosphonates, polycarboxylates, polyphosphates or mixtures thereof which couple with the surface of the solid particles, as described in U.S. 5,520,904.

A number of reaction schemes can be used to attach the peptides and peptidomimetics, Q, to the surfactant microsphere, X^3 . These are illustrated in following reaction schemes where S_f represents a surfactant moiety that forms the surfactant microsphere.

Acylation Reaction:




Y is a leaving group or active ester


Disulfide Coupling:



Sulfonamide Coupling:



Reductive Amidation:



In these reaction schemes, the substituents S_f and Q can be reversed as well.

The linking group L_n can serve several roles. First it provides a spacing group between the metal chelator or bonding moiety, C_h , the paramagnetic metal ion or heavy atom containing solid particle, X^2 , and the surfactant microsphere, X^3 , and the one or more of the peptides or peptidomimetics, Q , so as to minimize the possibility that the moieties C_h-X , C_h-X^1 , X^2 , and X^3 , will interfere with the interaction of the recognition sequences of Q with angiogenic tumor vasculature receptors. The necessity of incorporating a linking group in a reagent is dependent on the identity of Q , C_h-X , C_h-X^1 , X^2 , and X^3 . If C_h-X , C_h-X^1 , X^2 , and X^3 , cannot be attached to Q without substantially diminishing its affinity for the receptors, then a linking group is used. A linking group also provides a means of independently attaching multiple peptides and peptidomimetics, Q , to one group that is attached to C_h-X , C_h-X^1 , X^2 , or X^3 .

The linking group also provides a means of incorporating a pharmacokinetic modifier into the pharmaceuticals of the present invention. The pharmacokinetic modifier serves to direct the biodistribution of the injected pharmaceutical other than by the interaction of the targeting moieties, Q , with the receptors expressed in the tumor neovasculature. A wide variety of functional groups can serve as pharmacokinetic modifiers, including, but not limited to, carbohydrates, polyalkylene glycols, peptides or other polyamino acids, and cyclodextrins. The modifiers can be used to enhance or decrease hydrophilicity and to enhance or decrease the rate of blood clearance. The modifiers can also be used to direct the route of elimination of the pharmaceuticals. Preferred pharmacokinetic modifiers are those that result in moderate to fast blood clearance and enhanced renal excretion.

The metal chelator or bonding moiety, C_h , is selected to form stable complexes with the metal ion chosen for the particular application. Chelators or bonding moieties for diagnostic radiopharmaceuticals are selected to form stable complexes with the radioisotopes that have imageable gamma ray or positron emissions, such as ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{60}Cu , ^{64}Cu , ^{67}Ga , ^{68}Ga , ^{86}Y .

Chelators for technetium, copper and gallium isotopes are selected from diaminedithiols, monoamine-monoamidedithiols, triamide-monothiol, monoamine-diamide-monothiol, diaminedioximes, and hydrazines.

5 The chelators are generally tetradentate with donor atoms selected from nitrogen, oxygen and sulfur. Preferred reagents are comprised of chelators having amine nitrogen and thiol sulfur donor atoms and hydrazine bonding units. The thiol sulfur atoms and the hydrazines may bear a protecting group

10 which can be displaced either prior to using the reagent to synthesize a radiopharmaceutical or preferably in situ during the synthesis of the radiopharmaceutical.

Exemplary thiol protecting groups include those listed in Greene and Wuts, "Protective Groups in Organic Synthesis" John

15 Wiley & Sons, New York (1991), the disclosure of which is hereby incorporated by reference. Any thiol protecting group known in the art can be used. Examples of thiol protecting groups include, but are not limited to, the following: acetamidomethyl, benzamidomethyl, 1-ethoxyethyl, benzoyl, and

20 triphenylmethyl.

Exemplary protecting groups for hydrazine bonding units are hydrazones which can be aldehyde or ketone hydrazones having substituents selected from hydrogen, alkyl, aryl and heterocycle. Particularly preferred hydrazones are described

25 in co-pending U.S.S.N. 08/476,296 the disclosure of which is herein incorporated by reference in its entirety.

The hydrazine bonding unit when bound to a metal radionuclide is termed a hydrazido, or diazenido group and serves as the point of attachment of the radionuclide to the

30 remainder of the radiopharmaceutical. A diazenido group can be either terminal (only one atom of the group is bound to the radionuclide) or chelating. In order to have a chelating diazenido group at least one other atom of the group must also be bound to the radionuclide. The atoms bound to the metal

35 are termed donor atoms.

Chelators for ^{111}In and ^{86}Y are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, DO3A, 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-

tetraazacyclododecane-1-acetic-4,7,10-tris(methylacetic)acid,
2-benzyl-cyclohexyldiethylenetriaminepentaacetic acid, 2-
benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"-
tetra(carboxymethyl)aminomethyl)-4'-(3-amino-4-methoxyphenyl)-
5 2,2':6',2"-terpyridine. Procedures for synthesizing these
chelators that are not commercially available can be found in
Brechtel, M. and Gansow, O., *J. Chem. Soc. Perkin Trans.*
1992, 1, 1175; Brechtel, M. and Gansow, O., *Bioconjugate*
Chem. **1991**, 2, 187; Deshpande, S., et. al., *J. Nucl. Med.*
10 **1990**, 31, 473; Kruper, J., U.S. Patent 5,064,956, and Toner,
J., U.S. Patent 4,859,777, the disclosures of which are hereby
incorporated by reference in their entirety.

The coordination sphere of metal ion includes all the
ligands or groups bound to the metal. For a transition metal
15 radionuclide to be stable it typically has a coordination
number (number of donor atoms) comprised of an integer greater
than or equal to 4 and less than or equal to 8; that is there
are 4 to 8 atoms bound to the metal and it is said to have a
complete coordination sphere. The requisite coordination
20 number for a stable radionuclide complex is determined by the
identity of the radionuclide, its oxidation state, and the
type of donor atoms. If the chelator or bonding unit does not
provide all of the atoms necessary to stabilize the metal
radionuclide by completing its coordination sphere, the
25 coordination sphere is completed by donor atoms from other
ligands, termed ancillary or co-ligands, which can also be
either terminal or chelating.

A large number of ligands can serve as ancillary or
co-ligands, the choice of which is determined by a variety of
30 considerations such as the ease of synthesis of the
radiopharmaceutical, the chemical and physical properties of
the ancillary ligand, the rate of formation, the yield, and
the number of isomeric forms of the resulting
radiopharmaceuticals, the ability to administer said ancillary
35 or co-ligand to a patient without adverse physiological
consequences to said patient, and the compatibility of the
ligand in a lyophilized kit formulation. The charge and
lipophilicity of the ancillary ligand will effect the charge

and lipophilicity of the radiopharmaceuticals. For example, the use of 4,5-dihydroxy-1,3-benzene disulfonate results in radiopharmaceuticals with an additional two anionic groups because the sulfonate groups will be anionic under

5 physiological conditions. The use of N-alkyl substituted 3,4-hydroxypyridinones results in radiopharmaceuticals with varying degrees of lipophilicity depending on the size of the alkyl substituents.

Preferred technetium radiopharmaceuticals of the present
10 invention are comprised of a hydrazido or diazenido bonding unit and an ancillary ligand, A_{L1} , or a bonding unit and two types of ancillary A_{L1} and A_{L2} , or a tetradentate chelator comprised of two nitrogen and two sulfur atoms. Ancillary ligands A_{L1} are comprised of two or more hard donor atoms such
15 as oxygen and amine nitrogen (sp^3 hybridized). The donor atoms occupy at least two of the sites in the coordination sphere of the radionuclide metal; the ancillary ligand A_{L1} serves as one of the three ligands in the ternary ligand system. Examples of ancillary ligands A_{L1} include but are not
20 limited to dioxygen ligands and functionalized aminocarboxylates. A large number of such ligands are available from commercial sources.

Ancillary dioxygen ligands include ligands that coordinate to the metal ion through at least two oxygen donor
25 atoms. Examples include but are not limited to: glucoheptonate, gluconate, 2-hydroxyisobutyrate, lactate, tartrate, mannitol, glucarate, maltol, Kojic acid, 2,2-bis(hydroxymethyl)propionic acid, 4,5-dihydroxy-1,3-benzene disulfonate, or substituted or unsubstituted 1,2
30 or 3,4 hydroxypyridinones. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

Functionalized aminocarboxylates include ligands that have a combination of amine nitrogen and oxygen donor atoms.
35 Examples include but are not limited to: iminodiacetic acid, 2,3-diaminopropionic acid, nitrilotriacetic acid, N,N'-ethylenediamine diacetic acid, N,N,N'-ethylenediamine triacetic acid, hydroxyethylethylenediamine triacetic acid,

and N,N'-ethylenediamine bis-hydroxyphenylglycine. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

A series of functionalized aminocarboxylates are disclosed by Bridger et. al. in U.S. Patent 5,350,837, herein incorporated by reference, that result in improved rates of formation of technetium labeled hydrazino modified proteins. We have determined that certain of these aminocarboxylates result in improved yields of the radiopharmaceuticals of the present invention. The preferred ancillary ligands A_{L1} functionalized aminocarboxylates that are derivatives of glycine; the most preferred is tricine (tris(hydroxymethyl)methylglycine).

The most preferred technetium radiopharmaceuticals of the present invention are comprised of a hydrazido or diazenido bonding unit and two types of ancillary designated A_{L1} and A_{L2} , or a diaminedithiol chelator. The second type of ancillary ligands A_{L2} are comprised of one or more soft donor atoms selected from the group: phosphine phosphorus, arsine arsenic, imine nitrogen (sp^2 hybridized), sulfur (sp^2 hybridized) and carbon (sp hybridized); atoms which have p-acid character. Ligands A_{L2} can be monodentate, bidentate or tridentate, the denticity is defined by the number of donor atoms in the ligand. One of the two donor atoms in a bidentate ligand and one of the three donor atoms in a tridentate ligand must be a soft donor atom. We have disclosed in co-pending U.S.S.N. 08/415,908, and U.S.S.N. 60/013360 and 08/646,886, the disclosures of which are herein incorporated by reference in their entirety, that radiopharmaceuticals comprised of one or more ancillary or co-ligands A_{L2} are more stable compared to radiopharmaceuticals that are not comprised of one or more ancillary ligands, A_{L2} ; that is, they have a minimal number of isomeric forms, the relative ratios of which do not change significantly with time, and that remain substantially intact upon dilution.

The ligands A_{L2} that are comprised of phosphine or arsine donor atoms are trisubstituted phosphines, trisubstituted

arsines, tetrasubstituted diphosphines and tetrasubstituted diarsines. The ligands A_{L2} that are comprised of imine nitrogen are unsaturated or aromatic nitrogen-containing, 5 or 6-membered heterocycles. The ligands that are comprised of sulfur (sp^2 hybridized) donor atoms are thiocarbonyls, comprised of the moiety $C=S$. The ligands comprised of carbon (sp hybridized) donor atoms are isonitriles, comprised of the moiety CNR , where R is an organic radical. A large number of such ligands are available from commercial sources.

10 Isonitriles can be synthesized as described in European Patent 0107734 and in U.S. Patent 4,988,827, herein incorporated by reference.

Preferred ancillary ligands A_{L2} are trisubstituted phosphines and unsaturated or aromatic 5 or 6 membered heterocycles. The most preferred ancillary ligands A_{L2} are trisubstituted phosphines and unsaturated 5 membered heterocycles.

The ancillary ligands A_{L2} may be substituted with alkyl, aryl, alkoxy, heterocycle, aralkyl, alkaryl and arylalkaryl groups and may or may not bear functional groups comprised of heteroatoms such as oxygen, nitrogen, phosphorus or sulfur. Examples of such functional groups include but are not limited to: hydroxyl, carboxyl, carboxamide, nitro, ether, ketone, amino, ammonium, sulfonate, sulfonamide, phosphonate, and phosphonamide. The functional groups may be chosen to alter the lipophilicity and water solubility of the ligands which may affect the biological properties of the radiopharmaceuticals, such as altering the distribution into non-target tissues, cells or fluids, and the mechanism and rate of elimination from the body.

Chelators or bonding moieties for therapeutic radiopharmaceuticals are selected to form stable complexes with the radioisotopes that have alpha particle, beta particle, Auger or Coster-Kronig electron emissions, such as ^{186}Re , ^{188}Re , ^{153}Sm , ^{166}Ho , ^{177}Lu , ^{149}Pm , ^{90}Y , ^{212}Bi , ^{103}Pd , ^{109}Pd , ^{159}Gd , ^{140}La , ^{198}Au , ^{199}Au , ^{169}Yb , ^{175}Yb , ^{165}Dy , ^{166}Dy , ^{67}Cu , ^{105}Rh , ^{111}Ag , and ^{192}Ir . Chelators for rhenium, copper, palladium, platinum, iridium, rhodium, silver and gold

isotopes are selected from diaminedithiols, monoamine-monoamidedithiols, triamide-monothiol, monoamine-diamide-monothiol, diaminedioximes, and hydrazines. Chelators for yttrium, bismuth, and the lanthanide isotopes are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, DO3A, 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10-tris(methylacetic)acid, 2-benzyl-cyclohexyldiethylenetriaminepentaacetic acid, 2-benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"-tetra(carboxymethyl)aminomethyl]-4'-(3-amino-4-methoxyphenyl)-2,2':6',2"-terpyridine.

Chelators for magnetic resonance imaging contrast agents are selected to form stable complexes with paramagnetic metal ions, such as Gd(III), Dy(III), Fe(III), and Mn(II), are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, DO3A, 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10-tris(methylacetic)acid, 2-benzyl-cyclohexyldiethylenetriaminepentaacetic acid, 2-benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"-tetra(carboxymethyl)aminomethyl]-4'-(3-amino-4-methoxyphenyl)-2,2':6',2"-terpyridine.

The technetium and rhenium radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be easily prepared by admixing a salt of a radionuclide, a reagent of the present invention, an ancillary ligand A_{L1}, an ancillary ligand A_{L2}, and a reducing agent, in an aqueous solution at temperatures from 0 to 100 °C. The technetium and rhenium radiopharmaceuticals of the present invention comprised of a tetradentate chelator having two nitrogen and two sulfur atoms can be easily prepared by admixing a salt of a radionuclide, a reagent of the present invention, and a reducing agent, in an aqueous solution at temperatures from 0 to 100 °C.

When the bonding unit in the reagent of the present invention is present as a hydrazone group, then it must first be converted to a hydrazine, which may or may not be protonated, prior to complexation with the metal radionuclide.

The conversion of the hydrazone group to the hydrazine can occur either prior to reaction with the radionuclide, in which case the radionuclide and the ancillary or co-ligand or ligands are combined not with the reagent but with a hydrolyzed form of the reagent bearing the chelator or bonding unit, or in the presence of the radionuclide in which case the reagent itself is combined with the radionuclide and the ancillary or co-ligand or ligands. In the latter case, the pH of the reaction mixture must be neutral or acidic.

Alternatively, the radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be prepared by first admixing a salt of a radionuclide, an ancillary ligand A_{L1} , and a reducing agent in an aqueous solution at temperatures from 0 to 100 °C to form an intermediate radionuclide complex with the ancillary ligand A_{L1} then adding a reagent of the present invention and an ancillary ligand A_{L2} and reacting further at temperatures from 0 to 100 °C.

Alternatively, the radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be prepared by first admixing a salt of a radionuclide, an ancillary ligand A_{L1} , a reagent of the present invention, and a reducing agent in an aqueous solution at temperatures from 0 to 100 °C to form an intermediate radionuclide complex, and then adding an ancillary ligand A_{L2} and reacting further at temperatures from 0 to 100 °C.

The technetium and rhenium radionuclides are preferably in the chemical form of pertechnetate or perrhenate and a pharmaceutically acceptable cation. The pertechnetate salt form is preferably sodium pertechnetate such as obtained from commercial Tc-99m generators. The amount of pertechnetate used to prepare the radiopharmaceuticals of the present invention can range from 0.1 mCi to 1 Ci, or more preferably from 1 to 200 mCi.

The amount of the reagent of the present invention used to prepare the technetium and rhenium radiopharmaceuticals of the present invention can range from 0.01 µg to 10 mg, or more preferably from 0.5 µg to 200 µg. The amount used will

be dictated by the amounts of the other reactants and the identity of the radiopharmaceuticals of the present invention to be prepared.

The amounts of the ancillary ligands A_{L1} used can range
5 from 0.1 mg to 1 g, or more preferably from 1 mg to 100 mg. The exact amount for a particular radiopharmaceutical is a function of identity of the radiopharmaceuticals of the present invention to be prepared, the procedure used and the amounts and identities of the other reactants. Too large an
10 amount of A_{L1} will result in the formation of by-products comprised of technetium labeled A_{L1} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary ligand A_{L1} but without the ancillary ligand A_{L2} . Too small an amount of
15 A_{L1} will result in other by-products such as technetium labeled biologically active molecules with the ancillary ligand A_{L2} but without the ancillary ligand A_{L1} , or reduced hydrolyzed technetium, or technetium colloid.

The amounts of the ancillary ligands A_{L2} used can range
20 from 0.001 mg to 1 g, or more preferably from 0.01 mg to 10 mg. The exact amount for a particular radiopharmaceutical is a function of the identity of the radiopharmaceuticals of the present invention to be prepared, the procedure used and the amounts and identities of the other reactants. Too large an
25 amount of A_{L2} will result in the formation of by-products comprised of technetium labeled A_{L2} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary ligand A_{L2} but without the ancillary ligand A_{L1} . If the reagent bears
30 one or more substituents that are comprised of a soft donor atom, as defined above, at least a ten-fold molar excess of the ancillary ligand A_{L2} to the reagent of formula 2 is required to prevent the substituent from interfering with the coordination of the ancillary ligand A_{L2} to the metal
35 radionuclide.

Suitable reducing agents for the synthesis of the radiopharmaceuticals of the present invention include stannous salts, dithionite or bisulfite salts, borohydride salts, and

formamidinesulfinic acid, wherein the salts are of any pharmaceutically acceptable form. The preferred reducing agent is a stannous salt. The amount of a reducing agent used can range from 0.001 mg to 10 mg, or more preferably from 5 0.005 mg to 1 mg.

The specific structure of a radiopharmaceutical of the present invention comprised of a hydrazido or diazenido bonding unit will depend on the identity of the reagent of the present invention used, the identity of any ancillary ligand 10 A_{L1} , the identity of any ancillary ligand A_{L2} , and the identity of the radionuclide. Radiopharmaceuticals comprised of a hydrazido or diazenido bonding unit synthesized using concentrations of reagents of $<100 \mu\text{g/mL}$, will be comprised of one hydrazido or diazenido group. Those synthesized using >1 15 mg/mL concentrations will be comprised of two hydrazido or diazenido groups from two reagent molecules. For most applications, only a limited amount of the biologically active molecule can be injected and not result in undesired side-effects, such as chemical toxicity, interference with a 20 biological process or an altered biodistribution of the radiopharmaceutical. Therefore, the radiopharmaceuticals which require higher concentrations of the reagents comprised in part of the biologically active molecule, will have to be diluted or purified after synthesis to avoid such 25 side-effects.

The identities and amounts used of the ancillary ligands A_{L1} and A_{L2} will determine the values of the variables y and z . The values of y and z can independently be an integer from 1 to 2. In combination, the values of y and z will result in 30 a technetium coordination sphere that is made up of at least five and no more than seven donor atoms. For monodentate ancillary ligands A_{L2} , z can be an integer from 1 to 2; for bidentate or tridentate ancillary ligands A_{L2} , z is 1. The preferred combination for monodentate ligands is y equal to 1 35 or 2 and z equal to 1. The preferred combination for bidentate or tridentate ligands is y equal to 1 and z equal to 1.

The indium, copper, gallium, silver, palladium, rhodium, gold, platinum, bismuth, yttrium and lanthanide radiopharmaceuticals of the present invention can be easily prepared by admixing a salt of a radionuclide and a reagent of the present invention, in an aqueous solution at temperatures from 0 to 100 °C. These radionuclides are typically obtained as a dilute aqueous solution in a mineral acid, such as hydrochloric, nitric or sulfuric acid. The radionuclides are combined with from one to about one thousand equivalents of the reagents of the present invention dissolved in aqueous solution. A buffer is typically used to maintain the pH of the reaction mixture between 3 and 10.

The gadolinium, dysprosium, iron and manganese metallopharmaceuticals of the present invention can be easily prepared by admixing a salt of the paramagnetic metal ion and a reagent of the present invention, in an aqueous solution at temperatures from 0 to 100 °C. These paramagnetic metal ions are typically obtained as a dilute aqueous solution in a mineral acid, such as hydrochloric, nitric or sulfuric acid. The paramagnetic metal ions are combined with from one to about one thousand equivalents of the reagents of the present invention dissolved in aqueous solution. A buffer is typically used to maintain the pH of the reaction mixture between 3 and 10.

The total time of preparation will vary depending on the identity of the metal ion, the identities and amounts of the reactants and the procedure used for the preparation. The preparations may be complete, resulting in > 80% yield of the radiopharmaceutical, in 1 minute or may require more time. If higher purity metallopharmaceuticals are needed or desired, the products can be purified by any of a number of techniques well known to those skilled in the art such as liquid chromatography, solid phase extraction, solvent extraction, dialysis or ultrafiltration.

Buffers useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of said radiopharmaceuticals include but are not limited to phosphate, citrate, sulfosalicylate, and acetate.

A more complete list can be found in the United States Pharmacopeia.

Lyophilization aids useful in the preparation of diagnostic kits useful for the preparation of
5 radiopharmaceuticals include but are not limited to mannitol, lactose, sorbitol, dextran, Ficoll, and polyvinylpyrrolidone (PVP).

Stabilization aids useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the
10 preparation of radiopharmaceuticals include but are not limited to ascorbic acid, cysteine, monothioglycerol, sodium bisulfite, sodium metabisulfite, gentisic acid, and inositol.

Solubilization aids useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the
15 preparation of radiopharmaceuticals include but are not limited to ethanol, glycerin, polyethylene glycol, propylene glycol, polyoxyethylene sorbitan monooleate, sorbitan monooleate, polysorbates,
poly(oxyethylene)poly(oxypropylene)poly(oxyethylene) block
20 copolymers (Pluronic) and lecithin. Preferred solubilizing aids are polyethylene glycol, and Pluronic.

Bacteriostats useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the
preparation of radiopharmaceuticals include but are not
25 limited to benzyl alcohol, benzalkonium chloride, chlorbutanol, and methyl, propyl or butyl paraben.

A component in a diagnostic kit can also serve more than one function. A reducing agent can also serve as a
stabilization aid, a buffer can also serve as a transfer
30 ligand, a lyophilization aid can also serve as a transfer, ancillary or co-ligand and so forth.

The diagnostic radiopharmaceuticals are administered by intravenous injection, usually in saline solution, at a dose
of 1 to 100 mCi per 70 kg body weight, or preferably at a dose
35 of 5 to 50 mCi. Imaging is performed using known procedures.

The therapeutic radiopharmaceuticals are administered by intravenous injection, usually in saline solution, at a dose

of 0.1 to 100 mCi per 70 kg body weight, or preferably at a dose of 0.5 to 5 mCi per 70 kg body weight.

The magnetic resonance imaging contrast agents of the present invention may be used in a similar manner as other MRI agents as described in U.S. Patent 5,155,215; U.S. Patent 5,087,440; Margerstadt et al., Magn. Reson. Med., 1986, 3, 808; Runge et al., Radiology, 1988, 166, 835; and Bousquet et al., Radiology, 1988, 166, 693. Generally, sterile aqueous solutions of the contrast agents are administered to a patient intravenously in dosages ranging from 0.01 to 1.0 mmoles per kg body weight.

For use as X-ray contrast agents, the compositions of the present invention should generally have a heavy atom concentration of 1 mM to 5 M, preferably 0.1 M to 2 M. Dosages, administered by intravenous injection, will typically range from 0.5 mmol/kg to 1.5 mmol/kg, preferably 0.8 mmol/kg to 1.2 mmol/kg. Imaging is performed using known techniques, preferably X-ray computed tomography.

The ultrasound contrast agents of the present invention are administered by intravenous injection in an amount of 10 to 30 μ L of the echogenic gas per kg body weight or by infusion at a rate of approximately 3 μ L/kg/min. Imaging is performed using known techniques of sonography.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

Representative materials and methods that may be used in preparing the compounds of the invention are described further below.

Manual solid phase peptide synthesis was performed in 25 mL polypropylene filtration tubes purchased from BioRad Inc., or in 60 mL hour-glass reaction vessels purchased from Peptides International. Oxime resin (substitution level = 0.96 mmol/g) was prepared according to published procedure (DeGrado and Kaiser, *J. Org. Chem.* **1980**, 45, 1295), or was purchased

from Novabiochem (substitution level = 0.62 mmol/g). All chemicals and solvents (reagent grade) were used as supplied from the vendors cited without further purification. t-Butyloxycarbonyl (Boc) amino acids and other starting amino acids may be obtained commercially from Bachem Inc., Bachem Biosciences Inc. (Philadelphia, PA), Advanced ChemTech (Louisville, KY), Peninsula Laboratories (Belmont, CA), or Sigma (St. Louis, MO). 2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and TBTU were purchased from Advanced ChemTech. N-methylmorpholine (NMM), m-cresol, D-2-aminobutyric acid (Abu), trimethylacetylchloride, diisopropylethylamine (DIEA), 1,2,4-triazole, stannous chloride dihydrate, and tris(3-sulfonatophenyl)phosphine trisodium salt (TPPTS) were purchased from Aldrich Chemical Company. Bis(3-sulfonatophenyl)phenylphosphine disodium salt (TPPDS) was prepared by the published procedure (Kuntz, E., U.S. Patent 4,248,802). (3-Sulfonatophenyl)diphenylphosphine monosodium salt (TPPMS) was purchased from TCI America, Inc. Tricine was obtained from Research Organics, Inc. Technetium-99m-pertechnetate ($^{99m}\text{TcO}_4^-$) was obtained from a DuPont Pharma $^{99}\text{Mo}/^{99m}\text{Tc}$ Technelite® generator. In-111-chloride (Indichlor®) was obtained from Amersham Medi-Physics, Inc. Sm-153-chloride and Lutetium-177-chloride were obtained from the University of Missouri Research Reactor (MURR). Yttrium-90 chloride was obtained from the Pacific Northwest Research Laboratories. Dimethylformamide (DMF), ethyl acetate, chloroform (CHCl_3), methanol (MeOH), pyridine and hydrochloric acid (HCl) were obtained from Baker. Acetonitrile, dichloromethane (DCM), acetic acid (HOAc), trifluoroacetic acid (TFA), ethyl ether, triethylamine, acetone, and magnesium sulfate were commercially obtained. Absolute ethanol was obtained from Quantum Chemical Corporation.

General Procedure for Solid Phase Peptide Synthesis Using Boc-Chemistry on the Oxime Resin for the Preparation of Cyclic Peptides

The appropriately protected cyclic peptides, described in the Examples, were prepared by manual solid phase peptide synthesis using Boc-teabag chemistry (Houghton, 1985) on a p-nitrobenzophenone oxime solid support (DeGrado, 1982, Scarr and Findeis, 1990). The 5.0 cm x 5.0 cm teabags were made from 0.75 mm mesh polypropylene filters (Spectra Filters) and filled with 0.5 g (or 1 g) of the oxime resin. The coupling and deprotection steps were carried out in a polypropylene reactor using a table-top shaker for agitation. Synthesis of the protected pentapeptide-resin intermediate was achieved by first coupling Boc-Gly-OH to the oxime resin (substitution 0.69 mmol/g or 0.95 mmol/g). Attachment of Boc-Gly-OH onto the oxime resin was achieved by using five equivalents each of the amino acid, HBTU and diisopropylethylamine (DIPEA) in DMF. Coupling of the first amino acid generally occurred over 2-3 days. After thorough washing, substitution levels were determined using the picric acid assay (Stewart and Martin). Unreacted oxime groups on the resin were then capped with a solution of DIPEA and trimethylacetyl chloride in DMF. The boc-group was deprotected using 50% or 25% TFA in DCM (30 min). Coupling of the other protected boc-amino acids were performed in a similar manner by overnight shaking (1-2 days), and the coupling yields for each newly added amino acid was determined using the picric acid assay.

General Procedure for Solid Phase Peptide Synthesis Using Fmoc-Chemistry on the HMPB-BHA Resin for the Preparation of Cyclic Peptides

The appropriately protected linear peptide precursors to the cyclic peptides, described in the Examples, were also prepared by automated solid phase peptide synthesis using Fmoc chemistry on an Advanced ChemTech Model 90 Synthesizer and using HMPB-BHA resin as the solid support. Synthesis of the protected pentapeptide-resin intermediates was achieved by coupling (for 3 h) the Fmoc-amino acids sequentially to the commercially available (Novabiochem) Fmoc-Gly-HMPB-BHA resin (usually 2 g, substitution 0.47 to 0.60 mmol/g) by using three

to five equivalents each of the amino acid, HBTU, HOBt and diisopropylethylamine (DIPEA) in DMF. The Fmoc-group was deprotected using 20% piperidine in DMF (30 min). The peptides were cleaved from the HMPB-BHA resin using a solution of 1%
5 TFA/DCM and collecting the peptide solutions in a solution of pyridine in methanol (1:10). The linear protected peptides were isolated by removing the solvents and reagents in vacuo and triturating the crude residue in diethyl ether.

The syntheses of several amino acids that are not
10 commercially available are described in the following procedures.

Synthesis of Tfa-amino acids

Boc-HomoLys(Tfa)-OH and Boc-Cys(2-N-Tfa-aminoethyl)-OH
15 are prepared via the reaction of Boc-HomoLys-OH and Boc-Cys(2-aminoethyl)-OH, respectively, with ethyl thioltrifluoroacetate in Aq. NaOH, and purified by recrystallization from ethanol.

Synthesis of Boc-Orn(d-N-Benzylcarbamoyl)

To a solution of Boc-Orn (1 mmol) in DMF (30 mL) is added benzylisocyanate (2.2 mmol), and diisopropylamine (3 mmol). The reaction mixture is then stirred overnight at room temperature. The volatiles are removed in vacuo and the crude
25 material is purified by column chromatography to obtain the desired product.

Synthesis of Boc-Orn(d-N-1-Tos-2-ImidazolinyI)

A solution of Boc-Orn-OH (10 mmol), 1-tosyl-2-methylthio-
30 2-imidazoline (12 mmol, (which in turn is prepared from the reaction of the commercially available 2-methylthio-2-imidazoline hydriodide and p-toluenesulfonic anhydride in methylene chloride (0 °C to RT) in the presence of triethylamine)), and diisopropylethylamine (12 mmol) is
35 stirred at reflux, overnight. The volatiles are removed and the desired product isolated by chromatography.

Synthesis of Dap(b-(1-Tos-2-benzimidazolylacetyl))

To a solution of 1-Tos-2-benzimidazolylacetic acid (10 mmol, prepared using tosyl chloride and standard reported conditions) and N-methylmorpholine (10 mmol) in anhydrous DMF is added isobutyl chloroformate (10 mmol). After stirring at
5 ice bath temperature for 5-10 min., Boc-Orn-OH (10 mmol) and N-methylmorpholine (20 mmol) in anhydrous DMF is added in one portion. The reaction mixture is stirred overnight at room temperature, the volatiles removed in vacuo, and the product is isolated by chromatography. (Alternatively, Boc-Orn-OMe is
10 used and the product isolated is treated with aqueous LiOH to obtain the acid.)

The analytical HPLC methods utilized are described below:

HPLC Method 1


15 Instrument: HP1050
Column: Vydac C18(4.6 x 250 mm)
Detector: Diode array detector 220nm/500ref
Flow Rate: 1.0 mL/min.
Column Temp: 50 °C
20 Sample Size: 15 uL
Mobile Phase: A: 0.1% TFA in water
B: 0.1% TFA in ACN/Water (9:1)

Gradient A:	Time (min)	%A	%B
25	0	80	20
	20	0	100
	30	0	100
	31	80	20

Gradient B:	Time (min)	%A	%B
30	0	98	2
	16	63.2	36.8
	18	0	100
	28	0	100
35	30	98	2

Example 1

Synthesis of cyclo{Arg-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}




5

Part A: Preparation of cyclo{Arg(Tos)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val}

- 10 The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Tyr(N-Cbz-aminopropyl)-Val-Arg(Tos)-Gly-Oxime resin was removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with DCM, the resin was treated with 10% DIEA/DCM (2 x 10 min.). The resin was
- 15 subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.7474 g, 0.55 mmol/g) was then suspended in dimethylformamide (15 mL). Glacial acetic acid (55.0 μ L, 0.961 mmol) was added, and the reaction mixture was heated at 50 °C for 72 h. The resin was filtered, and washed with DMF
- 20 (2 x 10 mL). The filtrate was concentrated to an oil under high vacuum. The resulting oil was triturated with ethyl acetate. The solid thus obtained was filtered, washed with ethyl acetate, and dried under high vacuum to give 444.4 mg of the desired product. ESMS: Calcd. for C₅₁H₆₃N₉O₁₂S, 1025.43; Found, 1026.6 [M+H]⁺1. Analytical HPLC, Method 1A, R_t =
- 25 14.366 min, Purity = 75%.

Part B: Preparation of cyclo{Arg-Gly-Asp-D-Tyr(3-aminopropyl)-Val} Trifluoroacetic acid salt.



Cyclo{Arg(Tos)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-
 5 Val} (0.150 g, 0.146 mmol) was dissolved in trifluoroacetic
 acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic
 acid (0.5 mL) was added dropwise, maintaining the temperature
 at -10 °C. Anisole (0.1 mL) was added and the reaction
 mixture was stirred at -10 °C for 3 h. Diethyl ether was
 10 added, the reaction mixture cooled to -35 °C and then stirred
 for 30 min. The reaction mixture was cooled further to -50 °C
 and stirred for 30 min. The crude product obtained was
 filtered, washed with diethyl ether, dried under high vacuum,
 and purified by preparative HPLC Method 1, to give 29.7 mg
 15 (23%) of the desired product as a lyophilized solid. ESMS:
 Calcd. for C₂₉H₄₅N₉O₈, 647.34; Found, 648.5 [M+H]⁺.
 Analytical HPLC, Method 1B, R_t = 10.432 min, Purity = 91%.

Preparative HPLC Method 1

20 Instrument: Rainin Rabbit; Dynamax software
 Column: Vydac C-18 (21.2 mm x 25 cm)
 Detector: Knauer VWM
 Flow Rate: 15ml/min
 Column Temp: RT
 25 Mobile Phase: A: 0.1% TFA in H₂O
 B: 0.1%TFA in ACN/H₂O (9:1)
 Gradient:

Time (min)	%A	%B
0	98	2
16	63.2	36.8
18	0	100
28	0	100
30	98	2


30

Part C. Preparation of cyclo{Arg-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}

- 5 Cyclo{Arg-Gly-Asp-D-Tyr(3-aminopropyl)-Val}
 trifluoroacetic acid salt (0.020 g, 0.0228 mmol) was dissolved
 in DMF (1 mL). Triethylamine (9.5 μ L, 0.0648 mmol) was added,
 and after 5 min of stirring 2-[[[5-[(2,5-dioxo-1-
 pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-
 10 benzenesulfonic acid, monosodium salt (0.0121 g, 0.0274 mmol)
 was added. The reaction mixture was stirred for 7 days, and
 then concentrated to an oil under high vacuum. The oil was
 purified by preparative HPLC Method 1 to give 8.9 mg (37%) of
 the title product as a lyophilized solid (TFA salt). HRMS:
 15 Calcd. for $C_{42}H_{54}N_{12}O_{12}S + H$, 951.3783; Found, 951.3767.
 Analytical HPLC, Method 1B, R_t = 14.317 min, Purity = 95%.

Example 2

- Synthesis of cyclo{Arg-Gly-Asp-D-Tyr((N-[2-[[[5-[carbonyl]-2-
 20 pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-18-amino-14-
 aza-4,7,10-oxy-15-oxo-octadecoyl)-3-aminopropyl)-Val}




Part A: Preparation of 3-(N-(3-(2-(2-(3-((tert-butoxy)-carbonylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)-propanoic acid

5 N-(3-(2-(2-(3-Aminopropoxy)ethoxy)ethoxy)propyl)(tert-butoxy)formamide (1.5 g, 4.68 mmol) was added to DMF (15 mL). To this solution pyridine (15 mL), succinic anhydride (0.47 g, 4.68 mmol) were added, followed by dimethylaminopyridine (62 mL, 0.468 μ mol). The reaction mixture was stirred overnight
10 at 100 °C. The mixture was concentrated under high vacuum and the residue was brought up in water, acidified to pH 2.5 with 1N HCl, and extracted with ethyl acetate (3x). The combined organic extracts were dried over MgSO₄ and filtered. The filtrate was concentrated *in vacuo* to provide 1.24 g of an oil
15 product (63%). The desired product was used without further purification. ¹H NMR (CDCl₃) 3.67-3.45 (m, 11H), 3.41-3.28 (m, 2H), 3.21-3.09 (m, 2H), 2.95-2.82 (m, 2H), 2.80-2.35 (m, 3H), 1.81-1.68 (m, 4H), 1.50-1.35 (s, 9H); ESMS: Calculated for C₁₉H₃₆N₂O₈, 420.2471 Found 419.3 [M-H]-1.

20

Part B: Preparation of 3-(N-(3-(2-(2-(3-((tert-butoxy)-carbonylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)-propanoic acid succinimide ester




25

To a solution of 3-(N-(3-(2-(2-(3-((tert-butoxy)-carbonylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)-propanoic acid (1.12 g, 2.66 mmol), N-hydroxysuccinimide (0.40
30 g, 3.46 mmol), and N,N-dimethylformamide (40 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.67 g, 3.46 mmol). The reaction mixture was stirred at room temperature for 48 h. The mixture was concentrated under high vacuum and the residue was brought up in 0.1N HCl and extracted with
35 ethyl acetate (3x). The combined organic extracts were washed with water (2x) then saturated sodium chloride, dried over

MgSO₄, and filtered. The filtrate was concentrated *in vacuo* to give 1.0 g of the product as an oil (73%). The desired product was used without further purification. ESMS: Calculated for C₂₃H₃₉N₃O₁₀, 517.2635 Found 518.2 [M+H]⁺1.

5

Part C. Preparation of cyclo{Arg-Gly-Asp-D-Tyr(3-(3-(N-(3-(2-(2-(3-((tert-butoxy)-carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)-propanamido)propyl)-Val}



10

Cyclo{Arg-Gly-Asp-D-Tyr(3-aminopropyl)-Val}. TFA salt (0.040 g, 0.0457 mmol) was dissolved in DMF (2 mL). Triethylamine (19.1 μ L, 0.137 mmol) was added and after stirring for 5 minutes 3-(N-(3-(2-(2-(3-((tert-butoxy)-carbonylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)propanoic acid succinimide ester (0.0284 g, 0.0548 mmol) was added. The reaction mixture was stirred under N₂ for 48 h and then concentrated to an oil under high vacuum. The oil was trituated with ethyl acetate, the product filtered, washed with ethyl acetate, and dried under high vacuum. The crude product was purified by Preparative HPLC Method 1 to give 7.4 mg (14%) of the desired product as a lyophilized solid. ESMS: Calcd. for C₄₈H₇₉N₁₁O₁₅, 1049.58; Found, 1050.5 [M+H]⁺1.

Analytical HPLC, Method 1B, R_t = 20.417 min, Purity = 100%.

Part D. Preparation of cyclo{Arg-Gly-Asp-D-Tyr(3-(3-(N-(3-(2-(2-(3-(amino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)-propanamido)propyl)-Val}


Cyclo{Arg-Gly-Asp-D-Tyr(3-(3-(N-(3-(2-(2-(3-((tert-butoxy)-carbonylamino)propoxy)ethoxy)ethoxy)propyl)-carbamoyl)-propanamido)propyl)-Val} (6.0 mg, 0.00515 mmol) was dissolved in methylene chloride (1 mL) and trifluoroacetic acid (1 mL) was added. The solution stirred for 2 h and then concentrated to an oil under high vacuum. The oil was triturated with diethyl ether, the product filtered, washed with diethyl ether, and dried under high vacuum to give 6.0 mg (98%) of the desired product. ESMS: Calcd. for C₄₃H₇₁N₁₁O₁₃, 949.52; Found, 950.6 [M+H]⁺+1. Analytical HPLC, Method 1B, R_t = 14.821 min, Purity = 73%.

Part E. Preparation of cyclo{Arg-Gly-Asp-D-Tyr((N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-18-amino-14-aza-4,7,10-oxy-15-oxo-octadecoyl)-3-aminopropyl)-Val}

Cyclo{Arg-Gly-Asp-D-Tyr(3-(3-(N-(3-(2-(2-(3-(amino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)-propanamido)propyl)-Val} (5.0 mg, 0.00424 mmol) was dissolved in dimethylformamide (1 mL). Triethylamine (1.8 µL, 0.0127 mmol) was added, and after stirring for 5 min 2-[[[5-[[[2,5-dioxo-1-pyrrolidinyl)oxy]-carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (2.2 mg, 0.00509 mmol) was added. The reaction mixture was stirred for 24 h and then concentrated to an oil under high vacuum. The oil was purified by preparative HPLC Method 1 to give 2.2 mg (38%) of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for C₅₆H₈₀N₁₄O₁₇S, 1252.6; Found, 1253.7 (M+H⁺). Analytical HPLC, Method 1B, R_t = 17.328 min, Purity = 100%.

Example 3

Synthesis of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp}




Part A. Preparation of Boc-Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp}

5

Cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp} (0.040 g, 0.0457 mmol) was dissolved in dimethylformamide (2 mL). Triethylamine (19.1 μ L, 0.137 mmol) was added and the reaction mixture was stirred for 5 minutes. Boc-Glu(OSu)-OSu (0.0101 g, 0.0229 mmol) was added and the reaction mixture was stirred under N₂ for 18 h. The reaction mixture was then concentrated to an oil under high vacuum. The oil was triturated with ethyl acetate. The product was filtered, washed with ethyl acetate, and dried under high vacuum to give 38.0 mg (55%) of the desired product. ESMS: Calcd. for C₆₈H₁₀₃N₁₉O₂₀, 1505.76; Found, 1504.9 [M-H]⁻1. Analytical HPLC, Method 1B, R_t = 19.797 min, Purity = 73%.

Part B. Preparation of Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp}. TFA salt

20



Boc-Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp} (0.035 g, 0.0232 mmol) was dissolved in methylene chloride (1 mL). Trifluoroacetic acid (1 mL) was added, and the reaction mixture was stirred for 2 h, concentrated to an oil under high vacuum and triturated with ether. The product obtained was filtered, washed with diethyl ether, and dried under high vacuum to give 30.7 mg (76%) of the desired product. ESMS: Calcd. for C₆₃H₉₅N₁₉O₁₈, 1405.71; Found, 1404.7 [M-H]⁻1.

10 Analytical HPLC, Method 1B, R_t = 15.907 min, Purity = 77%.

Part C. Preparation of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp}

15

To a solution of Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp} (0.025 g, 0.0143 mmol) in dimethylformamide (2 mL) was added triethylamine (6.0 μL, 0.0429 mmol) and the reaction mixture was stirred for 5 min. 2-[[[5-[(2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0076 g, 0.0172 mmol) was added, and the reaction mixture was stirred for 5 days, then concentrated to an oil under high vacuum. The oil was purified by Preparative HPLC Method 1 to give 12.0 mg (43%) of the desired product as a lyophilized solid. ESMS: Calcd. for C₇₆H₁₀₄N₂₂O₂₂S, 1708.7; Found, 1710.1 (M+H⁺). Analytical HPLC, Method 1B, R_t = 17.218 min, Purity = 94%.

20


25

30

Example 4

Synthesis of cyclo(Arg-Gly-Asp-D-Tyr-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]]))

35




Part A. Preparation of cyclo{Arg(Tos)-Gly-Asp(OBzl)-D-Tyr(Bzl)-Lys(Cbz)}

5

The N-terminus Boc-protecting group of the peptide sequence Boc-Asp(OBzl)-D-Tyr(Bzl)-Lys(Z)-Arg(Tos)-Gly-oxime resin was removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with DCM, the resin was treated with 10% DIEA/DCM (2 x 10 min.). The resin was subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.8711 g, 0.44 mmol/g) was then suspended in DMF (15 mL). Glacial acetic acid (47.1 μ L, 0.823 mmol) was added, and the reaction was heated at 60 °C for 72 h. The resin was filtered, and washed with DMF (2 x 10 mL). The filtrate was concentrated to an oil under high vacuum. The resulting oil was triturated with ethyl acetate. The solid thus obtained was filtered, washed with ethyl acetate, and dried under high vacuum to give 653.7 mg of the desired product. ESMS: Calcd. for C₅₆H₆₅N₉O₁₂S, 1087.45; Found, 1088.7 [M+H]⁺+1. Analytical HPLC, Method 1A, R_t = 17.559 min, Purity = 82%.

Part B. Preparation of cyclo{Arg-Gly-Asp-D-Tyr-Lys}



Cyclo{Arg(Tos)-Gly-Asp(OBzl)-D-Tyr(Bzl)-Lys(Cbz)} (0.200 g, 0.184 mmol) was dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid (0.5 mL) was added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) was added and the reaction mixture was stirred at -10 °C for 3 h. Diethyl ether was added, the reaction was cooled to -50 °C, and stirred for 1 h. The crude product was filtered, washed with diethyl ether, and dried under high vacuum. The crude product was purified by Preparative HPLC Method 1, to give 15.2 mg (10%) of the desired product as a lyophilized solid. HRMS: Calcd. for C₂₇H₄₁N₉O₈ + H, 620.3156; Found, 620.3145. Analytical HPLC, Method 1B, R_t = 8.179 min, Purity = 100%.


Part C. Preparation of cyclo{Arg-Gly-Asp-D-Tyr-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])}

Cyclo{Arg-Gly-Asp-D-Tyr-Lys} TFA salt (0.010 g, 0.0118 mmol) was dissolved in DMF (1 mL). Triethylamine (5.0 µL, 0.0354 mmol) was added, and after stirring for 5 min 2-[[[5-[[[(2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.0062 g, 0.0142 mmol) was added. The reaction mixture was stirred for 20 h and then concentrated to an oil under high vacuum. The oil was purified by Preparative HPLC Method 1 to give 6.2 mg (46%) of the desired product as a lyophilized solid. HRMS: Calcd. for C₄₀H₅₀N₁₂O₁₂S + H, 923.3470; Found,

923.3486. Analytical HPLC, Method 1B, R_t = 11.954 min, Purity = 100%.

Example 5

- 5 Synthesis of cyclo{Arg-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])}




- 10 Part A. Preparation of cyclo{Arg(Tos)-Gly-Asp(OBzl)-D-Phe-Lys(Cbz)}

The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Phe-Lys(Z)-Arg(Tos)-Gly-Oxime resin
 15 was removed using standard deprotection (25% TFA in CH_2Cl_2). After eight washes with DCM, the resin was treated with 10% DIEA/DCM (2 x 10 min.). The resin was subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.7053 g, 0.44 mmol/g) was then suspended in dimethylformamide (15
 20 mL). Glacial acetic acid (43.0 μL , 0.750 mmol) was added, and the reaction was heated to 60 $^\circ\text{C}$ for 72 h. The resin was filtered, and washed with DMF (2 x 10 mL). The filtrate was concentrated to an oil under high vacuum. The resulting oil was triturated with ethyl acetate. The solid thus obtained
 25 was filtered, washed with ethyl acetate, and dried under high vacuum to give 510.3 mg of the desired product. ESMS: Calcd.

for C₄₉H₅₉N₉O₁₁S, 981.40; Found, 982.6 [M+H]⁺. Analytical HPLC, Method 1A, R_t = 15.574 min, Purity = 89%.

Part B. Preparation of cyclo{Arg-Gly-Asp-D-Phe-Lys}

5



Cyclo{Arg(Tos)-Gly-Asp(OBzl)-D-Phe-Lys(Cbz)} (0.200 g, 0.204 mmol) was dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid (0.5 mL) was added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) was added and the reaction was stirred at -10 °C for 3 h. Diethyl ether was added, the reaction was cooled to -50 °C, and stirred for 1 h. The crude product was filtered, washed with diethyl ether, dried under high vacuum and purified by Preparative HPLC Method 1, to give 121.1 mg (71%) of the desired product as a lyophilized solid. HRMS: Calcd. for C₂₇H₄₁N₉O₇ +H, 604.3207; Found, 604.3206. Analytical HPLC, Method 1B, R_t = 11.197 min, Purity = 100%.

20

Part C. Preparation of cyclo{Arg-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])}

25


Cyclo{Arg-Gly-Asp-D-Phe-Lys} TFA salt (0.040 g, 0.0481 mmol) was dissolved in DMF (2 mL). Triethylamine (20.1 µL, 0.144 mmol) was added, and after 5 min of stirring 2-[[[5-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.0254 g, 0.0577 mmol) was added. The reaction mixture was stirred for 20 h and then concentrated to an oil under

30

high vacuum. The oil was purified by Preparative HPLC Method 1 to give 38.2 mg (78%) of the desired product as a lyophilized solid. HRMS: Calcd. for $C_{40}H_{50}N_{12}O_{11}S + H$, 907.3521; Found, 907.3534. Analytical HPLC, Method 1B, $R_t =$
 5 14.122 min, Purity = 91%.


Example 6

Synthesis of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-
 benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-
 10 cyclo{Lys-Arg-Gly-Asp-D-Phe}



Part A. Preparation of Boc-Glu(OSu)-OSu

15



To a solution of Boc-Glu-OH (8.0 g, 32.25 mmol), N-hydroxysuccinimide (8.94 g, 77.64 mmol), and DMF (120 mL) was
 20 added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (14.88 g, 77.64 mmol). The reaction mixture was stirred at room temperature for 48 h. The mixture was concentrated under high vacuum and the residue was brought up in 0.1 N HCl and extracted with ethyl acetate (3x). The combined organic
 25 extracts were washed with water, saturated sodium bicarbonate and then saturated sodium chloride, dried over $MgSO_4$, and filtered. The filtrate was concentrated *in vacuo* and purified via reverse-phase HPLC (Vydac C18 column, 18 to 90 %

acetonitrile gradient containing 0.1% TFA, R_t = 9.413 min) to afford 8.5 g (60%) of the desired product as a white powder.

^1H NMR (CDCl_3): 2.98-2.70 (m, 11H), 2.65-2.25 (m, 2H), 1.55-1.40 (s, 9H); ESMS: Calculated for $\text{C}_{18}\text{H}_{23}\text{N}_3\text{O}_{10}$, 441.1383


5 Found 459.2 $[\text{M}+\text{NH}_4]^+$ 1.

Part B. Preparation of Boc-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe}

10 To a solution of cyclo(Lys-Arg-Gly-Asp-D-Phe) (0.050 g, 0.0601 mmol) in dimethylformamide (2 mL) was added triethylamine (25.1 μL , 0.183 mmol). After stirring for 5 minutes Boc-Glu(OSu)-OSu (0.0133 g, 0.0301 mmol) was added. The reaction mixture was stirred under N_2 for 20 h, then
 15 concentrated to an oil under high vacuum and triturated with ethyl acetate. The product thus obtained was filtered, washed with ethyl acetate, and dried under high vacuum to give 43.7 mg (44%) of the desired product. ESMS: Calcd. for $\text{C}_{64}\text{H}_{95}\text{N}_{19}\text{O}_{18}$, 1417.71; Found, 1418.8 $[\text{M}+\text{H}]^+$ 1. Analytical
 20 HPLC, Method 1B, R_t = 19.524 min, Purity = 73%.

Part C. Preparation of Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe} TFA salt.

25



To a solution of Boc-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe} (0.040 g, 0.0243 mmol) in
 30 methylene chloride (1 mL) was added trifluoroacetic acid (1 mL). The reaction mixture was stirred for 2 h, concentrated to an oil under high vacuum and triturated with diethyl ether. The product was filtered, washed with diethyl ether, and dried

under high vacuum to give 39.9 mg (100%) of the desired product. ESMS: Calcd. for $C_{59}H_{87}N_{19}O_{16}$, 1317.66; Found, 1318.9 $[M+H]^+$. Analytical HPLC, Method 1B, $R_t = 15.410$ min, Purity = 73%.


5

Part D. Preparation of [2-[[[5-[carbonyl]-2-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe}

10 To a solution of Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe} (0.030 g, 0.0183 mmol) in dimethylformamide (3 mL) was added triethylamine (7.6 μ L, 0.0549 mmol) and the reaction mixture was stirred for 5 min. 2-[[[5-[[[2,5-Dioxo-1-pyrrolidinyl]oxy]carbonyl]-2-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid, monosodium salt
15 (0.0096 g, 0.0220 mmol) was added, and the reaction mixture was stirred for 18 h, then concentrated to an oil under high vacuum. The oil was purified by Preparative HPLC Method 1 to give 11.0 mg (32%) of the desired product as a lyophilized
20 solid. ESMS: Calcd. for $C_{72}H_{96}N_{22}O_{20}S$, 1620.7; Found, 1620.1 ($M-H^+$). Analytical HPLC, Method 1B, $R_t = 16.753$ min, Purity = 91%.


Example 7

25 Synthesis of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Phe-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe}



30

Part A. Preparation of Phe-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-
cyclo{Lys-Arg-Gly-Asp-D-Phe}



5

A solution of Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-
cyclo{Lys-Arg-Gly-Asp-D-Phe} (23.4 mg, 0.014 mmol) and
triethylamine (7.8 μ L, 0.56 mmol) in DMF (2 mL) was stirred
for 5 min. To this was added Boc-Phe-OSu (5.1 mg, 0.014 mmol)
10 and the reaction mixture was stirred overnight at room
temperature under nitrogen. DMF was removed in vacuo, and the
resulting residue was dissolved in TFA (1.5 mL) and methylene
chloride (1.5 mL). The solution was stirred for 2 h and
concentrated in vacuo to provide 31 mg of the desired product
15 as the TFA salt. ESMS: Calcd. for $C_{68}H_{96}N_{20}O_{17}$, 1464.7;
Found, 1465.6 (M+H)+1. Analytical HPLC, Method 1B, R_t =
=15.48 min, Purity = 95%.

Part B. Preparation of [2-[[[5-[carbonyl]-2-
20 pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Phe-
Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe}


To a solution of Phe-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-
cyclo{Lys-Arg-Gly-Asp-D-Phe} (0.030 g, 0.016 mmol) in
25 dimethylformamide (2 mL) was added triethylamine (9 μ L, 0.064
mmol) and the reaction mixture was stirred for 5 min. 2-[[[5-
[[[2,5-Dioxo-1-pyrrolidinyl]oxy]carbonyl]-2-pyridinyl]-
hydrazono]methyl]-benzenesulfonic acid, monosodium salt
(0.0099 g, 0.0220 mmol) was added, and the reaction mixture
30 was stirred for 18 h, then concentrated under high vacuum.
The residue was purified by preparative RP-HPLC Method 1 to

give 7 mg (22%) of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for $C_{81}H_{105}N_{23}O_{21}S$, 1767.8; Found, 1768.8 ($M-H^+$). Analytical HPLC, Method 1B, $R_t = 17.68$ min, Purity = 99%.

5


Example 8

Synthesis of cyclo{Arg-Gly-Asp-D-Nal-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])}



10

Part A. Preparation of cyclo{Arg(Mtr)-Gly-Asp(OtBu)-D-Nal-Lys(Boc)}



15


The peptide Asp(OtBu)-D-Nal-Lys(Boc)-Arg(Mtr)-Gly was obtained by automated solid phase peptide synthesis using Fmoc chemistry. A 100 mL round bottom flask was charged with HBTU (349 mg, 0.92 mmol) and DMF (10 mL). The solution was stirred at 60 °C for 5 min. To this a solution of Asp(OtBu)-D-Nal-Lys(Boc)Arg(Mtr)-Gly (0.684 g) and Hunig's base (0.34 mL, 1.97 mmol.) in DMF (10 mL) was added and the solution stirred at 60 °C for 4 h under nitrogen. The solvent was then removed in vacuo and the residue was triturated with ethyl acetate. The

25

solids were filtered and washed with ethyl acetate (3 x 5 mL) and dried in vacuo to give the desired product (520 mg, 86%). ESMS: Calcd. for C₅₀H₇₁N₉O₁₂S, 1021.5; Found, 1022.5 [M+H]⁺. Analytical HPLC, Method 1A, R_t = 15.91 min (purity 99%).

5

Part B. Preparation of cyclo{Arg-Gly-Asp-D-Nal-Lys} bis TFA salt



10 A solution of cyclo{Arg(Mtr)-Gly-Asp(OtBu)-D-Nal-Lys(Boc)} (500 mg, 0.49 mmol), TFA (7 mL), triisopropylsilane (0.25 mL) and water (0.25 mL) was stirred at room temperature under nitrogen for 18 h. The solvents were removed in vacuo (over 3 h) and the residue triturated with diethyl ether to
15 give the desired product as the TFA salt (426 mg, 98%). ESMS: Calcd. for C₃₁H₄₃N₉O₇, 653.3; Found, 654.3 [M+H]⁺. Analytical HPLC, Method 1B, R_t = 13.30 min, Purity = 97%.


Part C. Preparation of cyclo{Arg-Gly-Asp-D-Nal-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])}

Cyclo{Arg-Gly-Asp-D-Nal-Lys} TFA salt (0.056 g, 0.064 mmol) was dissolved in DMF (2 mL). Triethylamine (27 μL, 0.19
25 mmol) was added, and after 5 min of stirring 2-[[[5-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]-hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.039 g, 0.089 mmol) was added. The reaction mixture was stirred overnight, under nitrogen, and then concentrated to an oil under high
30 vacuum. The oil was purified by Preparative HPLC Method 1 to give 49.3 mg (72%) of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for C₄₄H₅₂N₁₂O₁₁S, 956.4;

Found, 957.5 [M+H]⁺+1. Analytical HPLC, Method 1B, R_t = 16.19 min, Purity = 99%.

Example 9

- 5 Synthesis of [2-[[[5-[carbonyl]-2-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal}




10

Part A. Preparation of Boc-Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal}

- To a solution of cyclo{Lys-Arg-Gly-Asp-D-Nal} (0.052 g, 0.059 mmol) in dimethylformamide (2 mL) was added
 15 triethylamine (25 μ L). After stirring for 5 minutes Boc-Glu(OSu)-OSu (0.013 g, 0.029 mmol) was added. The reaction mixture was stirred under N₂ for 20 h, then concentrated to an oil under high vacuum and triturated with ethyl acetate. The
 20 product thus obtained was filtered, washed with ethyl acetate, and dried under high vacuum to give 35.2 mg of the desired product in crude form. ESMS: Calcd. for C₇₂H₉₉N₁₉O₁₈, 1517.7; Found, 760.1 [M+2H]⁺+2. Analytical HPLC, Method 1B, R_t = 21.07 min (65%).

25

Part B. Preparation of Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal}



To a solution of the crude Boc-Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal} (35.2 mg) in methylene chloride (1.5 mL) was added trifluoroacetic acid (1.5 mL). The reaction mixture was stirred for 2 h, concentrated to an oil under high vacuum and triturated with diethyl ether. The product was filtered, washed with diethyl ether, and dried under high vacuum to give 34.9 mg of the crude desired product (TFA salt). ESMS: Calcd. for C₆₇H₉₁N₁₉O₁₆, 1417.69; Found, 1418.7 [M+H]⁺+1. Analytical HPLC, Method 1B, R_t = 19.1 min, Purity = 62%.


Part C. Preparation of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal}

To a solution of Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal} (34.9 mg) in dimethylformamide (2 mL) was added triethylamine (10 μ L, 0.074 mmol) and the reaction mixture was stirred for 5 min. 2-[[[5-[(2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid, monosodium salt (15.2 mg, 0.0344 mmol) was added, and the reaction mixture was stirred for 18 h, then concentrated to an oil under high vacuum. The oil was purified by preparative RP-HPLC Method 1 to give 3 mg of the desired product (TFA salt). ESMS: Calcd. for C₈₀H₁₀₀N₂₂O₂₀S, 1720.7; Found, 1722.6 (M+H)⁺+1. Analytical HPLC, Method 1B, R_t = 19.78 min, Purity = 92%.

30

Example 10

Synthesis of cyclo{Arg-Gly-Asp-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid))-D-Val}



5

Part A. Preparation of cyclo{Arg(Tos)-Gly-Asp(OBzl)-Lys(Cbz)-D-Val}

The N-terminus Boc- protecting group of the peptide
 10 sequence Boc-Asp(OBzl)-Lys(Z)-D-Val-Arg(Tos)-Gly-Oxime resin
 was removed using standard deprotection (25% TFA in CH₂Cl₂).
 After eight washes with DCM, the resin was treated with 10%
 DIEA/DCM (2 x 10 min.). The resin was subsequently washed
 with DCM (x 5) and dried under high vacuum. The resin (1.3229
 15 g, 0.44 mmol/g) was then suspended in dimethylformamide (10
 mL). Glacial acetic acid (33.3 μ L, 0.582 mmol) was added, and
 the reaction was heated at 65 °C for 72 h. The resin was
 filtered, and washed with DMF (2 x 10 mL). The filtrate was
 concentrated to an oil under high vacuum. The resulting oil
 20 was triturated with ethyl acetate. The solid thus obtained
 was filtered, washed with ethyl acetate, dried under high
 vacuum, then purified by Preparative HPLC Method 2 to give
 93.0 mg of the desired product as a lyophilized solid. ESMS:
 Calcd. for C₄₅H₅₉N₉O₁₁S, 933.41; Found, 934.5 [M+H]⁺.
 25 Analytical HPLC, Method 1A, R_t = 14.078 min, Purity = 85%.

Preparative HPLC Method 2


Instrument: Rainin Rabbit; Dynamax software

Column: Vydac C-18 (21.2 mm x 25 cm)

Detector: Knauer VWM
 Flow Rate: 15ml/min
 Column Temp: RT
 Mobile Phase: A: 0.1% TFA in H₂O
 5 B: 0.1%TFA in ACN/H₂O (9:1)
 Gradient:

Time (min)	%A	%B
0	80	20
20	0	100
30	0	100
10 31	80	20

Part B. Preparation of cyclo{Arg-Gly-Asp-Lys-D-Val}



15


Cyclo{Arg(Tos)-Gly-Asp(OBzl)-Lys(Cbz)-D-Val} (0.080 g, 0.0856 mmol) was dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid (0.5 mL) was added dropwise, maintaining the temperature at -10 °C.
 20 Anisole (0.1 mL) was added and the reaction mixture was stirred at -10 °C for 3 h. Diethyl ether was added, the reaction mixture cooled to -50 °C and stirred for 30 mins. The crude product obtained was filtered, washed with ether, dried under high vacuum and purified by Preparative HPLC
 25 Method 1, to give 44.2 mg (66%) of the desired product as a lyophilized solid. ESMS: Calcd. for C₂₃H₄₁N₉O₇, 555.31; Found, 556.3 [M+H]⁺. Analytical HPLC, Method 1B, R_t = 8.959 min, Purity = 92%.

30 Part C. Preparation of cyclo{Arg-Gly-Asp-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Val}

To a solution of cyclo{Arg-Gly-Asp-Lys-D-Val} (0.036 g, 0.0459 mmol) in dimethylformamide (3 mL) was added triethylamine (19.2 μ L, 0.0138 mmol) and stirred for 5 min. Methyl sulfoxide was added (0.7 mL) followed by 2-[[[5-[[[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0243 g, 0.0551 mmol) and the reaction mixture stirred for 20 h. The reaction mixture was concentrated to an oil under high vacuum and purified by Preparative HPLC Method 1 to give 13.9 mg (31%) of the desired product as a lyophilized solid. HRMS: Calcd. for C₃₆H₅₀N₁₂O₁₁S +H, 859.3443; Found, 859.3503. Analytical HPLC, Method 1B, R_t = 13.479 min, Purity = 92%.

Example 11

Synthesis of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-cyclo{Lys-D-Val-Arg-Gly-Asp}




Part A. Preparation of Boc-Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-cyclo{Lys-D-Val-Arg-Gly-Asp}

To a solution of cyclo{Lys-D-Val-Arg-Gly-Asp} (0.400 g, 0.51 mmol) in dimethylformamide (7 mL) was added triethylamine (0.21 mL, 1.53 mmol). After stirring for 5 minutes Boc-Glu(OSu)-OSu (115 mg, 0.26 mmol) was added. The reaction mixture was stirred under N₂ for 20 h, then concentrated to an oil. The product thus obtained was partially purified by

preparative RP-HPLC to give 124 mg of product. ESMS: Calcd. for C₅₆H₉₅N₁₉O₁₈, 1321.71; Found, 1322.6 [M+H]⁺1.

Part B. Preparation of Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-
5 cyclo{Lys-D-Val-Arg-Gly-Asp}



To a solution of the impure Boc-Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-cyclo{Lys-D-Val-Arg-Gly-Asp} (0.124 g) in methylene chloride (5 mL) was added trifluoroacetic acid (5 mL). The reaction mixture was stirred for 2 h, concentrated to an oil under high vacuum and triturated with diethyl ether. The product was filtered, washed with diethyl ether, and dried under high vacuum to give 16.2 mg of the desired product after RP-HPLC (TFA salt). ESMS: Calcd. for C₅₁H₈₇N₁₉O₁₆, 1221.66; Found, 1222.6 [M+H]⁺1. Analytical HPLC, Method 1B, R_t = 11.43 min, Purity = 93%.


Part C. Preparation of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-
20 Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-cyclo{Lys-D-Val-Arg-Gly-Asp}

To a solution of Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-cyclo{Lys-D-Val-Arg-Gly-Asp} (0.016 g, 0.01 mmol) in dimethylformamide (2 mL) was added triethylamine (4.2 μL) and the reaction mixture was stirred for 5 min. 2-[[[5-[(2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt
25 (0.0063 g, 0.014 mmol) was added, and the reaction mixture was stirred for 18 h, then concentrated to an oil under high vacuum. The residue was purified by preparative RP-HPLC Method 1 to give the desired product (TFA salt). ESMS: Calcd.

for C₆₄H₉₆N₂₂O₂₀S, 1524.7; Found, 1525.7 (M+H)+1. Analytical HPLC, Method 1B, R_t = 13.20 min, Purity = 99%.

Example 12

- 5 Synthesis of {cyclo(Arg-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}




10

Part A: Preparation of cyclo(Arg(Tos)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly)

- The N-terminus Boc-protecting group of the peptide
 15 sequence Boc-Arg(Tos)-D-Val-D-Tyr(N-Cbz-aminopropyl)-D-Asp(OBzl)-Gly-Oxime resin was removed using standard deprotection (50% TFA in CH₂Cl₂). After washing with DCM (8x), the resin was neutralized with 10% DIEA/DCM (2 x 10 min). The resin was washed with DCM (5x) and dried under high
 20 vacuum overnight. The resin (1.08 g, 0.36 mmol/g) was then suspended in N,N-dimethylformamide (12 mL). Glacial acetic acid (67 mL, 1.16 mmol) was added and the reaction mixture was heated to 55 °C for 72 h. The resin was filtered and washed with DMF (3 x 10 mL). The filtrate was concentrated under
 25 high vacuum to give an oil. The resulting oil was triturated with ethyl acetate. The solid obtained was purified by reverse-phase HPLC (Vydac C18 column, 18 to 90% acetonitrile gradient containing 0.1% TFA, R_t=15.243 min) to afford 101 mg
 30 of a white powdered product (30%). ESMS: Calculated for C₄₄H₅₇N₉O₁₂S, 935.3847 Found 936.5 [M+H]+1.

Part B: Preparation of cyclo{Arg-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly}



5

The protected cyclic peptide cyclo{Arg(Tos)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly} (90 mg, 0.0961 mmol) was dissolved in trifluoroacetic acid (0.95 mL) and cooled to -10 °C in a dry ice/acetone bath. To this solution was added trifluoromethanesulfonic acid (0.1.16 mmol), followed by anisole (190 mL). The reaction mixture was stirred at -16 °C for 3 h. The dry ice/acetone bath was then cooled to -35 °C and cold ether (40 mL) was added to the solution. The mixture was stirred for 30 min at -35 °C, then cooled to -50 °C and stirred for another 30 min. The crude product was filtered, redissolved in water/acetonitrile (1/1), lyophilized, and purified by reverse-phase HPLC (Vydac C18 Column, 1.8 to 90% acetonitrile gradient containing 0.1% TFA, R_t =13.383 min) to generate 17 mg of the title product (27%). ESMS: Calculated for C₂₉H₄₅N₉O₈, 647.3391 Found 648.2 [M+H]⁺1.

20

Part C: Preparation of {cyclo{Arg-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}

25


A solution of cyclo{Arg-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly} (14 mg, 0.0216 mmol) in N,N-dimethylformamide (2 mL) was added triethylamine (15 mL, 0.108 mmol) and stirred at room temperature for 10 min. 2-[[[5-[(2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl-2-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid, monosodium salt (11 mg, 0.0260 mmol) was added, and the mixture was stirred for 18 h. The mixture was

30

concentrated under high vacuum and the residue was purified by reverse-phase HPLC (Vydac C18 Column, 1.8 to 90% acetonitrile gradient containing 0.1% TFA, R_t =16.264 min) to afford 10 mg of a white powdered product (49%). ESMS: Calculated for $C_{42}H_{54}N_{12}O_{12}S$, 950.3705 Found 951.3 [M+H]⁺1.

Example 13

Synthesis of cyclo{D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Phe-D-Asp-Gly-Arg}




Part A: Preparation of cyclo{D-Lys(Cbz)-D-Phe-D-Asp(OBzl)-Gly-Arg(Tos)}

The N-terminus Boc- protecting group of the peptide sequence Boc-Arg(Tos)-D-Lys(Cbz)-D-Phe-D-Asp(OBzl)-Gly-Oxime resin was removed using standard deprotection (25% TFA in CH_2Cl_2). After eight washes with DCM, the resin was treated with 10% DIEA/DCM (2 x 10 min.). The resin was subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.93 g, 0.44 mmol/g) was then suspended in dimethylformamide (15 mL). Glacial acetic acid (77 μ L) was added, and the reaction was heated to 60 °C for 72 h. The resin was filtered, and washed with DMF (2 x 10 mL). The filtrate was concentrated to an oil under high vacuum. The resulting oil was triturated with ethyl acetate. The solid thus obtained was filtered, washed with ethyl acetate, and dried under high vacuum to give the desired product which was then purified by preparative RP-HPLC (yield = 252 mg). ESMS: Calcd. for

C₄₉H₅₉N₉O₁₁S, 981.40; Found, 982.3 [M+H]⁺1. Analytical HPLC, Method 1A, R_t = 14.577 min.

Part B: Preparation of cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg} TFA salt



Cyclo{D-Lys(Cbz)-D-Phe-D-Asp(OBzl)-Gly-Arg(Tos)} (0.152 g, 0.155 mmol) was dissolved in trifluoroacetic acid (1.55 mL) and cooled to -16 °C. Trifluoromethanesulfonic acid (1.86 mL) was added dropwise, maintaining the temperature at -16 °C. Anisole (0.31 mL) was added and the reaction was stirred at -16 °C for 3 h. Diethyl ether was added, the reaction was cooled to -35 °C, and stirred for 20 min. The crude product was filtered, washed with diethyl ether, dried under high vacuum and purified by Preparative HPLC Method 1, to give 69 mg (~53%) of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for C₂₇H₄₁N₉O₇ +H, 604.3207; Found, 604.4. Analytical HPLC, Method 1B, R_t = 10.35 min, Purity = 93%.


Part C: Preparation of cyclo{D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Phe-D-Asp-Gly-Arg} TFA salt

Cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg} TFA salt (0.056 g, 0.0673 mmol) was dissolved in DMF (2 mL). Triethylamine (28 µL, 0.202 mmol) was added, and after 5 min of stirring 2-[[[5-[[[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.029 g, 0.0673 mmol) was added. The reaction mixture

was stirred for 70 h and then concentrated to an oil under high vacuum. The oil was purified by preparative HPLC Method 1 to give 14 mg (78%) of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for $C_{40}H_{50}N_{12}O_{11}S + H$,
 5 907.3521; Found, 907.3. Analytical HPLC, Method 1B, $R_t = 14.17$ min, Purity = 99%.

Example 14

Synthesis of [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-
 10 benzenesulfonic acid]-Glu(cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg})-
 cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg}




15 Part A. Preparation of Boc-Glu(cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg})-cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg}

To a solution of cyclo(D-Lys-D-Phe-D-Asp-Gly-Arg) (0.190 g, 0.228 mmol) in dimethylformamide (5 mL) was added
 20 triethylamine (95 μ L, 0.684 mmol). After stirring for 5 minutes Boc-Glu(OSu)-OSu (0.050 g, 0.114 mmol) was added. The reaction mixture was stirred under N_2 for 20 h, then concentrated to an oil under high vacuum and triturated with ethyl acetate. The product thus obtained was filtered, washed
 25 with ethyl acetate, and dried under high vacuum to give 172 mg of the desired product in crude form. ESMS: Calcd. for


C₆₄H₉₅N₁₉O₁₈, 1417.71; Found, 1418.7 [M+H]⁺+1. Analytical HPLC, Method 1B, R_t = 16.8 min.

Part B. Preparation of Glu(cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg})-
5 cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg}



To a solution of the crude Boc-Glu(cyclo{D-Lys-D-Phe-D-
10 Asp-Gly-Arg})-cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg} (0.172 g) in
methylene chloride (4.5 mL) was added trifluoroacetic acid
(4.5 mL). The reaction mixture was stirred for 2 h,
concentrated to an oil under high vacuum and triturated with
diethyl ether. The product was filtered, washed with diethyl
15 ether, and dried under high vacuum to give 38 mg of the
desired product after RP-HPLC as a lyophilized solid (TFA
salt). ESMS: Calcd. for C₅₉H₈₇N₁₉O₁₆, 1317.66; Found, 1318.9
[M+H]⁺+1. Analytical HPLC, Method 1B, R_t = 13.06 min, Purity =
20 93%.


Part C. Preparation of [2-[[[5-[carbonyl]-2-
pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{D-
Lys-D-Phe-D-Asp-Gly-Arg})-cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg}



To a solution of Glu(cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg})-
 cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg} (0.025 g, 0.015 mmol) in
 5 dimethylformamide (2 mL) was added triethylamine (6.3 μ L,
 0.045 mmol) and the reaction mixture was stirred for 5 min.
 2-[[[5-[[[(2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]-
 hydrazono]methyl]-benzenesulfonic acid, monosodium salt
 (0.0092 g, 0.0210 mmol) was added, and the reaction mixture
 10 was stirred for 18 h, then concentrated to an oil under high
 vacuum. The oil was purified by Preparative HPLC Method 1 to
 give 12.5 mg of the desired product as a lyophilized solid
 (TFA salt). ESMS: Calcd. for C₇₂H₉₆N₂₂O₂₀S, 1620.7; Found,
 1622.5 (M+H)⁺. Analytical HPLC, Method 1B, R_t = 14.62 min,
 15 Purity = 96%.

Example 15

Synthesis of cyclo{D-Phe-D-Lys([2-[[[5-[carbonyl]-2-
 pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Asp-Gly-
 20 Arg}




Part A. Preparation of cyclo{D-Phe-D-Lys(Cbz)-D-Asp(OBzl)-Gly-Arg(Tos)}

5

The N-terminus Boc- protecting group of the peptide sequence Boc-Arg(Tos)-D-Phe-D-Lys(Cbz)-D-Asp(OBzl)-Gly-Oxime resin was removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with DCM, the resin was treated with 10% DIEA/DCM (2 x 10 min.). The resin was subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.5 g, 0.44 mmol/g) was then suspended in dimethylformamide (12 mL). Glacial acetic acid (61 μ L) was added, and the reaction was heated to 60 °C for 72 h. The resin was filtered, and washed with DMF (2 x 10 mL). The filtrate was concentrated to an oil under high vacuum. The resulting oil was triturated with ethyl acetate. The solid thus obtained was filtered, washed with ethyl acetate, and dried under high vacuum to give the desired product (yield = 370 mg). ESMS: Calcd. for C₄₉H₅₉N₉O₁₁S, 981.40; Found, 982.4 [M+H]⁺. Analytical HPLC, Method 1A, R_t = 14.32 min (purity 60%).

Part B. Preparation of cyclo{D-Phe-D-Lys-D-Asp-Gly-Arg} bis TFA Salt




The crude cyclo{D-Phe-D-Lys(Cbz)-D-Asp(OBzl)-Gly-Arg(Tos)} (0.146 g) was dissolved in trifluoroacetic acid (1.5 mL) and cooled to -16 °C. Trifluoromethanesulfonic acid (1.8 mL) was added dropwise, maintaining the temperature at -16 °C. Anisole (0.3 mL) was added and the reaction was stirred at -16 °C for 3 h. Diethyl ether was added, the reaction was cooled to -35 °C, and stirred for 20 min. The crude product was filtered, washed with diethyl ether, dried under high vacuum and purified by Preparative HPLC Method 1, to give 100 mg of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for C₂₇H₄₁N₉O₇ + H, 604.3; Found, 604.3. Analytical HPLC, Method 1B, R_t = 10.25 min, Purity = 90%.

Part C. Preparation of cyclo{D-Phe-D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Asp-Gly-Arg}

Cyclo{D-Phe-D-Lys-D-Asp-Gly-Arg} TFA salt (0.090 g, 0.108 mmol) was dissolved in DMF (2 mL). Triethylamine (45 µL, 0.324 mmol) was added, and after 5 min of stirring 2-[[[5-[[[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.048 g, 0.108 mmol) was added. The reaction mixture was stirred for 70 h and then concentrated to an oil under high vacuum. The oil was purified by Preparative HPLC Method 1 to give 10 mg of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for C₄₀H₅₀N₁₂O₁₁S + H, 907.4; Found, 907.3. Analytical HPLC, Method 1B, R_t = 13.47 min, Purity = 89%.

Example 16

Synthesis of cyclo{N-Me-Arg-Gly-Asp-ATA-D-Lys([2-[[[5-
[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic
5 acid])}




Part A: Preparation of cyclo{N-Me-Arg(Tos)-Gly-Asp(OBzl)-ATA-D-Lys(Cbz)}

10

The N-terminus Boc-protecting group of the peptide sequence Boc-Asp(OBzl)-ATA-D-Lys(Z)-N-Me-Arg(Tos)-Gly-Oxime resin was removed using standard deprotection (50% TFA in CH₂Cl₂). After washing with DCM (8x), the resin was treated
15 with 10% DIEA/DCM (2 x 10 min). The resin was washed with DCM (5x) and dried under high vacuum overnight. The resin (1.24 g, 0.39 mmol/g) was then suspended in DMF (12 mL). Glacial acetic acid (67 mL, 1.16 mmol) was added and the reaction mixture was heated at 50 °C for 72 h. The resin was filtered
20 and washed with DMF (3 x 10 mL). The filtrate was concentrated under high vacuum to give an oil. The resulting oil was triturated with ethyl acetate. The solid obtained was purified by reverse-phase HPLC (Vydac C18 column, 18 to 90% acetonitrile gradient containing 0.1% TFA, R_t=14.129 min) to
25 afford 42 mg (9%) of the desired product as a lyophilized solid. ESMS: Calculated for C₄₆H₅₆N₁₀O₁₁S₂, 988.3571 Found 989.4 [M+H]⁺1.

Part B: Preparation of cyclo{N-Me-Arg-Gly-Asp-ATA-D-Lys}

30



Cyclo(N-Me-Arg(Tos)-Gly-Asp(OBzl)-ATA-D-Lys(Cbz)) (36 mg, 0.0364 mmol) was dissolved in trifluoroacetic acid (0.364 mL) and cooled to -10 °C in a dry ice/acetone bath. To this solution was added trifluoromethanesulfonic acid (0.437 mmol), followed by anisole (70 mL). The reaction mixture was stirred at -10 °C for 3 h. The dry ice/acetone bath was then cooled to -35 °C and cold ether (40 mL) was added to the solution. The mixture was stirred for 30 min at -35 °C, then cooled further to -50 °C and stirred for another 30 min. The crude product was filtered, redissolved in water/acetonitrile (1/1), and lyophilized to generate 35 mg of the title product (100%). ESMS: Calculated for C₂₄H₃₈N₁₀O₇S, 610.2646 Found 611.4 [M+H]⁺1.


Part C: Preparation of cyclo(N-Me-Arg-Gly-Asp-ATA-D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]))

To a solution of cyclo(N-Me-Arg-Gly-Asp-ATA-D-Lys) (31 mg, 0.051 mmol) in DMF (2 mL) was added triethylamine (28 mL, 0.204 mmol) and the reaction mixture stirred at room temperature for 10 min. 2-[[[5-[(2,5-Dioxo-1-pyrrolidinyl)-oxy]carbonyl-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (27 mg, 0.0612 mmol) was added, the mixture stirred for 18 h and then concentrated under high vacuum. The residue obtained was purified by reverse-phase HPLC (Shandon HS-BDS column, 3 to 10% acetonitrile, R_t=13.735 min) to afford 4 mg (8.8%) of the desired product as a lyophilized solid. ESMS: Calculated for C₃₇H₄₇N₁₃O₁₁S₂, 913.2959 Found 914.5 [M+H]⁺1.

Example 17

Synthesis of cyclo{Cit-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])}


5



Part A. Preparation of cyclo{Cit-Gly-Asp(OtBu)-D-Phe-Lys(Boc)}

10 The peptide Asp(OtBu)-D-Phe-Lys(Boc)-Cit-Gly was obtained by automated solid phase peptide synthesis using Fmoc chemistry (see general procedure). A 100 mL round bottom flask was charged with HBTU (271 mg, 0.71 mmol) and DMF (10 mL). The solution was stirred at 60 °C for 5 min. To this a
 15 solution of Asp(OtBu)-D-Phe-Lys(Boc)-Cit-Gly (0.456 g) and Hunig's base (0.27 mL, 1.53 mmol.) in DMF (10 mL) was added and the solution stirred at 60 °C for 4 h under nitrogen. The solvent was then removed in vacuo and the residue was trituated with ethyl acetate. The solids were filtered and
 20 washed with ethyl acetate (3 x 6 mL) and dried in vacuo to give the desired product (305 mg, 78%).
 ESMS: Calcd. for C₃₆H₅₆N₈O₁₀, 760.4; Found, 761.4 [M+H]⁺.
 Analytical HPLC, Method 1A, R_t = 11.8 min (purity 99%).

25 Part B. Preparation of cyclo{Cit-Gly-Asp(OtBu)-D-Phe-Lys(Boc)}




A solution of cyclo{Cit-Gly-Asp(OtBu)-D-Phe-Lys(Boc)} (287 mg, 0.38 mmol), TFA (6 mL), triisopropylsilane (0.25 mL) and water (0.25 mL) was stirred at room temperature under nitrogen for 4 h. The solvents were removed in vacuo (over 3 h) and the residue triturated with diethyl ether, filtered and washed with ether to give the desired product (315 mg) (TFA salt). ESMS: Calcd. for C₂₇H₄₀N₈O₈, 604.3; Found, 605.4 [M+H]⁺. Analytical HPLC, Method 1B, R_t = 9.6 min, Purity = 97%.

Part C. Preparation of cyclo{Cit-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]])}


Cyclo{Cit-Gly-Asp-D-Phe-Lys} TFA salt (0.044 g) was dissolved in DMF (2 mL). Triethylamine (22 µL, 0.156 mmol) was added, and after 5 min of stirring 2-[[[5-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.032 g, 0.073 mmol) was added. The reaction mixture was stirred overnight, under nitrogen, and then concentrated under high vacuum. The residue was purified by preparative RP-HPLC Method 1 to give 37 mg (70%) of the desired product as a lyophilized solid (TFA salt). ESMS: Calcd. for C₄₀H₄₉N₁₁O₁₂S, 907.3; Found, 908.4 [M+H]⁺. Analytical HPLC, Method 1B, R_t = 14.15 min, Purity = 99%.

Example 18A

Synthesis of tris(t-butyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid



Part A. Preparation of Phenylmethyl 2-(1,4,7,10-Tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)-acetate




A solution of tert-butyl (1,4,7,10-tetraaza-4,7-bis(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetate (0.922 g, 1.79 mmol), TEA (1.8 mL) and benzyl bromoacetate (0.86 mL, 5.37 mmol) in anhydrous DMF (24 mL) was stirred at ambient temperatures under a nitrogen atmosphere for 24 h. The DMF was removed under vacuum and the resulting oil was dissolved in EtOAc (300 mL). This solution was washed consecutively with water (2 x 50 mL) and saturated NaCl (50 mL), dried (MgSO₄), and concentrated to give the title compound as an amorphous solid (1.26 g). MS: m/e 663.5 [M+H].

Part B. Preparation of 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic acid

The product from Part A, above (165 mg, 0.25 mmol) was hydrogenolyzed over 10% Pd on carbon (50 mg) in EtOH (15 mL) at 60 psi for 24 h. The catalyst was removed by filtration through filter aid and washed with EtOH. The filtrates were concentrated to give the title compound as an amorphous solid (134 mg, 94%). MS: m/e 573.5 [M+H].


Example 18

Synthesis of 2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)-1-cyclododecyl)acetyl-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe}



Part A. Preparation of 2-(1,4,7,10-tetraaza-4,7,10-tris(t-butoxycarbonylmethyl)-1-cyclododecyl)acetyl-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe}

5



To a solution of tris(t-butyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (28 mg, 0.049 mmol) and Hunig's base (14 μ L) in DMF (2 mL) was added HBTU (17 mg, 0.0456 mmol) and the mixture stirred for 5 min. To this was added a solution of Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe} (54.1 mg, 0.0326 mmol) in DMF (1 mL) and the reaction mixture allowed to stir under nitrogen at room temperature for 4 h. The solvent was removed in vacuo and the residue purified by preparative RP-HPLC to give the product as a lyophilized solid (18.3 mg) (TFA salt). ESMS: Calcd. for $C_{87}H_{137}N_{23}O_{23}$, 1872.0; Found, 937.2 $[M+2H]^+$. Analytical HPLC, Method 1B, R_t = 19.98 min, Purity = 99%.


Part B. Preparation of 2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)-1-cyclododecyl)acetyl-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe}

5 A solution of 2-(1,4,7,10-tetraaza-4,7,10-tris(t-butoxycarbonylmethyl)-1-cyclododecyl)acetyl-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe} (18.3 mg, 8.71 mmol) in TFA (3 mL) was stirred at room temperature under nitrogen for 5 h. The solution was concentrated in vacuo and
10 the residue was purified by preparative RP-HPLC to give 8 mg (45%) of the desired product as the lyophilized solid (TFA salt). ESMS: Calcd. for $C_{75}H_{113}N_{23}O_{23}$, 1703.8; Found, 853.0 $[M+2H]^+$. Analytical HPLC, Method 1B, R_t = 13.13 min, Purity = 99%.

15

Example 19


Synthesis of cyclo{Arg-Gly-Asp-D-Phe-Lys(DTPA)}



20

To a solution of cyclo{Arg-Gly-Asp-D-Phe-Lys} (0.050 g, 0.0601 mmol) in DMF (2 mL) was added triethylamine (41.9 μ L, 0.301 mmol). This solution was added dropwise over 4 h to a solution of diethylenetriaminepentaacetic dianhydride (0.1074 g, 0.301 mmol) in DMF (2 mL) and methyl sulfoxide (2 mL). The
25 reaction mixture was then stirred for 16 h, concentrated to an oil under high vacuum and purified by Preparative HPLC Method 1 to give 29.9 mg (46%) of the desired product as a lyophilized solid. ESMS: Calcd. for $C_{41}H_{62}N_{12}O_{16}$, 978.4; Found, 977.5 $(M-H^+)$. Analytical HPLC, Method 1B, R_t = 11.916 min. Purity = 100%.

30

Example 20Synthesis of cyclo{Arg-Gly-Asp-D-Phe-Lys}₂ (DTPA)

5


The oil obtained in Example 9 upon purification by Preparative HPLC Method 1, also gave 21.5 mg (21%) of the title product as a lyophilized solid. ESMS: Calcd. for C₆₈H₁₀₁N₂₁O₂₂, 1563.7; Found, 1562.8 (M-H⁺). Analytical HPLC, Method 1B, R_t = 15.135 min, Purity = 93%.

10

Example 21

Synthesis of Cyclo{Arg-Gly-Asp-D-Tyr(N-DTPA-3-aminopropyl)-Val}

15



To a solution of cyclo{Arg-Gly-Asp-D-Tyr(3-aminopropyl)-Val} (0.050 g, 0.0571 mmol) in dimethylformamide (2 mL) was added triethylamine (39.8 μ L, 0.286 mmol). This solution was added dropwise over 5 h to a solution of diethylenetriamine-pentaacetic dianhydride (0.1020 g, 0.286 mmol) in methyl sulfoxide (2 mL). The reaction mixture was stirred for an


20

additional 18 h, then concentrated to an oil under high vacuum and purified by Preparative HPLC Method 1 to give 41.9 mg (65%) of the desired product as a lyophilized solid. ESMS: Calcd. for $C_{43}H_{66}N_{12}O_{17}$, 1022.5; Found, 1021.4 ($M-H^+$).

5 Analytical HPLC, Method 1B, R_t = 15.690 min, Purity = 96%.

Example 22

Synthesis of cyclo{Orn(d-N-2-ImidazolinyI)-Gly-Asp-D-Tyr(N-[2-
10 [[5-[carbonyl]-2-pyridinyI]hydrazono]methyl]-benzenesulfonic
acid]-3-aminopropyl)-Val}



Part A: Preparation of cyclo{Orn(d-N-1-Tos-2-ImidazolinyI)-
15 Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val}

The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Tyr(N-Cbz-aminopropyl)-Val-Orn(d-N-1-Tos-2-ImidazolinyI)-Gly-Oxime resin is removed using standard
20 deprotection (25% TFA in CH_2Cl_2). After eight washes with DCM, the resin is treated with 10% DIEA/DCM (2 x 10 min.). The resin is subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.75 g, 0.55 mmol/g) is then suspended in dimethylformamide (15 mL). Glacial acetic acid
25 (55.0 μ L, 0.961 mmol) is added, and the reaction mixture is heated at 50 °C for 72 h. The resin is filtered, and washed with DMF (2 x 10 mL). The filtrate is concentrated to an oil under high vacuum. The resulting oil is triturated with ethyl

acetate. The solid is filtered, washed with ethyl acetate, and is dried under high vacuum to obtain the desired product.

Part B: Preparation of cyclo{Orn(d-N-2-Imidazolinyl)-Gly-Asp-D-Tyr(3-aminopropyl)-Val}. Trifluoroacetic acid salt.


Cyclo{Orn(d-N-1-Tos-2-Imidazolinyl)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val} (0.146 mmol) is dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid (0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction mixture is stirred at -10 °C for 3 h. Diethyl ether is added, the reaction mixture cooled to -35 °C and then stirred for 30 min. The reaction mixture is cooled further to -50 °C and stirred for 30 min. The crude product is filtered, washed with diethyl ether, dried under high vacuum, and is purified by preparative HPLC to obtain the desired product.

Part C. Preparation of cyclo{Orn(d-N-2-Imidazolinyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}

Cyclo{Orn(d-N-2-Imidazolinyl)-Gly-Asp-D-Tyr(3-aminopropyl)-Val} trifluoroacetic acid salt (0.0228 mmol) is dissolved in DMF (1 mL). Triethylamine (0.0648 mmol) is added, and after 5 min of stirring 2-[[[5-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0274 mmol) is added. The reaction mixture is stirred for 1-2 days, and then concentrated to an oil under high vacuum. The oil is purified by preparative HPLC to obtain the desired product.

Example 23

Synthesis of cyclo{Lys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}



Part A: Preparation of cyclo{Lys(Tfa)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val}

5

The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Tyr(N-Cbz-aminopropyl)-Val-Lys(Tfa)-Gly-Oxime resin is removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with DCM, the resin is treated with 10% DIEA/DCM (2 x 10 min.). The resin is subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.75 g, 0.55 mmol/g) is then suspended in dimethylformamide (15 mL). Glacial acetic acid (55.0 μ L, 0.961 mmol) is added, and the reaction mixture is heated at 50 °C for 72 h. The resin is filtered, and washed with DMF (2 x 10 mL). The filtrate is concentrated to an oil under high vacuum. The resulting oil is triturated with ethyl acetate. The solid thus obtained is filtered, washed with ethyl acetate, and is dried under high vacuum to obtain the desired product.

10
15
20

Part B: Preparation of cyclo{Lys(Tfa)-Gly-Asp-D-Tyr(3-aminopropyl)-Val} Trifluoroacetic acid salt.

25 Cyclo{Lys(Tfa)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val} (0.146 mmol) is dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid (0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction mixture is stirred


at -10 °C for 3 h. Diethyl ether is added, the reaction mixture cooled to -35 °C and then stirred for 30 min. The reaction mixture is cooled further to -50 °C and stirred for 30 min. The crude product obtained is filtered, washed with diethyl ether, dried under high vacuum, and is purified by preparative HPLC to obtain the desired product.

Part C. Preparation of cyclo{Lys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}

Cyclo{Lys(Tfa)-Gly-Asp-D-Tyr(3-aminopropyl)-Val} trifluoroacetic acid salt (0.0228 mmol) is dissolved in DMF (1 mL). Triethylamine (0.0648 mmol) is added, and after 5 min of stirring 2-[[[5-[[[2,5-dioxo-1-pyrrolidinyl]oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0274 mmol) is added. The reaction mixture is stirred for 1-2 days, and then concentrated to an oil under high vacuum. The oil is treated with 20% piperidine in DMF, and the crude material is purified by preparative HPLC to obtain the desired product.

Example 24

Synthesis of cyclo{Cys(2-aminoethyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}



Part A: Preparation of cyclo{Cys(2-N-Tfa-aminoethyl)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val}

The N-terminus Boc- protecting group of the peptide
5 sequence Boc-Asp(OBzl)-D-Tyr(N-Cbz-aminopropyl)-Val-Cys(2-N-Tfa-aminoethyl)-Gly-Oxime resin is removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with DCM, the resin is treated with 10% DIEA/DCM (2 x 10 min.). The resin is subsequently washed with DCM (x 5) and dried
10 under high vacuum. The resin (1.75 g, 0.55 mmol/g) is then suspended in dimethylformamide (15 mL). Glacial acetic acid (55.0 µL, 0.961 mmol) is added, and the reaction mixture is heated at 50 °C for 72 h. The resin is filtered, and washed with DMF (2 x 10 mL). The filtrate is concentrated to an oil
15 under high vacuum. The resulting oil is triturated with ethyl acetate. The solid thus obtained is filtered, washed with ethyl acetate, and dried under high vacuum to obtain the desired product.

20 Part B: Preparation of cyclo{Cys(2-N-Tfa-aminoethyl)-Gly-Asp-D-Tyr(3-aminopropyl)-Val}. Trifluoroacetic acid salt.

Cyclo{Cys(2-N-Tfa-aminoethyl)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val} (0.146 mmol) is dissolved in
25 trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid (0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction mixture is stirred at -10 °C for 3 h. Diethyl ether is added, the reaction mixture cooled to -35 °C
30 and then stirred for 30 min. The reaction mixture is cooled further to -50 °C and stirred for 30 min. The crude product obtained is filtered, washed with diethyl ether, dried under high vacuum, and is purified by preparative HPLC to obtain the desired product.


35

Part C. Preparation of cyclo{Cys(2-aminoethyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}

Cyclo{Cys(2-N-Tfa-aminoethyl)-Gly-Asp-D-Tyr(3-aminopropyl)-Val} trifluoroacetic acid salt (0.0228 mmol) is dissolved in DMF (1 mL). Triethylamine (9.5 μ L, 0.0648 mmol) is added, and after 5 min of stirring 2-[[[5-[[[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0121 g, 0.0274 mmol) is added. The reaction mixture is stirred for 1-2 days, and then concentrated to an oil under high vacuum. The oil is treated with 20% piperidine in DMF, and the crude material is purified by preparative HPLC to obtain the desired product.

Example 25

Synthesis of cyclo{HomoLys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}



Part A: Preparation of cyclo{HomoLys(Tfa)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val}

The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Tyr(N-Cbz-aminopropyl)-Val-HomoLys(Tfa)-Gly-Oxime resin is removed using standard deprotection (25% TFA in CH_2Cl_2). After eight washes with DCM, the resin is treated with 10% DIEA/DCM (2 x 10 min.). The resin is subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.75 g, 0.55 mmol/g) is then

suspended in dimethylformamide (15 mL). Glacial acetic acid (55.0 μ L, 0.961 mmol) is added, and the reaction mixture is heated at 50 °C for 72 h. The resin is filtered, and washed with DMF (2 x 10 mL). The filtrate is concentrated to an oil under high vacuum. The resulting oil is triturated with ethyl acetate. The solid thus obtained is filtered, washed with ethyl acetate, and dried under high vacuum to obtain the desired product.

- 10 Part B: Preparation of cyclo{HomoLys(Tfa)-Gly-Asp-D-Tyr(3-aminopropyl)-Val}, Trifluoroacetic acid salt.

Cyclo{HomoLys(Tfa)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val} (0.146 mmol) is dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid (0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction mixture is stirred at -10 °C for 3 h. Diethyl ether is added, the reaction mixture cooled to -35 °C and then stirred for 30 min. The reaction mixture is cooled further to -50 °C and stirred for 30 min. The crude product obtained is filtered, washed with diethyl ether, dried under high vacuum, and is purified by preparative HPLC to obtain the desired product.


- 25 Part C. Preparation of cyclo{HomoLys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}

Cyclo{HomoLys(Tfa)-Gly-Asp-D-Tyr(3-aminopropyl)-Val} trifluoroacetic acid salt (0.0228 mmol) is dissolved in DMF (1 mL). Triethylamine (9.5 μ L, 0.0648 mmol) is added, and after 5 min of stirring 2-[[[5-[[[2,5-dioxo-1-pyrrolidinyl]oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0121 g, 0.0274 mmol) is added. The reaction mixture is stirred for 1-2 days, and then concentrated to an oil under high vacuum. The oil is treated with 20% piperidine in DMF, and the crude material is purified by preparative HPLC to obtain the desired product.

Example 26

Synthesis of cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Tyr(N-
[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-
benzenesulfonic acid]-3-aminopropyl)-Val}

5



Part A: Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-
Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val}

10

The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Tyr(N-Cbz-aminopropyl)-Val-Orn(d-N-Benzylcarbamoyl)-Gly-Oxime resin is removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with DCM, the resin is treated with 10% DIEA/DCM (2 x 10 min.). The resin is subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.75 g, 0.55 mmol/g) is then suspended in dimethylformamide (15 mL). Glacial acetic acid (55.0 μL, 0.961 mmol) is added, and the reaction mixture is heated at 50 °C for 72 h. The resin is filtered, and washed with DMF (2 x 10 mL). The filtrate is concentrated to an oil under high vacuum. The resulting oil is triturated with ethyl acetate. The solid thus obtained is filtered, washed with ethyl acetate, and dried under high vacuum to obtain the desired product.

15

20

25


Part B: Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Tyr(3-aminopropyl)-Val}. Trifluoroacetic acid salt.

- Cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val} (0.146 mmol) is dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C.
- 5 Trifluoromethanesulfonic acid (0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction mixture is stirred at -10 °C for 3 h. Diethyl ether is added, the reaction mixture cooled to -35 °C and then stirred for 30 min. The reaction mixture is cooled
- 10 further to -50 °C and stirred for 30 min. The crude product obtained is filtered, washed with diethyl ether, dried under high vacuum, and is purified by preparative HPLC to obtain the desired product.
- 15 Part C. Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}

- Cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Tyr(3-aminopropyl)-Val} trifluoroacetic acid salt (0.0228 mmol) is
- 20 dissolved in DMF (1 mL). Triethylamine (9.5 µL, 0.0648 mmol) is added, and after 5 min of stirring 2-[[[5-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0121 g, 0.0274 mmol)
- 25 is added. The reaction mixture is stirred for 1-2 days, and then concentrated to an oil under high vacuum. The oil is purified by preparative HPLC to obtain the desired product.

Example 27

- 30 Synthesis of cyclo{Dap(b-(2-benzimidazolylacetyl))-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}



Part A: Preparation of cyclo{Dap(b-(1-Tos-2-
benzimidazolylacetyl))-Gly-Asp(OBzl)-D-Tyr(N-Cbz-3-
5 aminopropyl)-Val}

The N-terminus Boc- protecting group of the peptide
sequence Boc-Asp(OBzl)-D-Tyr(N-Cbz-aminopropyl)-Val-Dap(b-(1-
Tos-2-benzimidazolylacetyl))-Gly-Oxime resin is removed using
10 standard deprotection (25% TFA in CH₂Cl₂). After eight washes
with DCM, the resin is treated with 10% DIEA/DCM (2 x 10
min.). The resin is subsequently washed with DCM (x 5) and
dried under high vacuum. The resin (1.75 g, 0.55 mmol/g) is
then suspended in dimethylformamide (15 mL). Glacial acetic
15 acid (55.0 μ L, 0.961 mmol) is added, and the reaction mixture
is heated at 50 °C for 72 h. The resin is filtered, and
washed with DMF (2 x 10 mL). The filtrate is concentrated to
an oil under high vacuum. The resulting oil is triturated
with ethyl acetate. The solid thus obtained is filtered,
20 washed with ethyl acetate, and dried under high vacuum to
obtain the desired product.

Part B: Preparation of cyclo{Dap(b-(2-benzimidazolylacetyl))-
Gly-Asp-D-Tyr(3-aminopropyl)-Val}. Trifluoroacetic acid salt.
25

Cyclo{Dap(b-(1-Tos-2-benzimidazolylacetyl))-Gly-
Asp(OBzl)-D-Tyr(N-Cbz-3-aminopropyl)-Val} (0.146 mmol) is
dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10


'C. Trifluoromethanesulfonic acid (0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction mixture is stirred at -10 °C for 3 h. Diethyl ether is added, the reaction mixture cooled to -35 °C and then stirred for 30 min. The reaction mixture is cooled further to -50 °C and stirred for 30 min. The crude product obtained is filtered, washed with diethyl ether, dried under high vacuum, and purified by preparative HPLC to obtain the desired product.

Part C. Preparation of cyclo{Dap(b-(2-benzimidazolylacetyl))-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val}

Cyclo{Dap(b-(2-benzimidazolylacetyl))-Gly-Asp-D-Tyr(3-aminopropyl)-Val} trifluoroacetic acid salt (0.0228 mmol) is dissolved in DMF (1 mL). Triethylamine (9.5 µL, 0.0648 mmol) is added, and after 5 min of stirring 2-[[[5-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0121 g, 0.0274 mmol) is added. The reaction mixture is stirred for 1-2 days, and then concentrated to an oil under high vacuum. The oil is purified by the method described below to obtain the desired product.

Example 28

Synthesis of cyclo{Orn(d-N-2-Imidazolinyl)-Gly-Asp-D-Phe-Lys(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]])}



Part A: Preparation of cyclo{Orn(d-N-1-Tos-2-Imidazolinyl)-
Gly-Asp(OBzl)-D-Phe-Lys(Cbz)}

5

The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Phe-Lys(Z)-Orn(d-N-1-Tos-2-Imidazolinyl)-Gly-Oxime resin is removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with
10 DCM, the resin is treated with 10% DIEA/DCM (2 x 10 min.). The resin is subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.75 g, 0.55 mmol/g) is then suspended in dimethylformamide (15 mL). Glacial acetic acid (55.0 μL, 0.961 mmol) is added, and the reaction mixture is
15 heated at 50 °C for 72 h. The resin is filtered, and washed with DMF (2 x 10 mL). The filtrate is concentrated to an oil under high vacuum. The resulting oil is triturated with ethyl acetate. The solid thus obtained is filtered, washed with ethyl acetate, and dried under high vacuum to obtain the
20 desired product.

Part B. Preparation of cyclo{Orn(d-N-2-Imidazolinyl)-Gly-Asp-D-Phe-Lys}

25

Cyclo{Orn(d-N-1-Tos-2-Imidazolinyl)-Gly-Asp(OBzl)-D-Phe-Lys(Cbz)} (0.204 mmol) is dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid


(0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction is stirred at -10 °C for 3 h. Diethyl ether is added, the reaction is cooled to -50 °C, and stirred for 1 h. The crude product is
5 filtered, washed with diethyl ether, dried under high vacuum and purified by preparative HPLC to obtain the desired product.

Part C. Preparation of cyclo{Orn(d-N-2-ImidazolinyI)-Gly-Asp-
10 D-Phe-Lys(N-[2-[[[5-[carbonyl]-2-pyridinyI]hydrazono]methyl]-benzenesulfonic acid]])}

Cyclo{Orn(d-N-2-ImidazolinyI)-Gly-Asp-D-Phe-Lys} TFA salt (0.0481 mmol) is dissolved in DMF (2 mL). Triethylamine (20.1
15 µL, 0.144 mmol) is added, and after 5 min of stirring 2-[[[5-[[[2,5-dioxo-1-pyrrolidinyI]oxy]carbonyl]-2-pyridinyI]hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.0254 g, 0.0577 mmol) is added. The reaction mixture is stirred for 20 h and then concentrated to an oil under high
20 vacuum. The oil is purified by preparative HPLC to obtain the desired product.

Example 29

Synthesis of cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Phe-
25 Lys(N-[2-[[[5-[carbonyl]-2-pyridinyI]hydrazono]methyl]-benzenesulfonic acid]])}



Part A: Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-
Asp(OBzl)-D-Phe-Lys(Cbz)}

5

The N-terminus Boc- protecting group of the peptide sequence Boc-Asp(OBzl)-D-Phe-Lys(Z)-Orn(d-N-Benzylcarbamoyl)-Gly-Oxime resin is removed using standard deprotection (25% TFA in CH₂Cl₂). After eight washes with DCM, the resin is
10 treated with 10% DIEA/DCM (2 x 10 min.). The resin is subsequently washed with DCM (x 5) and dried under high vacuum. The resin (1.75 g, 0.55 mmol/g) is then suspended in dimethylformamide (15 mL). Glacial acetic acid (55.0 μ L, 0.961 mmol) is added, and the reaction mixture is heated at 50
15 °C for 72 h. The resin is filtered, and washed with DMF (2 x 10 mL). The filtrate is concentrated to an oil under high vacuum. The resulting oil is triturated with ethyl acetate. The solid thus obtained is filtered, washed with ethyl acetate, and dried under high vacuum to obtain the desired
20 product.

Part B. Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-
Asp-D-Phe-Lys}

25

Cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp(OBzl)-D-Phe-Lys(Cbz)} (0.204 mmol) is dissolved in trifluoroacetic acid (0.6 mL) and cooled to -10 °C. Trifluoromethanesulfonic acid


(0.5 mL) is added dropwise, maintaining the temperature at -10 °C. Anisole (0.1 mL) is added and the reaction is stirred at -10 °C for 3 h. Diethyl ether is added, the reaction is cooled to -50 °C, and stirred for 1 h. The crude product is
 5 filtered, washed with diethyl ether, dried under high vacuum and purified by preparative HPLC to obtain the desired product.

Part C. Preparation of cyclo(Orn(d-N-Benzylcarbamoyl)-Gly-
 10 Asp-D-Phe-Lys(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]]))

Cyclo(Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Phe-Lys) TFA salt (0.0481 mmol) is dissolved in DMF (2 mL). Triethylamine
 15 (20.1 µL, 0.144 mmol) is added, and after 5 min of stirring 2-[[[5-[[[2,5-dioxo-1-pyrrolidinyl]oxy]carbonyl]-2-pyridinyl]hydrazono]-methyl]-benzenesulfonic acid, monosodium salt (0.0254 g, 0.0577 mmol) is added. The reaction mixture is stirred for 20 h and then concentrated to an oil under high
 20 vacuum. The oil is purified by preparative HPLC to obtain the desired product.

Example 30

Synthesis of cyclo(Lys-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-
 25 aminopropyl)-D-Asp-Gly}



30 Part A: Preparation of cyclo(Lys(Tfa)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly}

The N-terminus Boc-protecting group of the peptide sequence Boc-Lys(Tfa)-D-Val-D-Tyr(N-Cbz-aminopropyl)-D-Asp(OBzl)-Gly-Oxime resin is removed using standard deprotection (50% TFA in CH₂Cl₂). After washing with DCM (8x), the resin is neutralized with 10% DIEA/DCM (2 x 10 min). The resin is washed with DCM (5x) and dried under high vacuum overnight. The resin (1.0 g, about 0.36 mmol/g) is then suspended in N,N-dimethylformamide (12 mL). Glacial acetic acid (67 mL, 1.16 mmol) is added and the reaction mixture is heated to 55 °C for 72 h. The resin is filtered and washed with DMF (3 x 10 mL). The filtrate is concentrated under high vacuum to give an oil. The resulting oil is triturated with ethyl acetate. The desired product is purified by reverse-phase HPLC.

15

Part B: Preparation of cyclo(Lys-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly), Trifluoroacetic acid salt.

The protected cyclic peptide cyclo{Lys(Tfa)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly} (0.10 mmol) is dissolved in trifluoroacetic acid (0.95 mL) and cooled to -10 °C in a dry ice/acetone bath. To this solution is added trifluoromethanesulfonic acid (0.12 mmol), followed by anisole (190 mL). The reaction mixture is stirred at -16 °C for 3 h. The dry ice/acetone bath is then cooled to -35 °C and cold ether (40 mL) is added to the solution. The mixture is stirred for 30 min at -35 °C, then cooled to -50 °C and stirred for another 30 min. The crude product is filtered, redissolved in water/acetonitrile (1/1), lyophilized, and purified by reverse-phase HPLC to give the desired product.

Part C: Preparation of cyclo{Lys-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}

35


A solution of cyclo{Lys(Tfa)-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly} (0.0216 mmol) in N,N-dimethylformamide (2 mL) is added triethylamine (15 mL, 0.108 mmol) and stirred at room

temperature for 10 min. 2-[[[5-[[[2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl-2-pyridinyl]-hydrazono]methyl-benzenesulfonic acid, monosodium salt (0.0260 mmol) is added, and the mixture is stirred for 18 h. The mixture is concentrated under high vacuum, the oil is treated with 20% piperidine in DMF, and is again concentrated in vacuo. The residue is purified by reverse-phase HPLC to give the desired product.

10

Example 31

Synthesis of cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}



15

Part A: Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly}

20

The N-terminus Boc-protecting group of the peptide sequence Boc-Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(N-Cbz-aminopropyl)-D-Asp(OBzl)-Gly-Oxime resin is removed using standard deprotection (50% TFA in CH₂Cl₂). After washing with DCM (8x), the resin is neutralized with 10% DIEA/DCM (2 x 10 min). The resin is washed with DCM (5x) and dried under high vacuum overnight. The resin (1.0 g, about 0.36 mmol/g) is then suspended in N,N-dimethylformamide (12 mL). Glacial acetic acid (67 mL, 1.16 mmol) is added and the reaction mixture is heated to 55 °C for 72 h. The resin is filtered and washed with DMF (3 x 10 mL). The filtrate is concentrated under high vacuum to give an oil. The resulting oil is

30

trituated with ethyl acetate. The desired product is purified by reverse-phase HPLC.

Part B: Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly}, Trifluoroacetic acid salt.

The protected cyclic peptide cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly} (0.10 mmol) is dissolved in trifluoroacetic acid (0.95 mL) and cooled to -10 °C in a dry ice/acetone bath. To this solution is added trifluoromethanesulfonic acid (0.12 mmol), followed by anisole (190 mL). The reaction mixture is stirred at -16 °C for 3 h. The dry ice/acetone bath is then cooled to -35 °C and cold ether (40 mL) is added to the solution. The mixture is stirred for 30 min at -35 °C, then cooled to -50 °C and stirred for another 30 min. The crude product is filtered, redissolved in water/acetonitrile (1/1), lyophilized, and purified by reverse-phase HPLC to give the desired product.

20


Part C: Preparation of cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}

A solution of cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly} (0.0216 mmol) in N,N-dimethylformamide (2 mL) is added triethylamine (15 mL, 0.108 mmol) and stirred at room temperature for 10 min. 2-[[[5-[[[(2,5-Dioxo-1-pyrrolidinyl)oxy]carbonyl-2-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid, monosodium salt (0.0260 mmol) is added, and the mixture is stirred for 18 h. The mixture is concentrated under high vacuum and the residue is purified by reverse-phase HPLC to give the desired product.

35

Example 32

Synthesis of cyclo{Orn(d-N-2-Imidazoliny1)-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}



Part A: Preparation of cyclo{Orn(d-N-1-Tos-2-Imidazoliny1)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly}

The N-terminus Boc-protecting group of the peptide sequence Boc-Orn(d-N-1-Tos-2-Imidazoliny1)-D-Val-D-Tyr(N-Cbz-aminopropyl)-D-Asp(OBzl)-Gly-Oxime resin is removed using standard deprotection (50% TFA in CH₂Cl₂). After washing with DCM (8x), the resin is neutralized with 10% DIEA/DCM (2 x 10 min). The resin is washed with DCM (5x) and dried under high vacuum overnight. The resin (1.0 g, about 0.36 mmol/g) is then suspended in N,N-dimethylformamide (12 mL). Glacial acetic acid (67 mL, 1.16 mmol) is added and the reaction mixture is heated to 55 °C for 72 h. The resin is filtered and washed with DMF (3 x 10 mL). The filtrate is concentrated under high vacuum to give an oil. The resulting oil is triturated with ethyl acetate. The desired product is purified by reverse-phase HPLC.

Part B: Preparation of cyclo{Orn(d-N-2-Imidazoliny1)-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly}, Trifluoroacetic acid salt.

The protected cyclic peptide cyclo{Orn(d-N-1-Tos-2-Imidazoliny1)-D-Val-D-Tyr(N-Cbz-3-aminopropyl)-D-Asp(OBzl)-Gly} (0.10 mmol) is dissolved in trifluoroacetic acid (0.95 mL) and cooled to -10 °C in a dry ice/acetone bath. To this solution is added trifluoromethanesulfonic acid (0.12 mmol), followed by anisole (190 mL). The reaction mixture is stirred at -16 °C for 3 h. The dry ice/acetone bath is then cooled to -35 °C and cold ether (40 mL) is added to the

solution. The mixture is stirred for 30 min at -35 °C, then cooled to -50 °C and stirred for another 30 min. The crude product is filtered, redissolved in water/acetonitrile (1/1), lyophilized, and purified by reverse-phase HPLC to give the
5 desired product.

Part C: Preparation of cyclo{Orn(d-N-2-ImidazolinyI)-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyI]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}

10

A solution of cyclo{Orn(d-N-2-ImidazolinyI)-D-Val-D-Tyr(3-aminopropyl)-D-Asp-Gly} (0.0216 mmol) in N,N-dimethylformamide (2 mL) is added triethylamine (15 mL, 0.108 mmol) and stirred at room temperature for 10 min. 2-[[[5-
15 [[(2,5-Dioxo-1-pyrrolidinyI)oxy]carbonyl-2-pyridinyI]-hydrazono)methyl-benzenesulfonic acid, monosodium salt (0.0260 mmol) is added, and the mixture is stirred for 18 h. The mixture is concentrated under high vacuum and the residue is purified by reverse-phase HPLC to give the desired product.

20

Radiopharmaceutical Examples

The following procedures (A, B) describe the synthesis of radiopharmaceuticals of the present invention of the formula $^{99m}\text{Tc}(\text{VnA})(\text{tricine})(\text{phosphine})$, in which (VnA) represents the
25 vitronectin receptor antagonist compound bonded to the Tc through a diazenido (-N=N-) or hydrazido (=N-NH-) moiety. The diazenido or hydrazido moiety results from the reaction of the hydrazinonicotinamido group, present either as the free hydrazine or protected as a hydrazone, with the Tc-99m. The
30 other two ligands in the Tc coordination sphere are tricine and a phosphine.

Procedure A

Synthesis of Tc-99m Vitronectin Receptor Antagonist Complexes
35 of the Formula $^{99m}\text{Tc}(\text{VnA})(\text{tricine})(\text{phosphine})$ Using Stannous Reducing Agent

10-30 μg (0.2-0.4 mL) of a reagent of the present invention dissolved in saline or 50% aqueous ethanol, 40 mg (0.4 mL) of tricine in water, 1-7 mg (0.10-0.30 mL) of phosphine dissolved in water or ethanol, 25 μg (25 μL) $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ dissolved in 0.1 M HCl, 0-0.25 mL ethanol and 50-150 mCi $^{99\text{m}}\text{TcO}_4^-$ in saline were combined in a 10 cc vial. The kit was heated in a 100°C water bath for 10-20 minutes, then a 50 μL sample analyzed by HPLC Method 3. If necessary, the complex was purified by performing a 300-400 μL injection on the HPLC and collecting the fraction into a shielded flask. The collected fraction was evaporated to dryness, redissolved in saline containing 0-5 vol% Tween 80, and then re-analyzed using HPLC Method 3.

15 Procedure B

Synthesis of Tc-99m Vitronectin Receptor Antagonist Complexes of the Formula $^{99\text{m}}\text{Tc}(\text{VnA})(\text{tricine})(\text{TPPTS})$ Without Using Stannous Reducing Agent

20 To a lyophilized vial containing 4.84 mg TPPTS, 6.3 mg tricine, 40 mg mannitol and 0.25 mmol succinate buffer, pH 4.8, was added 0.2-0.4 mL (20-40 μg) of a reagent of the present invention dissolved in saline or 50% aqueous ethanol, 50-100 mCi $^{99\text{m}}\text{TcO}_4^-$ in saline, and additional saline to give
25 a total volume of 1.3-1.5 mL. The kit is heated in an 100°C water bath for 10-15 minutes, and a sample was then analyzed by HPLC Method 3. If necessary, the complex was purified by performing a 300-400 μL injection on the HPLC and collecting the fraction into a shielded flask. The collected fraction
30 was evaporated to dryness, redissolved in saline containing 0-5 vol% Tween 80, and then re-analyzed using HPLC Method 3.

Table 1. Analytical and Yield Data for $^{99\text{m}}\text{Tc}(\text{VnA})(\text{tricine})(\text{Phosphine})$ Complexes

Complex No.	Ex. No.	Reagent Ex. No.	Phosphine	% Yield	RT (min)
33	1		TPPTS	88	8.2
34	2		TPPTS	96	19.5

35	3	TPPTS	91	33.7
36	4	TPPTS	92	21.8
37	5	TPPTS	65	25.1
38	6	TPPTS	91	41.7
39	7	TPPTS	89	20.4
40	8	TPPTS	93	16.4
41	9	TPPTS	90	13.4
42	10	TPPTS	93	12.9
43	12	TPPMS	94	23.5
44	12	TPPDS	93	18.1
45	12	TPPTS	93	13.6
46	13	TPPTS	93	11.2
47	14	TPPTS	79	11.0
48	15	TPPTS	94	11.2
49	16	TPPTS	81	9.2
50	17	TPPTS	97	10.4

The following example describes the synthesis of radiopharmaceuticals of the present invention of the formula $^{99m}\text{Tc}(\text{VnA})(\text{tricine})(\text{L})$ (L = Imine-Nitrogen Containing Heterocycle), in which (VnA) represents the vitronectin receptor antagonist compound bonded to the Tc through a diazenido (-N=N-) or hydrazido (=N-NH-) moiety. The other two ligands in the Tc coordination sphere are tricine and an imine-nitrogen containing heterocycle.

10

Example 51

Synthesis of Tc-99m Vitronectin Receptor Antagonist Complex $^{99m}\text{Tc}(\text{VnA})(\text{tricine})(1,2,4\text{-triazole})$

15 30 μg of the Reagent of Example 1 (0.30 mL 50/50 EtOH/H₂O), 40 mg tricine (0.25 mL/H₂O), 8 mg 1, 2, 4-triazole (0.25 mL/H₂O), 25 μg SnCl₂ (25 μL /0.1 N HCl), 0.50 mL water and 0.20 mL 50 \pm 5 mCi $^{99m}\text{TcO}_4^-$ were combined in a shielded 10 cc vial and heated at 100 °C for 10 minutes. 50 μL of the kit
20 contents were analyzed by HPLC using Method listed below. The

product eluted at a retention time of 8.33 min and had a radiochemical purity of 88.1%.

Reagents of the present invention comprised of either a DOTA (Example 18), DTPA monoamide (Examples 19 and 20) or DTPA bisamide chelator (Example 21) readily form complexes with metal ions of elements 31, 39, 49, and 58-71. The following examples demonstrate the synthesis of complexes with ^{153}Sm , ^{177}Lu , and ^{90}Y , beta particle emitting isotopes used in radiopharmaceutical therapy, and ^{111}In , a gamma emitting isotope used in radiopharmaceutical imaging agents. In both types of complexes, the metal ion is bound to the DOTA, DTPA monoamide or DTPA bisamide chelator moiety of the reagents.

Examples 52 and 53

Synthesis of Y-90 and Lu-177 DOTA-Containing Vitronectin Antagonist Complexes

To a clean sealed 10 mL vial was added 0.5 mL of the reagent of Example 18 (200 $\mu\text{g/mL}$ in 0.25 M ammonium acetate buffer, pH 7.0), followed by 0.05 - 0.1 mL of gentisic acid (sodium salt, 10 mg/mL in 0.25 M ammonium acetate buffer, pH 7.0) solution, 0.3 mL of 0.25 M ammonium acetate buffer (pH 7.0), and 0.05 mL of $^{177}\text{LuCl}_3$ solution or $^{90}\text{YCl}_3$ solution (100 - 200 mCi/mL) in 0.05 N HCl. The resulting mixture was heated at 100 °C for 35 min. After cooling to room temperature, a sample of the resulting solution was analyzed by radio-HPLC and ITLC. The complex of Example 53 was analyzed by mass spectroscopy (Found $[\text{M}+\text{H}^+] = 1877.6$, Calcd. 1875.8 for $\text{C}_{75}\text{H}_{110}\text{N}_{23}\text{O}_{23}\text{Lu}$) which confirmed identity.

Example 54

Synthesis of a ^{111}In DOTA-Containing Vitronectin Antagonist Complex

35

To a lead shielded 300 μL autosampler vial was added 50 μL of gentisic acid (10 mg/mL in 0.1 M ammonium acetate buffer, pH 6.75) solution, followed by 100 μL of the reagent

of Example 18 (200 µg/mL in 0.2 M ammonium acetate buffer, pH 5.0), and 50 µL of $^{111}\text{InCl}_3$ solution (10 mCi/mL) in 0.04 N HCl. The pH of the reaction mixture was about 4.0. The solution was heated at 100 °C for 25 min. A sample of the resulting solution was analyzed by radio-HPLC and ITLC.

Table 1A: Analytical and Yield Data for Y-90, In-111, and Lu-177 Complexes of DOTA-Conjugated Vitronectin Receptor Antagonists.

Complex Ex. No.	Reagent Ex. No.	Isotope	%Yield	HPLC Ret. Time (min)
52	18	Y-90	96	16.5
53	18	Lu-177	96	16.5
54	18	In-111	95	16.5

10

Examples 55 and 56

Synthesis of In-111 DTPA-monoamide or DTPA-bisamide Containing Vitronectin Antagonist Complexes

0.2 mL of $^{111}\text{InCl}_3$ (1.7 mCi) in 0.1 N HCl, 0.2 mL of 1.0 M ammonium acetate buffer (pH 6.9) and 0.1 ml of the reagent of the present invention dissolved in water were combined in a 10cc glass vial and allowed to react at room temperature for 30 min. The reaction mixture was analyzed by HPLC Method 3.

20

Table 2. Analytical and Yield Data for ^{111}In Complexes

Complex Ex. No.	Reagent Ex. No.	%Yield	HPLC Ret. Time (min)
55	19	86	11.1
56	20	96	18.8

Examples 57-59

Synthesis of Sm-153 Vitronectin Antagonist Complexes

25

0.25 mL of a $^{153}\text{SmCl}_3$ stock solution (54 mCi/µmol Sm, 40 mCi/mL) in 0.1 N HCl was combined with the reagent of the present invention (50-fold molar excess) dissolved in 1 N ammonium acetate buffer in a 10cc glass vial. The reaction was

allowed to proceed at room temperature for ~30 min and was then analyzed by ITLC and HPLC (Method 3). If necessary, the complex was purified by performing a 300-400 μ L injection on the HPLC and collecting the fraction into a shielded flask.




- 5 The collected fraction was evaporated to dryness, redissolved in saline, and then re-analyzed using HPLC Method 3.

Table 3. Analytical and Yield Data for ^{153}Sm Complexes

Complex Ex. No.	Reagent Ex. No.	%Yield	HPLC Ret. Time (min)
57	19	91	11.7
58	20	84	13.1
59	21	96	16.9

- 10 The non-radioactive (naturally occurring) samarium analog of the Radiopharmaceutical of Example 59 was prepared by combining 3.3. mg (2.9 μ mol) of the Reagent of Example 21 dissolved in 2 mL of 1 M ammonium acetate buffer, pH 7, and 0.29 mL of 0.01 M solution of SmCl_3 in 0.1 N HCl. The
- 15 reaction was allowed to proceed for ~ 5 h at room temperature and then the product was isolated by HPLC Method 3. The volatiles were removed by lyophilization. The identity of the complex was confirmed by mass spectroscopy. (API-ESMS:Found $[\text{M}+2\text{H}^+ = 1172.4$, Calcd. 1172.9 for $\text{C}_{43}\text{H}_{64}\text{N}_{12}\text{O}_{17}\text{Sm}$] A stock
- 20 solution of the complex was made in water and its concentration determined by ICP analysis for use in determining the binding affinity of the complex for the vitronectin receptor $\alpha_v\beta_3$.

- 25 The structures of representative In-111 (Example 56), Y-90 (Example 52) and Sm-153 (Example 59) radiopharmaceuticals of the present invention are shown below.



5

Examples 60-62

Synthesis of Lu-177 Vitronectin Antagonist Complexes

- 10 5×10^{-9} mol of a reagent of the present invention was dissolved in 1.0 mL of 0.1 N acetate buffer, pH 6.8. 1×10^{-9} mol of Lu-177 (40 μ l, 3 mCi) dissolved in 0.1 N HCl was added and the reaction allowed to proceed at room temperature for 30-45 min. The reaction mixtures were analyzed by HPLC Method
- 15 3.

Table 4. Analytical and Yield Data for ^{177}Lu Complexes

Complex Ex. No.	Reagent Ex. No.	%Yield	HPLC Ret. Time (min)
60	19	98	11.0
61	20	98	15.6
62	21	98	11.7

Example 63

5 The gadolinium complex of the reagent of Example 21 was prepared according to the following procedure. 3-3.5 mg of the reagent was dissolved in 2 mL 1 M ammonium acetate buffer at pH 7.0, and one equivalent $\text{Gd}(\text{NO}_3)_3$ solution (0.02 M in water) was added to it. The reaction mixture was allowed to

10 stay at room temperature for 3-5 hours and the product was isolated by HPLC Method 4. The fraction containing the complex was lyophilized and dissolved in 1 mL H_2O resulting in a solution approximately 2 mM in Gd as determined by ICP analysis. The identity of the complex was confirmed by mass

15 spectroscopy. (API-ESMS: Found $[\text{M}+2\text{H}^+] = 1176.9$, Calcd. 1176.2 for $\text{C}_{43}\text{H}_{64}\text{N}_{12}\text{O}_{17}\text{Gd}$).


The following examples describe the synthesis of ultrasound contrast agents of the present invention comprised

20 of targeting moieties for tumor neovasculature that are $\alpha_v\beta_3$ receptor antagonists.

Example 64

Part A. Synthesis of 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-(cyclo(Arg-Gly-Asp-D-Phe-Lys)-dodecane-1,12-dione

25




A solution of disuccinimidyl dodecane-1,12-dioate (0.424 g, 1 mmol), 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine (1.489 g, 1 mmol) and cyclo(Arg-Gly-Asp-D-Phe-Lys) TFA salt (0.831 g, 1 mmol) in 25 ml chloroform is stirred for 5 min. Sodium carbonate (1 mmol) and sodium sulfate (1 mmol) are added and the solution is stirred at room temperature under nitrogen for 18 h. DMF is removed in vacuo and the crude product is purified to obtain the title compound.

Part B. Preparation of Contrast Agent Composition

The Synthesis of 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-(cyclo(Arg-Gly-Asp-D-Phe-Lys)-dodecane-1,12-dione is admixed with three other lipids, 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine in relative amounts of 1 wt. %:6 wt. %:54 wt. %:41 wt. %. An aqueous solution of this lipid admixture (1 mg/mL), sodium chloride (7 mg/mL), glycerin (0.1 mL/mL), propylene glycol (0.1 mL/mL), at pH 6-7 is then prepared in a 2 cc glass vial. The air in the vial is evacuated and replaced with perfluoropropane and the vial is sealed. The ultrasound contrast agent composition is completed by agitating the sealed vial in a dental amalgamator for 30-45 sec. to form a milky white solution.

Example 65

Part A. Preparation of (ω -amino-PEG₃₄₀₀- α -carbonyl)-
cyclo(Arg-Gly-Asp-D-Phe-Lys)




5

To a solution of N-Boc- ω -amino-PEG₃₄₀₀- α -carboxylate
succinimidyl ester (1 mmol) and cyclo(Arg-Gly-Asp-D-Phe-Lys) (1
mmol) in DMF (25 mL) is added triethylamine (3 mmol). The
reaction mixture is stirred under nitrogen at room temperature
overnight and the solvent is removed in vacuo. The crude
10 product is dissolved in 50% trifluoroacetic
acid/dichloromethane and is stirred for 4 h. The volatiles
are removed and the title compound is isolated as the TFA salt
via trituration in diethyl ether.

15

Part B. Preparation of 1-(1,2-Dipalmitoyl-sn-glycero-3-
phosphoethanolamino)-12-(ω -amino-PEG₃₄₀₀- α -carbonyl)-
cyclo(Arg-Gly-Asp-D-Phe-Lys)-Dodecane-1,12-Dione



20

A solution of disuccinimidyl dodecanoate (1 mmol), 1,2-
dipalmitoyl-sn-glycero-3-phosphoethanolamine (1 mmol) and (ω -
amino-PEG₃₄₀₀- α -carbonyl)-cyclo(Arg-Gly-Asp-D-Phe-Lys) TFA
25 salt (1 mmol) in 25 ml chloroform is stirred for 5 min.

Sodium carbonate (1 mmol) and sodium sulfate (1 mmol) are added and the solution is stirred at room temperature under nitrogen for 18 h. DMF is removed in vacuo and the crude product is purified to obtain the title compound.

5


Part C. Preparation of Contrast Agent Composition

The 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-((ω -amino-PEG₃₄₀₀- α -carbonyl)-cyclo(Arg-Gly-Asp-D-Phe-Lys))-Dodecane-1,12-Dione is admixed with three other lipids, 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine in relative amounts of 1 wt. %:6 wt. %:54 wt. %:41 wt. %. An aqueous solution of this lipid admixture (1 mg/mL), sodium chloride (7 mg/mL), glycerin (0.1 mL/mL), propylene glycol (0.1 mL/mL), at pH 6-7 is then prepared in a 2 cc glass vial. The air in the vial is evacuated and replaced with perfluoropropane and the vial is sealed. The ultrasound contrast agent composition is completed by agitating the sealed vial in a dental amalgamator for 30-45 sec. to form a milky white solution.

25


Example 66

Part A. Preparation of Synthesis of (ω -amino-PEG₃₄₀₀- α -carbonyl)-Glu-(cyclo(Arg-Gly-Asp-D-Phe-Lys))₂



To a solution of N-Boc- ω -amino-PEG₃₄₀₀- α -carboxylate
 succinimidyl ester (1 mmol) and Glu-(cyclo(Arg-Gly-Asp-D-Phe-
 5 Lys))₂ (1 mmol) in DMF (25 mL) is added triethylamine (3
 mmol). The reaction mixture is stirred under nitrogen at room
 temperature overnight and the solvent is removed in vacuo.
 The crude product is dissolved in 50% trifluoroacetic
 acid/dichloromethane and is stirred for 4 h. The volatiles
 10 are removed and the title compound is isolated as the TFA salt
 via trituration in diethyl ether.

Part B. Preparation of 1-(1,2-Dipalmitoyl-sn-glycero-3-
 phosphoethanolamino)-12-((ω -amino-PEG₃₄₀₀- α -carbonyl)-Glu-
 15 (cyclo(Arg-Gly-Asp-D-Phe-Lys))₂)-Dodecane-1,12-Dione



A solution of disuccinimidyl dodecanoate (1 mmol), 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine (1 mmol) and (ω -amino-PEG₃₄₀₀- α -carbonyl)-Glu-(cyclo(Arg-Gly-Asp-D-Phe-Lys))₂ TFA salt (1 mmol) in 25 ml chloroform is stirred for 5 min. Sodium carbonate (1 mmol) and sodium sulfate (1 mmol) are added and the solution is stirred at room temperature under nitrogen for 18 h. DMF is removed in vacuo and the crude product is purified to obtain the title compound.

Part C. Preparation of Contrast Agent Composition

The 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-((ω -amino-PEG₃₄₀₀- α -carbonyl)-Glu-(cyclo(Arg-Gly-Asp-D-Phe-Lys))₂)-Dodecane-1,12-Dione is admixed with three other lipids, 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine in relative amounts of 1 wt. %:6 wt. %:54 wt. %:41 wt. %. An aqueous solution of this lipid admixture (1 mg/mL), sodium chloride (7 mg/mL), glycerin (0.1 mL/mL), propylene glycol (0.1 mL/mL), at pH 6-7 is then prepared in a 2 cc glass vial. The air in the vial is evacuated and replaced with perfluoropropane and the vial is

sealed. The ultrasound contrast agent composition is completed by agitating the sealed vial in a dental amalgamator for 30-45 sec. to form a milky white solution.

5 Analytical Methods

HPLC Method 3

Column: Zorbax C18, 25 cm x 4.6 mm or Vydac C18, 25 cm x 4.6 mm

Column Temperature: ambient

10 Flow: 1.0 mL/min

Solvent A: 10 mM sodium phosphate buffer pH 6

Solvent B: 100% Acetonitrile

Detector: sodium iodide (NaI) radiometric probe or beta detector

15

Gradient A (Exs. 33, 51)

t (min)	0	20	30	31	40
%B	0	75	75	0	0

20 Gradient B (Exs. 39, 40, 43, 44, 45, 46, 48, 50)

t (min)	0	20	30	31	35	36	40
%B	0	25	25	75	75	0	0

Gradient C (Examples 34, 35, 36, 37, 38, 42):

25

t (min)	0	40	41	46	47	55
%B	0	35	75	75	0	0

Gradient D (Ex. 49)

30

t (min)	0	20	30	31	40
%B	0	25	25	0	0

Gradient E (Exs. 55, 56):

t (min)	0	20	21	30	31	40
%B	0	20	50	50	0	0

35

Gradient F (Exs. 57, 58):

t (min)	0	15	16	25	26	35
%B	0	20	75	75	0	0

Gradient G (Ex. 59):

t (min)	0	20	21	30	31	40
%B	0	20	75	75	0	0

5

Gradient H (Exs. 60, 61, 62):

t (min)	0	15	16	21	22	40
%B	0	20	50	50	0	0

10 Gradient I (Exs. 52, 53, 54)

t (min)		0	20	21	30	31	40
% Solvent B		5	20	60	60	5	5

Gradient J (Ex. 41)

15 t (min)		0	20	30	31	40
% Solvent B		0	50	50	0	0

Gradient K (Ex. 47)

20 t (min)		0	20	21	30	31	40
% Solvent B		10	20	60	60	10	10

HPLC Method 4

Column: Zorbax C18, 25 cm x 4.6 mm

Flow: 1.0 mL/min

25 Solvent A: 10 mM ammonium acetate

Solvent B: 100% methanol

Gradient:

t (min)	0	23	26	27
%B	8	100	100	8

30 UV Detection

ITLC Method

Gelman ITLC-SG strips (2 cm x 7.5 cm)

Solvent System: 1:1 acetone:saline

35 Detection using a Bioscan System 200.

UTILITY

The pharmaceuticals of the present invention are useful for imaging angiogenic tumor vasculature in a patient or for treating cancer in a patient. The radiopharmaceuticals of the present invention comprised of a gamma emitting isotope are useful for imaging of pathological processes involving angiogenic neovasculature, including cancer, diabetic retinopathy, macular degeneration, restenosis of blood vessels after angioplasty, and wound healing. Diagnostic utilities also include imaging of unstable coronary syndromes (e.g., unstable coronary plaque). The radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope are useful for treatment of pathological processes involving angiogenic neovasculature, by delivering a cytotoxic dose of radiation to the locus of the angiogenic neovasculature. The treatment of cancer is affected by the systemic administration of the radiopharmaceuticals resulting in a cytotoxic radiation dose to tumors.

The compounds of the present invention comprised of one or more paramagnetic metal ions selected from gadolinium, dysprosium, iron, and manganese, are useful as contrast agents for magnetic resonance imaging (MRI) of pathological processes involving angiogenic neovasculature.

The compounds of the present invention comprised of one or more heavy atoms with atomic number of 20 or greater are useful as X-ray contrast agents for X-ray imaging of pathological processes involving angiogenic neovasculature.

The compounds of the present invention comprised of an echogenic gas containing surfactant microsphere are useful as ultrasound contrast agents for sonography of pathological processes involving angiogenic neovasculature.

Representative compounds of the present invention were tested in the following in vitro and in vivo assays and models and were found to be active.

35

Immobilized Human Placental $\alpha_v\beta_3$ Receptor Assay

The assay conditions were developed and validated using [I-125]vitronectin. Assay validation included Scatchard

format analysis (n=3) where receptor number (Bmax) and Kd (affinity) were determined. Assay format is such that compounds are preliminarily screened at 10 and 100 nM final concentrations prior to IC50 determination. Three standards (vitronectin, anti- $\alpha_v\beta_3$ antibody, LM609, and anti- $\alpha_v\beta_5$, P1F6) and five reference peptides have been evaluated for IC50 determination. Briefly, the method involves immobilizing previously isolated receptors in 96 well plates and incubating overnight. The receptors were isolated from normal, fresh, non-infectious (HIV, hepatitis B and C, syphilis, and HTLV free) human placenta. The tissue was lysed and tissue debris removed via centrifugation. The lysate was filtered. The receptors were isolated by affinity chromatography using the immobilized $\alpha_v\beta_3$ antibody. The plates are then washed 3x with wash buffer. Blocking buffer is added and plates incubated for 120 minutes at room temperature. During this time, compounds to be tested and [I-125]vitronectin are premixed in a reservoir plate. Blocking buffer is removed and compound mixture pipetted. Competition is carried out for 60 minutes at room temperature. Unbound material is then removed and wells are separated and counted via gamma scintillation.

Other Receptor Binding Assays

Whole cell assays for the determination of the binding affinity of pharmaceuticals of the present invention for the VEGF receptors, Flk-1/KDR and Flt-1, are described in Ortega, et. al., Amer. J. Pathol., 1997, 151, 1215-1224, and Dougher, et. al., Growth Factors, 1997, 14, 257-268. An in vitro assay for determining the affinity of pharmaceuticals of the present invention for the bFGF receptor is described in Yayon, et. al., Proc. Natl. Acad. Sci USA, 1993, 90, 10643-10647. Gho et. al., Cancer Research, 1997, 57, 3733-40, describe assays for angiogenin receptor binding peptides. Senger, et. al., Proc. Natl. Acad. Sci USA, 1997, 94, 13612-13617 describe assays for antagonists of the integrins $\alpha_1\beta_1$ and $\alpha_2\beta_1$. U.S. 5,536,814 describes assays for compounds that bind to the integrin $\alpha_5\beta_1$.

Oncomouse® Imaging

The study involves the use of the c-Neu Oncomouse® and FVB mice simultaneously as controls. The mice are anesthetized with sodium pentobarbital and injected with approximately 0.5 mCi of radiopharmaceutical. Prior to injection, the tumor locations on each Oncomouse® are recorded and tumor size measured using calipers. The animals are positioned on the camera head so as to image the anterior or posterior of the animals. 5 Minute dynamic images are acquired serially over 2 hours using a 256x256 matrix and a zoom of 2x. Upon completion of the study, the images are evaluated by circumscribing the tumor as the target region of interest (ROI) and a background site in the neck area below the carotid salivary glands.

This model can also be used to assess the effectiveness of the radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope. The radiopharmaceuticals are administered in appropriate amounts and the uptake in the tumors can be quantified either non-invasively by imaging for those isotopes with a coincident imageable gamma emission, or by excision of the tumors and counting the amount of radioactivity present by standard techniques. The therapeutic effect of the radiopharmaceuticals can be assessed by monitoring the rate of growth of the tumors in control mice versus those in the mice administered the radiopharmaceuticals of the present invention.

This model can also be used to assess the compounds of the present invention comprised of paramagnetic metals as MRI contrast agents. After administration of the appropriate amount of the paramagnetic compounds, the whole animal can be placed in a commercially available magnetic resonance imager to image the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of heavy atoms as X-ray contrast agents. After administration of the appropriate

amount of the X-ray absorbing compounds, the whole animal can be placed in a commercially available X-ray imager to image the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of an echogenic gas containing surfactant microsphere as ultrasound contrast agents. After administration of the appropriate amount of the echogenic compounds, the tumors in the animal can be imaging using an ultrasound probe held proximate to the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

15

Rabbit Matrigel Model

This model was adapted from a matrigel model intended for the study of angiogenesis in mice. Matrigel (Becton & Dickinson, USA) is a basement membrane rich in laminin, collagen IV, entactin, HSPG and other growth factors. When combined with growth factors such as bFGF [500 ng/ml] or VEGF [2 µg/ml] and injected subcutaneously into the mid-abdominal region of the mice, it solidifies into a gel and stimulates angiogenesis at the site of injection within 4-8 days. In the rabbit model, New Zealand White rabbits (2.5-3.0 kg) are injected with 2.0 ml of matrigel, plus 1 µg bFGF and 4 µg VEGF. The radiopharmaceutical is then injected 7 days later and the images obtained.

This model can also be used to assess the effectiveness of the radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope. The radiopharmaceuticals are administered in appropriate amounts and the uptake at the angiogenic sites can be quantified either non-invasively by imaging for those isotopes with a coincident imageable gamma emission, or by excision of the angiogenic sites and counting the amount of radioactivity present by standard techniques. The therapeutic effect of the radiopharmaceuticals can be assessed by monitoring the rate of

growth of the angiogenic sites in control rabbits versus those in the rabbits administered the radiopharmaceuticals of the present invention.

This model can also be used to assess the compounds of the present invention comprised of paramagnetic metals as MRI contrast agents. After administration of the appropriate amount of the paramagnetic compounds, the whole animal can be placed in a commercially available magnetic resonance imager to image the angiogenic sites. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of heavy atoms as X-ray contrast agents. After administration of the appropriate amount of the X-ray absorbing compounds, the whole animal can be placed in a commercially available X-ray imager to image the angiogenic sites. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of an echogenic gas containing surfactant microsphere as ultrasound contrast agents. After administration of the appropriate amount of the echogenic compounds, the angiogenic sites in the animal can be imaging using an ultrasound probe held proximate to the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

30

Canine Spontaneous Tumor Model

Adult dogs with spontaneous mammary tumors were sedated with xylazine (20 mg/kg)/atropine (1 ml/kg). Upon sedation the animals were intubated using ketamine (5 mg/kg)/diazepam (0.25 mg/kg) for full anesthesia. Chemical restraint was continued with ketamine (3 mg/kg)/xylazine (6 mg/kg) titrating as necessary. If required the animals were ventilated with room air via an endotracheal tube (12 strokes/min, 25 ml/kg)

during the study. Peripheral veins were catheterized using 20G I.V. catheters, one to serve as an infusion port for compound while the other for exfusion of blood samples. Heart rate and EKG were monitored using a cardiometer (Biotech, Grass Quincy, MA) triggered from a lead II electrocardiogram generated by limb leads. Blood samples are generally taken at ~10 minutes (control), end of infusion, (1 minute), 15 min, 30 min, 60 min, 90 min, and 120 min for whole blood cell number and counting. Radiopharmaceutical dose was 300 $\mu\text{Ci}/\text{kg}$ administered as an i.v. bolus with saline flush. Parameters were monitored continuously on a polygraph recorder (Model 7E Grass) at a paper speed of 10 mm/min or 10 mm/sec.

Imaging of the laterals were for 2 hours with a 256x256 matrix, no zoom, 5 minute dynamic images. A known source is placed in the image field (20-90 μCi) to evaluate region of interest (ROI) uptake. Images were also acquired 24 hours post injection to determine retention of the compound in the tumor. The uptake is determined by taking the fraction of the total counts in an inscribed area for ROI/source and multiplying the known μCi . The result is μCi for the ROI.

This model can also be used to assess the effectiveness of the radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope. The radiopharmaceuticals are administered in appropriate amounts and the uptake in the tumors can be quantified either non-invasively by imaging for those isotopes with a coincident imageable gamma emission, or by excision of the tumors and counting the amount of radioactivity present by standard techniques. The therapeutic effect of the radiopharmaceuticals can be assessed by monitoring the size of the tumors over time.

This model can also be used to assess the compounds of the present invention comprised of paramagnetic metals as MRI contrast agents. After administration of the appropriate amount of the paramagnetic compounds, the whole animal can be placed in a commercially available magnetic resonance imager to image the tumors. The effectiveness of the contrast agents

can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

5 This model can also be used to assess the compounds of the present invention comprised of heavy atoms as X-ray contrast agents. After administration of the appropriate amount of the X-ray absorbing compounds, the whole animal can be placed in a commercially available X-ray imager to image the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals
10 that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of an echogenic gas containing surfactant microsphere as ultrasound contrast agents. After administration of the appropriate amount of the echogenic
15 compounds, the tumors in the animal can be imaging using an ultrasound probe held proximate to the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

20 Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise that as specifically described herein.

WHAT IS CLAIMED IS:


1. A compound, comprising: a targeting moiety and a chelator, wherein the targeting moiety is bound to the chelator, is a peptide or peptidomimetic, and binds to a receptor that is upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.

2. A compound according to Claim 1, wherein the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

3. A compound according to Claim 2, the receptor is the integrin $\alpha_v\beta_3$ and the compound is of the formula:

(Q)_d-L_n-C_h or (Q)_d-L_n-(C_h)_{d'}

wherein, Q is a peptide independently selected from the group:



K is an L-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine, δ -N-2-imidazolinylnornithine, δ -N-benzylcarbamoylnornithine, and β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

K' is a D-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine,

δ -N-2-imidazolinylnornithine,
 δ -N-benzylcarbamoynornithine, and
 β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

5 L is independently selected at each occurrence from the group:
glycine, L-alanine, and D-alanine;

M is L-aspartic acid;

10 M' is D-aspartic acid;

R¹ is an amino acid substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, L-valine, D-valine, alanine, leucine,
15 isoleucine, norleucine, 2-aminobutyric acid,
2-aminohexanoic acid, tyrosine, phenylalanine,
thienylalanine, phenylglycine, cyclohexylalanine,
homophenylalanine, 1-naphthylalanine, lysine, serine,
ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
20 acid, cysteine, penicillamine, and methionine;

R² is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, valine, alanine, leucine, isoleucine,
25 norleucine, 2-aminobutyric acid, 2-aminohexanoic acid,
tyrosine, L-phenylalanine, D-phenylalanine,
thienylalanine, phenylglycine, biphenylglycine,
cyclohexylalanine, homophenylalanine,
L-1-naphthylalanine, D-1-naphthylalanine, lysine, serine,
30 ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
acid, cysteine, penicillamine, methionine, and
2-aminothiazole-4-acetic acid;

R³ is an amino acid, substituted with 0-1 bonds to L_n,
35 independently selected at each occurrence from the group:
glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
acid, D-tyrosine, D-phenylalanine, D-thienylalanine,

D-phenylglycine, D-cyclohexylalanine,
D-homophenylalanine, D-1-naphthylalanine, D-lysine,
D-serine, D-ornithine, D-1,2-diaminobutyric acid,
D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
5 and D-methionine;


R⁴ is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
10 D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
D-phenylglycine, D-cyclohexylalanine,
D-homophenylalanine, D-1-naphthylalanine, D-lysine,
D-serine, D-ornithine, D-1,2-diaminobutyric acid,
15 D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
D-methionine, and 2-aminothiazole-4-acetic acid;

R⁵ is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
20 glycine, L-valine, L-alanine, L-leucine, L-isoleucine,
L-norleucine, L-2-aminobutyric acid, L-2-aminohexanoic
acid, L-tyrosine, L-phenylalanine, L-thienylalanine,
L-phenylglycine, L-cyclohexylalanine,
L-homophenylalanine, L-1-naphthylalanine, L-lysine,
25 L-serine, L-ornithine, L-1,2-diaminobutyric acid,
L-1,2-diaminopropionic acid, L-cysteine, L-penicillamine,
L-methionine, and 2-aminothiazole-4-acetic acid;

provided that one of R¹, R², R³, R⁴, and R⁵ in each Q is
30 substituted with a bond to L_n, further provided that when
R² is 2-aminothiazole-4-acetic acid, K is
N-methylarginine, further provided that when R⁴ is
2-aminothiazole-4-acetic acid, K and K' are
N-methylarginine, and still further provided that when R⁵
35 is 2-aminothiazole-4-acetic acid, K' is N-methylarginine;

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

L_n is a linking group having the formula:



5 provided that $g+h+g'+k+h'+g''+h''+g'''$ is other than 0;

W is independently selected at each occurrence from the group:

O, S, NH, NHC(=O), C(=O)NH, C(=O), C(=O)O, OC(=O),
 NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'},
 10 (OCH₂CH₂CH₂)_{s''}, (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R¹⁰,
 15 C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁰, and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 substituted with 0-3 R¹⁰;

20 R⁶, R^{6a}, R⁷, R^{7a}, R⁸, R^{8a}, R⁹ and R^{9a} are independently selected
 at each occurrence from the group: H, =O, COOH, SO₃H,
 PO₃H, C₁₋₅ alkyl substituted with 0-3 R¹⁰, aryl
 substituted with 0-3 R¹⁰, benzyl substituted with 0-3
 R¹⁰, and C₁₋₅ alkoxy substituted with 0-3 R¹⁰,
 25 NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and a
 bond to C_H;

R¹⁰ is independently selected at each occurrence from the
 group: a bond to C_H, COOR¹¹, OH, NHR¹¹, SO₃H, PO₃H, aryl
 30 substituted with 0-3 R¹¹, C₁₋₅ alkyl substituted with 0-1
 R¹², C₁₋₅ alkoxy substituted with 0-1 R¹², and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 substituted with 0-3 R¹¹;

35

R¹¹ is independently selected at each occurrence from the
 group: H, aryl substituted with 0-1 R¹², a 5-10 membered
 heterocyclic ring system containing 1-4 heteroatoms

independently selected from N, S, and O and substituted
 with 0-1 R¹², C₃₋₁₀ cycloalkyl substituted with 0-1 R¹²,
 polyalkylene glycol substituted with 0-1 R¹²,
 carbohydrate substituted with 0-1 R¹², cyclodextrin
 5 substituted with 0-1 R¹², amino acid substituted with 0-1
 R¹², polycarboxyalkyl substituted with 0-1 R¹²,
 polyazaalkyl substituted with 0-1 R¹², peptide
 substituted with 0-1 R¹², wherein the peptide is
 comprised of 2-10 amino acids, and a bond to C_h;

10

R¹² is a bond to C_h;

k is selected from 0, 1, and 2;

h is selected from 0, 1, and 2;

15

h' is selected from 0, 1, 2, 3, 4, and 5;

h" is selected from 0, 1, 2, 3, 4, and 5;

g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

g" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

20

g"' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;



s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;


t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

25

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

C_h is a metal bonding unit having a formula selected from the
 group:





5 A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , and A^8 are independently selected at each occurrence from the group N, NR^{13} , $NR^{13}R^{14}$, S, SH, S(Pg), O, OH, PR^{13} , $PR^{13}R^{14}$, $P(O)R^{15}R^{16}$, and a bond to L_n ;

10 E is a bond, CH, or a spacer group independently selected at each occurrence from the group: C_1 - C_{10} alkyl substituted with 0-3 R^{17} , aryl substituted with 0-3 R^{17} , C_3 - C_{10} cycloalkyl substituted with 0-3 R^{17} , heterocyclo- C_1 - C_{10} alkyl substituted with 0-3 R^{17} , wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C_6 - C_{10} aryl- C_1 - C_{10} alkyl substituted with 0-3 R^{17} , C_1 - C_{10} alkyl- C_6 - C_{10} aryl- substituted with 0-3 R^{17} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{17} ;

20 R^{13} , and R^{14} are each independently selected from the group: a bond to L_n , hydrogen, C_1 - C_{10} alkyl substituted with 0-3 R^{17} , aryl substituted with 0-3 R^{17} , C_1 - C_{10} cycloalkyl substituted with 0-3 R^{17} , heterocyclo- C_1 - C_{10} alkyl substituted with 0-3 R^{17} , wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C_6 - C_{10} aryl- C_1 - C_{10} alkyl substituted with 0-3 R^{17} , C_1 - C_{10} alkyl- C_6 - C_{10} aryl- substituted with 0-3 R^{17} , a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and

30

substituted with 0-3 R^{17} , and an electron, provided that when one of R^{13} or R^{14} is an electron, then the other is also an electron;

5 alternatively, R^{13} and R^{14} combine to form $=C(R^{20})(R^{21})$;

R^{15} and R^{16} are each independently selected from the group: a bond to L_n , -OH, C₁-C₁₀ alkyl substituted with 0-3 R^{17} , C₁-C₁₀ alkyl substituted with 0-3 R^{17} , aryl substituted
10 with 0-3 R^{17} , C₃-₁₀ cycloalkyl substituted with 0-3 R^{17} , heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R^{17} , wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl
15 substituted with 0-3 R^{17} , C₁-₁₀ alkyl-C₆-₁₀ aryl-substituted with 0-3 R^{17} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{17} ;

20 R^{17} is independently selected at each occurrence from the group: a bond to L_n , =O, F, Cl, Br, I, -CF₃, -CN, -CO₂ R^{18} , -C(=O) R^{18} , -C(=O)N(R^{18})₂, -CHO, -CH₂OR¹⁸, -OC(=O) R^{18} , -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R^{18})₂,
25 -NR¹⁹C(=O) R^{18} , -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R^{18})₂, -NR¹⁹SO₂N(R^{18})₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a}, -SR¹⁸, -S(=O) R^{18a} , -SO₂N(R^{18})₂, -N(R^{18})₂, -NHC(=S)NHR¹⁸, =NOR¹⁸, NO₂, -C(=O)NHOR¹⁸, -C(=O)NHNR¹⁸R^{18a}, -OCH₂CO₂H, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆
30 cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆ alkoxyalkyl, aryl substituted with 0-2 R^{18} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O;


35 R^{18} , R^{18a} , and R^{19} are independently selected at each occurrence from the group: a bond to L_n , H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy, halide, nitro, cyano, and trifluoromethyl;

Pg is a thiol protecting group;

R²⁰ and R²¹ are independently selected from the group: H,
 5 C₁-C₁₀ alkyl, -CN, -CO₂R²⁵, -C(=O)R²⁵, -C(=O)N(R²⁵)₂,
 C₂-C₁₀ 1-alkene substituted with 0-3 R²³, C₂-C₁₀ 1-alkyne
 substituted with 0-3 R²³, aryl substituted with 0-3 R²³,
 unsaturated 5-10 membered heterocyclic ring system
 containing 1-4 heteroatoms independently selected from N,
 10 S, and O and substituted with 0-3 R²³, and unsaturated
 C₃-C₁₀ carbocycle substituted with 0-3 R²³;

alternatively, R²⁰ and R²¹, taken together with the divalent
 carbon radical to which they are attached form:

15



R²² and R²³ are independently selected from the group: H, R²⁴,
 C₁-C₁₀ alkyl substituted with 0-3 R²⁴, C₂-C₁₀ alkenyl
 substituted with 0-3 R²⁴, C₂-C₁₀ alkynyl substituted with
 20 0-3 R²⁴, aryl substituted with 0-3 R²⁴, a 5-10 membered
 heterocyclic ring system containing 1-4 heteroatoms
 independently selected from N, S, and O and substituted
 with 0-3 R²⁴, and C₃-C₁₀ carbocycle substituted with 0-3
 R²⁴;

25

alternatively, R²², R²³ taken together form a fused aromatic
 or a 5-10 membered heterocyclic ring system containing
 1-4 heteroatoms independently selected from N, S, and O;

30 a and b indicate the positions of optional double bonds and n
 is 0 or 1;

R^{24} is independently selected at each occurrence from the group: =O, F, Cl, Br, I, $-CF_3$, $-CN$, $-CO_2R^{25}$, $-C(=O)R^{25}$, $-C(=O)N(R^{25})_2$, $-N(R^{25})_3^+$, $-CH_2OR^{25}$, $-OC(=O)R^{25}$, $-OC(=O)OR^{25a}$, $-OR^{25}$, $-OC(=O)N(R^{25})_2$, $-NR^{26}C(=O)R^{25}$, $-NR^{26}C(=O)OR^{25a}$, $-NR^{26}C(=O)N(R^{25})_2$, $-NR^{26}SO_2N(R^{25})_2$, $-NR^{26}SO_2R^{25a}$, $-SO_3H$, $-SO_2R^{25a}$, $-SR^{25}$, $-S(=O)R^{25a}$, $-SO_2N(R^{25})_2$, $-N(R^{25})_2$, $=NOR^{25}$, $-C(=O)NHOR^{25}$, $-OCH_2CO_2H$, and 2-(1-morpholino)ethoxy; and,

R^{25} , R^{25a} , and R^{26} are each independently selected at each occurrence from the group: hydrogen and C_1 - C_6 alkyl;

and a pharmaceutically acceptable salt thereof.

15

4. A compound according to Claim 3, the present invention provides a compound, wherein:

L is glycine;

20

R^1 is an amino acid, optionally substituted with a bond to L_n , independently selected at each occurrence from the group: L-valine, D-valine, alanine, leucine, isoleucine, norleucine, 2-aminobutyric acid, tyrosine, phenylalanine, phenylglycine, cyclohexylalanine, homophenylalanine, lysine, ornithine, 1,2-diaminobutyric acid, and 1,2-diaminopropionic acid;

25

R^2 is an amino acid, optionally substituted with a bond to L_n , independently selected at each occurrence from the group: valine, alanine, leucine, isoleucine, norleucine, 2-aminobutyric acid, tyrosine, L-phenylalanine, D-phenylalanine, thienylalanine, phenylglycine, biphenylglycine, cyclohexylalanine, homophenylalanine, L-1-naphthylalanine, D-1-naphthylalanine, lysine, ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic acid, and 2-aminothiazole-4-acetic acid;

35

- R^3 is an amino acid, optionally substituted with a bond to L_n , independently selected at each occurrence from the group:
 D-valine, D-alanine, D-leucine, D-isoleucine,
 D-norleucine, D-2-aminobutyric acid, D-tyrosine,
 5 D-phenylalanine, D-phenylglycine, D-cyclohexylalanine,
 D-homophenylalanine, D-lysine, D-serine, D-ornithine,
 D-1,2-diaminobutyric acid, and D-1,2-diaminopropionic acid;
- 10 R^4 is an amino acid, optionally substituted with a bond to L_n , independently selected at each occurrence from the group:
 D-valine, D-alanine, D-leucine, D-isoleucine,
 D-norleucine, D-2-aminobutyric acid, D-tyrosine,
 D-phenylalanine, D-thienylalanine, D-phenylglycine,
 15 D-cyclohexylalanine, D-homophenylalanine,
 D-1-naphthylalanine, D-lysine, D-ornithine,
 D-1,2-diaminobutyric acid, D-1,2-diaminopropionic acid,
 and 2-aminothiazole-4-acetic acid;
- 20 R^5 is an amino acid, optionally substituted with a bond to L_n , independently selected at each occurrence from the group:
 L-valine, L-alanine, L-leucine, L-isoleucine,
 L-norleucine, L-2-aminobutyric acid, L-tyrosine,
 L-phenylalanine, L-thienylalanine, L-phenylglycine,
 25 L-cyclohexylalanine, L-homophenylalanine,
 L-1-naphthylalanine, L-lysine, L-ornithine,
 L-1,2-diaminobutyric acid, L-1,2-diaminopropionic acid,
 and 2-aminothiazole-4-acetic acid;
- 30 d is selected from 1, 2, and 3;
- W is independently selected at each occurrence from the group:
 O, NH, $NHC(=O)$, $C(=O)NH$, $C(=O)$, $C(=O)O$, $OC(=O)$,
 $NHC(=S)NH$, $NHC(=O)NH$, SO_2 , $(OCH_2CH_2)_s$, $(CH_2CH_2O)_s$,
 35 $(OCH_2CH_2CH_2)_s$, and $(CH_2CH_2CH_2O)_t$.
- Z is selected from the group: aryl substituted with 0-1 R^{10} ,
 C_{3-10} cycloalkyl substituted with 0-1 R^{10} , and a 5-10

membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹⁰;

- 5 R⁶, R^{6a}, R⁷, R^{7a}, R⁸, R^{8a}, R⁹, and R^{9a} are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, C₁-C₅ alkyl substituted with 0-1 R¹⁰, aryl substituted with 0-1 R¹⁰, benzyl substituted with 0-1 R¹⁰, and C₁-C₅ alkoxy substituted with 0-1 R¹⁰,
10 NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and a bond to C_H;

- R¹⁰ is independently selected at each occurrence from the group: COOR¹¹, OH, NHR¹¹, SO₃H, aryl substituted with
15 0-1 R¹¹, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹¹, C₁-C₅ alkyl substituted with 0-1 R¹², C₁-C₅ alkoxy substituted with 0-1 R¹², and a bond to C_H;

- 20 R¹¹ is independently selected at each occurrence from the group: H, aryl substituted with 0-1 R¹², a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted
25 with 0-1 R¹², polyalkylene glycol substituted with 0-1 R¹², carbohydrate substituted with 0-1 R¹², cyclodextrin substituted with 0-1 R¹², amino acid substituted with 0-1 R¹², and a bond to C_H;

- 30 k is 0 or 1;
h is 0 or 1;
h' is 0 or 1;
s is selected from 0, 1, 2, 3, 4, and 5;
s' is selected from 0, 1, 2, 3, 4, and 5;
35 s'' is selected from 0, 1, 2, 3, 4, and 5;
t is selected from 0, 1, 2, 3, 4, and 5;

A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently selected at each occurrence from the group: NR¹³, NR¹³R¹⁴, S, SH, S(Pg), OH, and a bond to L_n;

5 E is a bond, CH, or a spacer group independently selected at each occurrence from the group: C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₃-₁₀ cycloalkyl substituted with 0-3 R¹⁷, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms
10 independently selected from N, S, and O and substituted with 0-3 R¹⁷;

R¹³, and R¹⁴ are each independently selected from the group:
a bond to L_n, hydrogen, C₁-C₁₀ alkyl substituted with 0-3
15 R¹⁷, aryl substituted with 0-3 R¹⁷, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷, and an electron, provided that when one of R¹³ or R¹⁴ is an electron, then the other is also an
20 electron;

alternatively, R¹³ and R¹⁴ combine to form =C(R²⁰)(R²¹);


R¹⁷ is independently selected at each occurrence from the
25 group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹⁸, -C(=O)R¹⁸, -C(=O)N(R¹⁸)₂, -CH₂OR¹⁸, -OC(=O)R¹⁸, -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R¹⁸)₂, -NR¹⁹C(=O)R¹⁸, -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R¹⁸)₂, -NR¹⁹SO₂N(R¹⁸)₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a}, -S(=O)R^{18a}, -SO₂N(R¹⁸)₂,
30 -N(R¹⁸)₂, -NHC(=S)NHR¹⁸, =NOR¹⁸, -C(=O)NHN(R¹⁸)R^{18a}, -OCH₂CO₂H, and 2-(1-morpholino)ethoxy;

R¹⁸, R^{18a}, and R¹⁹ are independently selected at each
occurrence from the group: a bond to L_n, H, and C₁-C₆
35 alkyl;

R²⁰ and R²¹ are independently selected from the group: H, C₁-C₅ alkyl, -CO₂R²⁵, C₂-C₅ 1-alkene substituted with 0-3

5 R^{23} , C_2-C_5 1-alkyne substituted with 0-3 R^{23} , aryl substituted with 0-3 R^{23} , and unsaturated 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{23} ;

alternatively, R^{20} and R^{21} , taken together with the divalent carbon radical to which they are attached form:



10 R^{22} and R^{23} are independently selected from the group: H, and R^{24} ;

15 alternatively, R^{22} , R^{23} taken together form a fused aromatic or a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O;


20 R^{24} is independently selected at each occurrence from the group: $-CO_2R^{25}$, $-C(=O)N(R^{25})_2$, $-CH_2OR^{25}$, $-OC(=O)R^{25}$, $-OR^{25}$, $-SO_3H$, $-N(R^{25})_2$, and $-OCH_2CO_2H$; and,

R^{25} is independently selected at each occurrence from the group: H and C_1-C_3 alkyl.

25

5. A compound according to Claim 4, the present invention provides a compound, wherein:

30 Q is a peptide selected from the group:



R^1 is L-valine, D-valine, D-lysine optionally substituted on the ϵ amino group with a bond to L_n or L-lysine optionally substituted on the ϵ amino group with a bond to L_n ;

R^2 is L-phenylalanine, D-phenylalanine, D-1-naphthylalanine, 2-aminothiazole-4-acetic acid, L-lysine optionally substituted on the ϵ amino group with a bond to L_n or tyrosine, the tyrosine optionally substituted on the hydroxy group with a bond to L_n ;

R^3 is D-valine, D-phenylalanine, or L-lysine optionally substituted on the ϵ amino group with a bond to L_n ;

R^4 is D-phenylalanine, D-tyrosine substituted on the hydroxy group with a bond to L_n , or L-lysine optionally substituted on the ϵ amino group with a bond to L_n ;

provided that one of R^1 and R^2 in each Q is substituted with a bond to L_n , and further provided that when R^2 is 2-aminothiazole-4-acetic acid, K is N-methylarginine;

d is 1 or 2;

W is independently selected at each occurrence from the group: $\text{NHC}(=\text{O})$, $\text{C}(=\text{O})\text{NH}$, $\text{C}(=\text{O})$, $(\text{CH}_2\text{CH}_2\text{O})_s$, and $(\text{CH}_2\text{CH}_2\text{CH}_2\text{O})_t$;

R^6 , R^{6a} , R^7 , R^{7a} , R^8 , R^{8a} , R^9 , and R^{9a} are independently selected at each occurrence from the group: H, $\text{NHC}(=\text{O})\text{R}^{11}$, and a bond to C_h ;

k is 0;

t is 1 or 2;



20

30


35

R^{18} is a bond to L_n ;

R^{24} is selected from the group: $-\text{CO}_2R^{25}$, $-\text{OR}^{25}$, $-\text{SO}_3\text{H}$, and $-\text{N}(\text{R}^{25})_2$;

5

R^{25} is independently selected at each occurrence from the group: hydrogen and methyl;



alternatively, C_h is

10

A^1 , A^2 , A^3 , and A^4 are each N;

A^5 , A^6 , and A^8 are each OH;

15

A^7 is a bond to L_n ;

E is a C_2 alkyl substituted with 0-1 R^{17} ; and,

R^{17} is =O.

20

6. A compound according to Claim 3, the present invention provides a compound selected from the group:

25

(a) cyclo{Arg-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};

30

(b) cyclo{Arg-Gly-Asp-D-Tyr((N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-18-

amino-14-aza-4,7,10-oxy-15-oxo-octadecoyl)-3-aminopropyl)-Val}};

- 5 (c) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Glu(cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp})-cyclo{D-Tyr(3-aminopropyl)-Val-Arg-Gly-Asp});
- 10 (d) cyclo{Arg-Gly-Asp-D-Tyr-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])});
- (e) cyclo{Arg-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])});
- 15 (f) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};
- (g) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Phe-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};
- 20 (h) cyclo{Arg-Gly-Asp-D-Nal-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])});
- 25 (i) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-Arg-Gly-Asp-D-Nal})-cyclo{Lys-Arg-Gly-Asp-D-Nal};
- 30 (j) cyclo{Arg-Gly-Asp-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid])-D-Val};
- (k) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-Glu(cyclo{Lys-D-Val-Arg-Gly-Asp})-cyclo{Lys-D-Val-Arg-Gly-Asp};
- 35

- (l) {cyclo(Arg-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly)};
- 5 (m) cyclo{D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Phe-D-Asp-Gly-Arg)};
- (n) [2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Glu(cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg))-cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg)};
- 10 (o) cyclo{D-Phe-D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Asp-Gly-Arg)};
- 15 (p) cyclo{N-Me-Arg-Gly-Asp-ATA-D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])});
- 20 (q) cyclo{Cit-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])});
- (r) 2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)-1-cyclododecyl)acetyl-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe))-cyclo{Lys-Arg-Gly-Asp-D-Phe)};
- 25 (s) cyclo{Arg-Gly-Asp-D-Phe-Lys(DTPA)};
- (t) cyclo{Arg-Gly-Asp-D-Phe-Lys}₂(DTPA);
- 30 (u) Cyclo{Arg-Gly-Asp-D-Tyr(N-DTPA-3-aminopropyl)-Val)};
- (v) cyclo{Orn(d-N-2-Imidazoliny)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-Val)};
- 35

- (w) cyclo{Lys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 5 (x) cyclo{Cys(2-aminoethyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- (y) cyclo{HomoLys-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 10 (z) cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 15 (aa) cyclo{Dap(b-(2-benzimidazolylacetyl))-Gly-Asp-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-Val};
- 20 (bb) cyclo{Orn(d-N-2-Imidazolinyl)-Gly-Asp-D-Phe-Lys(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]]);
- 25 (cc) cyclo{Orn(d-N-Benzylcarbamoyl)-Gly-Asp-D-Phe-Lys(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]]);
- (dd) cyclo{Lys-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly};
- 30 (ee) cyclo{Orn(d-N-Benzylcarbamoyl)-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyl]hydrazono)methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly}; and,
- 35

(ff) cyclo{Orn(d-N-2-ImidazolinyI)-D-Val-D-Tyr(N-[2-[[[5-[carbonyl]-2-pyridinyI]hydrazono]methyl]-benzenesulfonic acid]-3-aminopropyl)-D-Asp-Gly);

5 or a pharmaceutically acceptable salt form thereof.

7. A kit comprising a compound of Claim 3, or a pharmaceutically acceptable salt form thereof and a
10 pharmaceutically acceptable carrier.

8. A kit according to Claim 7, wherein the kit further comprises one or more ancillary ligands and a reducing agent.
15

9. A kit according to Claim 8, wherein the ancillary ligands are tricine and TPPTS.

20 10. A kit according to Claim 9, wherein the reducing agent is tin(II).

25 11. A diagnostic or therapeutic metallopharmaceutical composition, comprising: a metal, a chelator capable of chelating the metal and a targeting moiety, wherein the targeting moiety is bound to the chelator, is a peptide or peptidomimetic and binds to a receptor that is upregulated
30 during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.

12. A composition according to Claim 11, wherein the
35 metallopharmaceutical is a diagnostic radiopharmaceutical, the metal is a radioisotope selected from the group: ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{64}Cu , ^{67}Ga , and ^{68}Ga , the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from

the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

5

13. A composition according to Claim 12, wherein the targeting moiety is a cyclic pentapeptide and the receptor is $\alpha_v\beta_3$.

10

14. A composition according to Claim 13, wherein the radioisotope is ^{99m}Tc or ^{95}Tc , the radiopharmaceutical further comprises a first ancillary ligand and a second ancillary ligand capable of stabilizing the radiopharmaceutical.

15

15. A composition according to Claim 14, wherein the radioisotope is ^{99m}Tc .

20

16. A composition according to Claim 15, wherein the radiopharmaceutical is selected from the group:

25 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{Arg-Gly-Asp-D-Tyr}(\text{N}-[[5-[\text{carbonyl}]-2\text{-pyridinyl}]\text{diazenido}]-3\text{-aminopropyl})-\text{Val}));$

$^{99m}\text{Tc}(\text{tricine})(\text{TPPMS})(\text{cyclo}(\text{Arg-D-Val-D-Tyr}(\text{N}-[[5-[\text{carbonyl}]-2\text{-pyridinyl}]\text{diazenido}]-3\text{-aminopropyl})-\text{D-Asp-Gly}));$

30

$^{99m}\text{Tc}(\text{tricine})(\text{TPPDS})(\text{cyclo}(\text{Arg-D-Val-D-Tyr}(\text{N}-[[5-[\text{carbonyl}]-2\text{-pyridinyl}]\text{diazenido}]-3\text{-aminopropyl})-\text{D-Asp-Gly}));$

35 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{Arg-D-Val-D-Tyr}(\text{N}-[[5-[\text{carbonyl}]-2\text{-pyridinyl}]\text{diazenido}]-3\text{-aminopropyl})-\text{D-Asp-Gly}));$

$^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{Arg-Gly-Asp-D-Phe-Lys}(\text{N}-[[5-[\text{carbonyl}]-2\text{-pyridinyl}]\text{diazenido}]]));$

- $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{Arg-Gly-Asp-D-Tyr-Lys}(\text{N}-[[5-\text{[carbonyl]}-2\text{-pyridinyl]diazenido}]]));$
- 5 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})([2-[[[5-\text{[carbonyl]}-2\text{-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-Phe-Glu}(\text{cyclo}(\text{Lys-Arg-Gly-Asp-D-Phe}))- \text{cyclo}(\text{Lys-Arg-Gly-Asp-D-Phe}));$
- 10 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{Arg-Gly-Asp-D-Nal-Lys}([2-[[[5-\text{[carbonyl]}-2\text{-pyridinyl]hydrazono]methyl]-benzenesulfonic acid}]]));$
- 15 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})([2-[[[5-\text{[carbonyl]}-2\text{-pyridinyl]-hydrazono]methyl]-benzenesulfonic acid]-Glu}(\text{cyclo}(\text{Lys-Arg-Gly-Asp-D-Nal}))- \text{cyclo}(\text{Lys-Arg-Gly-Asp-D-Nal}));$
- 20 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{Arg-Gly-Asp-D-Tyr}(\text{N}-[[5-\text{[carbonyl]}-2\text{-pyridinyl]diazenido]-18\text{-amino-14-aza-4,7,10-oxo-15-oxo-octadecoyl]-3-aminopropyl}-\text{Val}));$
- 25 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{N}-[[5-\text{[carbonyl]}-2\text{-pyridinyl]diazenido]-Glu}(\text{O-cyclo}(\text{Lys-Arg-Gly-Asp-D-Phe}))- \text{O-cyclo}(\text{Lys-Arg-Gly-Asp-D-Phe}));$
- 30 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{N}-[[5-\text{[carbonyl]}-2\text{-pyridinyl]diazenido]-Glu}(\text{O-cyclo}(\text{D-Tyr}(3\text{-aminopropyl})-\text{Val-Arg-Gly-Asp}))- \text{O-cyclo}(\text{D-Tyr}(3\text{-aminopropyl})-\text{Val-Arg-Gly-Asp}));$
- 35 $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{Arg-Gly-Asp-Lys}(\text{N}-[[5-\text{[carbonyl]}-2\text{-pyridinyl]diazenido}])- \text{D-Val}));$
- $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{cyclo}(\text{D-Lys}([2-[[[5-\text{[carbonyl]}-2\text{-pyridinyl]hydrazono]methyl]-benzenesulfonic acid}])- \text{D-Phe-D-Asp-Gly-Arg}));$
- $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})([2-[[[5-\text{[carbonyl]}-2\text{-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]-$

Glu(cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg})-cyclo{D-Lys-D-Phe-D-Asp-Gly-Arg});

5 ^{99m}Tc (tricine)(TPPTS)(cyclo{D-Phe-D-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid])-D-Asp-Gly-Arg});

10 ^{99m}Tc (tricine)(TPPTS)(cyclo{N-Me-Arg-Gly-Asp-ATA-D-Lys(N-[5-[carbonyl]-2-pyridinyl]diazenido)}));

^{99m}Tc (tricine)(TPPTS)(cyclo{Cit-Gly-Asp-D-Phe-Lys([2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid]]}); and,

15 ^{99m}Tc (tricine)(1,2,4-triazole)(cyclo{Arg-Gly-Asp-D-Tyr(N-[5-[carbonyl]-2-pyridinyl]diazenido]-3-aminopropyl)-Val)).

20 17. A composition according to Claim 13, wherein the radioisotope is ^{111}In .

25 18. A composition according to Claim 17, wherein the radiopharmaceutical is selected from the group:

(DOTA- ^{111}In)-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};

cyclo{Arg-Gly-Asp-D-Phe-Lys(DTPA- ^{111}In)}; and,

30 cyclo{Arg-Gly-Asp-D-Phe-Lys}₂(DTPA- ^{111}In).

35 19. A composition according to Claim 11, wherein the metallopharmaceutical is a therapeutic radiopharmaceutical, the metal is a radioisotope selected from the group: ^{186}Re , ^{188}Re , ^{153}Sm , ^{166}Ho , ^{177}Lu , ^{149}Pm , ^{90}Y , ^{212}Bi , ^{103}Pd , ^{109}Pd , ^{159}Gd , ^{140}La , ^{198}Au , ^{199}Au , ^{169}Yb , ^{175}Yb , ^{165}Dy , ^{166}Dy , ^{67}Cu ,

10⁵Rh, ¹¹¹Ag, and ¹⁹²Ir, the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and
5 $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

20. A composition according to Claim 19, wherein the
10 targeting moiety is a cyclic pentapeptide and the receptor is $\alpha_v\beta_3$.

21. A composition according to Claim 20, wherein the
15 radioisotope is ¹⁵³Sm.

22. A composition according to Claim 21, wherein the radiopharmaceutical is selected from the group:
20
cyclo(Arg-Gly-Asp-D-Phe-Lys(DTPA-¹⁵³Sm));
cyclo(Arg-Gly-Asp-D-Phe-Lys)₂(DTPA-¹⁵³Sm); and,
25 cyclo(Arg-Gly-Asp-D-Tyr(N-DTPA(¹⁵³Sm)-3-aminopropyl)-Val).

23. A composition according to Claim 20, wherein the
30 radioisotope is ¹⁷⁷Lu.

24. A composition according to Claim 23, wherein the radiopharmaceutical is selected from the group:
35 cyclo(Arg-Gly-Asp-D-Phe-Lys(DTPA-¹⁷⁷Lu));
(DOTA-¹⁷⁷Lu)-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-Gly-Asp-D-Phe};

cyclo(Arg-Gly-Asp-D-Phe-Lys)₂(DTPA-¹⁷⁷Lu); and,

cyclo(Arg-Gly-Asp-D-Tyr(N-DTPA(¹⁷⁷Lu)-3-aminopropyl)-Val).

5

25. A composition according to Claim 20, wherein the radioisotope is ⁹⁰Y.

10

26. A composition according to Claim 25, wherein the radiopharmaceutical is:

(DOTA-⁹⁰Y)-Glu(cyclo{Lys-Arg-Gly-Asp-D-Phe})-cyclo{Lys-Arg-
15 Gly-Asp-D-Phe};

27. A composition according to Claim 11, wherein the metallopharmaceutical is a MRI contrast agent, the metal is a
20 paramagnetic metal ion selected from the group: Gd(III), Dy(III), Fe(III), and Mn(II), the targeting moiety is a peptide or a mimetic thereof and the receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$,
25 $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking group is present between the targeting moiety and chelator.

28. A composition according to Claim 27, wherein the
30 targeting moiety is a cyclic pentapeptide and the receptor is $\alpha_v\beta_3$.

29. A composition according to Claim 28, wherein the
35 metal ion is Gd(III).

30. A composition according to Claim 29, wherein the contrast agent is:

cyclo(Arg-Gly-Asp-D-Tyr(N-DTPA(Gd(III))-3-aminopropyl)-Val).

5

31. A composition according to Claim 11, wherein the metallopharmaceutical is a X-ray contrast agent, the metal is selected from the group: Re, Sm, Ho, Lu, Pm, Y, Bi, Pd, Gd, La, Au, Au, Yb, Dy, Cu, Rh, Ag, and Ir, the targeting moiety is a cyclic pentapeptide, the receptor is $\alpha_v\beta_3$, and the linking group is present between the targeting moiety and chelator.

15

32. A composition of Claim 11 which is for use in treating rheumatoid arthritis.

20 33. A composition of Claim 11 which is for use in treating cancer.

25 34. A composition of Claim 11 which is for use in imaging the formation of new blood vessels.

30 35. A composition of Claim 12 which is for use in imaging cancer with planar or SPECT gamma scintigraphy, or positron emission tomography.

35 36. A composition of Claim 27 which is for use in imaging cancer with magnetic resonance imaging.


37. A composition of Claim 31 which is for use in imaging cancer with X-ray computed tomography.

38. A compound, comprising: a targeting moiety and a
 5 surfactant, wherein the targeting moiety is bound to the
 receptor that is upregulated during angiogenesis and the
 compound has 0-1 linking groups between the targeting moiety
 and surfactant.


10

39. A compound according to Claim 38, wherein the
 targeting moiety is a peptide or a mimetic thereof and the
 receptor is selected from the group: EGFR, FGFR, PDGFR, Flk-
 1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin,
 15 Axl, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_5\beta_1$, $\alpha_4\beta_1$, $\alpha_1\beta_1$, and $\alpha_2\beta_2$ and the linking
 group is present between the targeting moiety and surfactant.

40. A compound according to Claim 39, wherein the
 20 receptor is the integrin $\alpha_v\beta_3$ and the compound is of the
 formula:



wherein, Q is a cyclic pentapeptide independently selected
 25 from the group:



K is an L-amino acid independently selected at each occurrence
 30 from the group: arginine, citrulline, N-methylarginine,
 lysine, homolysine, 2-aminoethylcysteine,
 δ -N-2-imidazolinylnornithine,
 δ -N-benzylcarbamoylnornithine, and
 β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

35

K' is a D-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine, δ -N-2-imidazolinylnornithine,
5 δ -N-benzylcarbamoylnornithine, and β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

L is independently selected at each occurrence from the group: glycine, L-alanine, and D-alanine;

10

M is L-aspartic acid;

M' is D-aspartic acid;

15 R¹ is an amino acid substituted with 0-1 bonds to L_n, independently selected at each occurrence from the group: glycine, L-valine, D-valine, alanine, leucine, isoleucine, norleucine, 2-aminobutyric acid, 2-aminohexanoic acid, tyrosine, phenylalanine,
20 thienylalanine, phenylglycine, cyclohexylalanine, homophenylalanine, 1-naphthylalanine, lysine, serine, ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic acid, cysteine, penicillamine, and methionine;

25 R² is an amino acid, substituted with 0-1 bonds to L_n, independently selected at each occurrence from the group: glycine, valine, alanine, leucine, isoleucine, norleucine, 2-aminobutyric acid, 2-aminohexanoic acid, tyrosine, L-phenylalanine, D-phenylalanine,
30 thienylalanine, phenylglycine, biphenylglycine, cyclohexylalanine, homophenylalanine, L-1-naphthylalanine, D-1-naphthylalanine, lysine, serine, ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic acid, cysteine, penicillamine, methionine, and
35 2-aminothiazole-4-acetic acid;

R³ is an amino acid, substituted with 0-1 bonds to L_n, independently selected at each occurrence from the group:

- glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
 D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
 acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
 D-phenylglycine, D-cyclohexylalanine,
 5 D-homophenylalanine, D-1-naphthylalanine, D-lysine,
 D-serine, D-ornithine, D-1,2-diaminobutyric acid,
 D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
 and D-methionine;
- 10 R^4 is an amino acid, substituted with 0-1 bonds to L_n ,
 independently selected at each occurrence from the group:
 glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
 D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
 acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
 15 D-phenylglycine, D-cyclohexylalanine,
 D-homophenylalanine, D-1-naphthylalanine, D-lysine,
 D-serine, D-ornithine, D-1,2-diaminobutyric acid,
 D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
 D-methionine, and 2-aminothiazole-4-acetic acid;
 20
- R^5 is an amino acid, substituted with 0-1 bonds to L_n ,
 independently selected at each occurrence from the group:
 glycine, L-valine, L-alanine, L-leucine, L-isoleucine,
 L-norleucine, L-2-aminobutyric acid, L-2-aminohexanoic
 acid, L-tyrosine, L-phenylalanine, L-thienylalanine,
 25 L-phenylglycine, L-cyclohexylalanine,
 L-homophenylalanine, L-1-naphthylalanine, L-lysine,
 L-serine, L-ornithine, L-1,2-diaminobutyric acid,
 L-1,2-diaminopropionic acid, L-cysteine, L-penicillamine,
 30 L-methionine, and 2-aminothiazole-4-acetic acid;

provided that one of R^1 , R^2 , R^3 , R^4 , and R^5 in each Q is
 substituted with a bond to L_n , further provided that when
 R^2 is 2-aminothiazole-4-acetic acid, K is
 35 N-methylarginine, further provided that when R^4 is
 2-aminothiazole-4-acetic acid, K and K' are
 N-methylarginine, and still further provided that when R^5
 is 2-aminothiazole-4-acetic acid, K' is N-methylarginine;

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

S_f is a surfactant which is a lipid or a compound of the

5 formula: A^9-E-A^{10} ;

A⁹ is selected from the group: OH and OR²⁷;

A¹⁰ is OR²⁷;

10

R²⁷ is C(=O)C₁₋₂₀ alkyl;

E¹ is C₁₋₁₀ alkylene substituted with 1-3 R²⁸;


15 R²⁸ is independently selected at each occurrence from the group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹, -C(=O)N(R²⁹)₂, -CH₂OR²⁹, -OR²⁹, -N(R²⁹)₂, C₁-C₅ alkyl, and C₂-C₄ alkenyl;

20 R²⁹ is independently selected at each occurrence from the group: R³⁰, H, C₁-C₆ alkyl, phenyl, benzyl, and trifluoromethyl;

R³⁰ is a bond to L_n;

25

L_n is a linking group having the formula:



30 W is independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂O)₂₀₋₂₀₀, (OCH₂CH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂CH₂O)₂₀₋₂₀₀, and (aa)_t;

35 aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R¹⁰,
C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁰, and a 5-10
membered heterocyclic ring system containing 1-4
heteroatoms independently selected from N, S, and O and
5 substituted with 0-3 R¹⁰;

R⁶, R^{6a}, R⁷, R^{7a}, R⁸, R^{8a}, R⁹ and R^{9a} are independently selected
at each occurrence from the group: H, =O, COOH, SO₃H,
PO₃H, C₁₋₅ alkyl substituted with 0-3 R¹⁰, aryl
10 substituted with 0-3 R¹⁰, benzyl substituted with 0-3
R¹⁰, and C₁₋₅ alkoxy substituted with 0-3 R¹⁰,
NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and a
bond to S_f;

15 R¹⁰ is independently selected at each occurrence from the
group: a bond to S_f, COOR¹¹, OH, NHR¹¹, SO₃H, PO₃H, aryl
substituted with 0-3 R¹¹, C₁₋₅ alkyl substituted with 0-1
R¹², C₁₋₅ alkoxy substituted with 0-1 R¹², and a 5-10
membered heterocyclic ring system containing 1-4
20 heteroatoms independently selected from N, S, and O and
substituted with 0-3 R¹¹;

R¹¹ is independently selected at each occurrence from the
group: H, aryl substituted with 0-1 R¹², a 5-10 membered
25 heterocyclic ring system containing 1-4 heteroatoms
independently selected from N, S, and O and substituted
with 0-1 R¹², C₃₋₁₀ cycloalkyl substituted with 0-1 R¹²,
amino acid substituted with 0-1 R¹², and a bond to S_f;

30 R¹² is a bond to S_f;

k is selected from 0, 1, and 2;
h is selected from 0, 1, and 2;
h' is selected from 0, 1, 2, 3, 4, and 5;
35 h'' is selected from 0, 1, 2, 3, 4, and 5;
g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
g'' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;


g'' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

and a pharmaceutically acceptable salt thereof.


5

41. A compound according to Claim 40, wherein the compound is of the formula:



10

wherein, Q is a cyclic pentapeptide independently selected from the group:



15

K is an L-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine, δ -N-2-imidazolinylnornithine, δ -N-benzylcarbamoylnornithine, and β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

20

K' is a D-amino acid independently selected at each occurrence from the group: arginine, citrulline, N-methylarginine, lysine, homolysine, 2-aminoethylcysteine, δ -N-2-imidazolinylnornithine, δ -N-benzylcarbamoylnornithine, and β -2-benzimidazolylacetyl-1,2-diaminopropionic acid;

25

L is independently selected at each occurrence from the group: glycine, L-alanine, and D-alanine;

30

M is L-aspartic acid;

35 M' is D-aspartic acid;

R¹ is an amino acid substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, L-valine, D-valine, alanine, leucine,
5 isoleucine, norleucine, 2-aminobutyric acid,
2-aminohexanoic acid, tyrosine, phenylalanine,
thienylalanine, phenylglycine, cyclohexylalanine,
homophenylalanine, 1-naphthylalanine, lysine, serine,
ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
10 acid, cysteine, penicillamine, and methionine;

R² is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, valine, alanine, leucine, isoleucine,
15 norleucine, 2-aminobutyric acid, 2-aminohexanoic acid,
tyrosine, L-phenylalanine, D-phenylalanine,
thienylalanine, phenylglycine, biphenylglycine,
cyclohexylalanine, homophenylalanine,
L-1-naphthylalanine, D-1-naphthylalanine, lysine, serine,
20 ornithine, 1,2-diaminobutyric acid, 1,2-diaminopropionic
acid, cysteine, penicillamine, methionine, and
2-aminothiazole-4-acetic acid;

R³ is an amino acid, substituted with 0-1 bonds to L_n,
25 independently selected at each occurrence from the group:
glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic
acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
D-phenylglycine, D-cyclohexylalanine,
30 D-homophenylalanine, D-1-naphthylalanine, D-lysine,
D-serine, D-ornithine, D-1,2-diaminobutyric acid,
D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
and D-methionine;

35 R⁴ is an amino acid, substituted with 0-1 bonds to L_n,
independently selected at each occurrence from the group:
glycine, D-valine, D-alanine, D-leucine, D-isoleucine,
D-norleucine, D-2-aminobutyric acid, D-2-aminohexanoic

acid, D-tyrosine, D-phenylalanine, D-thienylalanine,
 D-phenylglycine, D-cyclohexylalanine,
 D-homophenylalanine, D-1-naphthylalanine, D-lysine,
 D-serine, D-ornithine, D-1,2-diaminobutyric acid,
 5 D-1,2-diaminopropionic acid, D-cysteine, D-penicillamine,
 D-methionine, and 2-aminothiazole-4-acetic acid;

R⁵ is an amino acid, substituted with 0-1 bonds to L_n,
 independently selected at each occurrence from the group:
 10 glycine, L-valine, L-alanine, L-leucine, L-isoleucine,
 L-norleucine, L-2-aminobutyric acid, L-2-aminohexanoic
 acid, L-tyrosine, L-phenylalanine, L-thienylalanine,
 L-phenylglycine, L-cyclohexylalanine,
 L-homophenylalanine, L-1-naphthylalanine, L-lysine,
 15 L-serine, L-ornithine, L-1,2-diaminobutyric acid,
 L-1,2-diaminopropionic acid, L-cysteine, L-penicillamine,
 L-methionine, and 2-aminothiazole-4-acetic acid;

provided that one of R¹, R², R³, R⁴, and R⁵ in each Q is
 20 substituted with a bond to L_n, further provided that when
 R² is 2-aminothiazole-4-acetic acid, K is
 N-methylarginine, further provided that when R⁴ is
 2-aminothiazole-4-acetic acid, K and K' are
 N-methylarginine, and still further provided that when R⁵
 25 is 2-aminothiazole-4-acetic acid, K' is N-methylarginine;

S_f is a surfactant which is a lipid or a compound of the

formula: $A^9-E^1-A^{10}$;

30 A⁹ is OR²⁷;

A¹⁰ is OR²⁷;

R²⁷ is C(=O)C₁₋₁₅ alkyl;

35


E¹ is C₁₋₄ alkylene substituted with 1-3 R²⁸;

R²⁸ is independently selected at each occurrence from the group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹, -CH₂OR²⁹, -OR²⁹, and C₁-C₅ alkyl;

5 R²⁹ is independently selected at each occurrence from the group: R³⁰, H, C₁-C₆ alkyl, phenyl, and benzyl;

R³⁰ is a bond to L_n;

10 L_n is a linking group having the formula:



W is independently selected at each occurrence from the group:
 15 O, S, NH, NHC(=O), C(=O)NH, C(=O), C(=O)O, OC(=O),
 NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂O)₂₀₋₂₀₀,
 200, (OCH₂CH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂CH₂O)₂₀₋₂₀₀, and (aa)_t;

aa is independently at each occurrence an amino acid;

20

Z is selected from the group: aryl substituted with 0-3 R¹⁰,
 C₃-10 cycloalkyl substituted with 0-3 R¹⁰, and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 25 substituted with 0-3 R¹⁰;

R⁶, R^{6a}, R⁷, R^{7a}, R⁸, R^{8a}, R⁹ and R^{9a} are independently selected
 at each occurrence from the group: H, =O, C₁-C₅ alkyl
 substituted with 0-3 R¹⁰, and C₁-C₅ alkoxy substituted
 30 with 0-3 R¹⁰, and a bond to S_f;

R¹⁰ is independently selected at each occurrence from the
 group: a bond to S_f, COOR¹¹, OH, NHR¹¹, C₁-5 alkyl
 substituted with 0-1 R¹², and C₁-5 alkoxy substituted
 35 with 0-1 R¹²;

R¹¹ is independently selected at each occurrence from the
 group: H, aryl substituted with 0-1 R¹², C₃-10 cycloalkyl

substituted with 0-1 R¹², amino acid substituted with 0-1 R¹², and a bond to S_f;

R¹² is a bond to S_f;

- 5
k is selected from 0, 1, and 2;
h is selected from 0, 1, and 2;
h' is selected from 0, 1, 2, 3, 4, and 5;
h'' is selected from 0, 1, 2, 3, 4, and 5;
10 g is selected from 0, 1, 2, 3, 4, and 5;
g' is selected from 0, 1, 2, 3, 4, and 5;
g'' is selected from 0, 1, 2, 3, 4, and 5;
g''' is selected from 0, 1, 2, 3, 4, and 5;
s is selected from 0, 1, 2, 3, 4, and 5;
15 s' is selected from 0, 1, 2, 3, 4, and 5;
s'' is selected from 0, 1, 2, 3, 4, and 5;
t is selected from 0, 1, 2, 3, 4, and 5;
t' is selected from 0, 1, 2, 3, 4, and 5;
20 and a pharmaceutically acceptable salt thereof.

42. A compound according to Claim 41, wherein the present invention provides a compound selected from the group:

- 25
1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-(cyclo(Arg-Gly-Asp-D-Phe-Lys))-dodecane-1,12-dione;

1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-((ω-
30 amino-PEG₃₄₀₀-α-carbonyl)-cyclo(Arg-Gly-Asp-D-Phe-Lys))-dodecane-1,12-dione; and,

1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)-12-((ω-
amino-PEG₃₄₀₀-α-carbonyl)-Glu-(cyclo(Arg-Gly-Asp-D-Phe-
35 Lys)))₂-Dodecane-1,12-dione.

43. An ultrasound contrast agent composition,
comprising:

- (a) a compound of Claim 40, comprising: a cyclic
pentapeptide that binds to the integrin $\alpha_v\beta_3$, a surfactant and
5 a linking group between the cyclicpentapeptide and the
surfactant;
(b) a parenterally acceptable carrier; and,
(c) an echogenic gas.

10

44. An ultrasound contrast agent composition, further
comprising: 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid,
1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-
(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-
15 sn-glycero-3-phosphatidylethanolamine.

20

45. An ultrasound contrast agent composition, wherein,
the echogenic gas is a C₂₋₅ perfluorocarbon.

25

46. A composition of Claim 40 which is for use in
imaging cancer with sonography.

30

47. A composition of Claim 40 which is for use in
imaging formation of new blood vessels.

35

48. A therapeutic radiopharmaceutical composition,
comprising:

- (a) a therapeutic radiopharmaceutical of Claim 11; and,
(b) a parenterally acceptable carrier.

49. A diagnostic radiopharmaceutical composition,
comprising:

- (a) a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11; and,
(b) a parenterally acceptable carrier.

5

50. A therapeutic radiopharmaceutical composition, comprising: a radiolabelled targeting moiety, wherein the targeting moiety is a compound Q of Claim 3 and the radiolabel is a therapeutic isotope selected from the group: ^{35}S , ^{32}P ,
10 ^{125}I , ^{131}I , and ^{211}At .

51. A therapeutic radiopharmaceutical composition, comprising: a radiolabelled targeting moiety, wherein the
15 targeting moiety is a compound Q of Claim 5 and the radiolabel is a therapeutic isotope which is ^{131}I .

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 7/64, 7/56, A61K 49/00, 51/08	A3	(11) International Publication Number: WO 99/58162 (43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/US99/06826 (22) International Filing Date: 29 March 1999 (29.03.99) (30) Priority Data: 60/080,150 31 March 1998 (31.03.98) US 60/112,715 18 December 1998 (18.12.98) US (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; 974 Centre Road, WR-1ST18, Wilmington, DE 19807 (US). (72) Inventors: RAJOPADHYE, Miland; 21 Honeysuckle Road, Westford, MA 01886 (US). EDWARDS, D., Scott; 123 Farms Drive, Burlington, MA (US). HARRIS, Thomas, D.; 56 Zion Hill Road, Samel, NH 03079 (US). HEMINWAY, Stuart, J.; 21 Fidler Terrace, Lowell, MA 01850 (US). LIU, Shuang; 17 Judith Road, Chelmsford, MA 01824 (US). SINGH, Prahlad, R.; 144 Charlton Street, Arlington, MA 02174 (US). (74) Agent: VANCE, David, H.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).	(81) Designated States: AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SI, SK, UA, VN, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 6 April 2000 (06.04.00)	
(54) Title: PHARMACEUTICALS FOR THE IMAGING OF ANGIOGENIC DISORDERS (57) Abstract <p>The present invention describes novel compounds of the formula: $(Q)_t-L_n-C_h$, useful for the diagnosis and treatment of cancer, methods of imaging tumors in a patient, and methods of treating cancer in a patient. The present invention also provides novel compounds useful for monitoring therapeutic angiogenesis treatment and destruction of new angiogenic vasculature. The pharmaceuticals are comprised of a targeting moiety that binds to a receptor that is upregulated during angiogenesis, an optional linking group, and a therapeutically effective radioisotope or diagnostically effective imageable moiety. The imageable moiety is a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, an X-ray contrast agent, or an ultrasound contrast agent.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/06826

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07K7/64 C07K7/56 A61K49/00 A61K51/08		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 16474 A (BRACCO RESEARCH) 9 May 1997 (1997-05-09) the whole document ---	1-51
T	WO 98 16256 A (THE BURNHAM INSTITUTE) 23 April 1998 (1998-04-23) the whole document ---	1-51
T	WO 99 13329 A (THE BURNHAM INSTITUTE) 18 March 1999 (1999-03-18) the whole document ---	1-51
A	WO 99 40947 A (RESOLUTION PHARMACEUTICALS) 19 August 1999 (1999-08-19) the whole document ---	1-51
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 17 February 2000		Date of mailing of the international search report 25/02/2000
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Masturzo, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/06826

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	DE 197 25 368 A (MERCK PATENT CO.) 17 December 1998 (1998-12-17) the whole document ----	1-6
P, X	WO 98 14220 A (DU PONT MERCK) 9 April 1998 (1998-04-09) the whole document -----	1-51

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 06826

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
Claims: 1-5, 7-15, 17, 9-21, 23, 25, 27-29, 31-41, 43-51

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99 06826

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-5, 7-15, 17, 19-21, 23, 25, 27-29, 31-41, 43-51

Present claims 1-5, 7-15, 17, 19-21, 23, 25, 27-29, 31-41, 43-51 relate to an extremely large number of possible compounds/products. In fact, the claims contain so many options and variables that a lack of clarity (and/or conciseness) within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear (and/or concise), namely the compounds mentioned in claims 6, 16, 18, 22, 24, 26, 30 and 42. Moreover, claims 44-45 and 48-51, which do not indicate to be dependent on any previous claims, were searched as if they were dependent on claim 42.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/US 99/06826

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9716474 A	09-05-1997	EP 0800545 A JP 10512617 T US 5910300 A	15-10-1997 02-12-1998 08-06-1999
WO 9816256 A	23-04-1998	AU 4753997 A	11-05-1998
WO 9913329 A	18-03-1999	AU 9477398 A	29-03-1999
WO 9940947 A	19-08-1999	AU 2406699 A	30-08-1999
DE 19725368 A	17-12-1998	NONE	
WO 9814220 A	09-04-1998	AU 4892597 A US 5888970 A	24-04-1998 30-03-1999